



**Cochrane**  
**Library**

**Cochrane** Database of Systematic Reviews

## **Decision aids for people facing health treatment or screening decisions (Review)**

Stacey D, Légaré F, Lewis K, Barry MJ, Bennett CL, Eden KB, Holmes-Rovner M, Llewellyn-Thomas H, Lyddiatt A, Thomson R, Trevena L

Stacey D, Légaré F, Lewis K, Barry MJ, Bennett CL, Eden KB, Holmes-Rovner M, Llewellyn-Thomas H, Lyddiatt A, Thomson R, Trevena L.

Decision aids for people facing health treatment or screening decisions.

*Cochrane Database of Systematic Reviews* 2017, Issue 4. Art. No.: CD001431.

DOI: 10.1002/14651858.CD001431.pub5.

**[www.cochranelibrary.com](http://www.cochranelibrary.com)**

## TABLE OF CONTENTS

HEADER . . . . .	1
ABSTRACT . . . . .	1
PLAIN LANGUAGE SUMMARY . . . . .	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON . . . . .	4
BACKGROUND . . . . .	7
OBJECTIVES . . . . .	8
METHODS . . . . .	8
Figure 1. . . . .	10
RESULTS . . . . .	12
Figure 2. . . . .	14
Figure 3. . . . .	15
Figure 4. . . . .	17
Figure 5. . . . .	19
Figure 6. . . . .	20
DISCUSSION . . . . .	28
AUTHORS' CONCLUSIONS . . . . .	31
ACKNOWLEDGEMENTS . . . . .	31
REFERENCES . . . . .	32
CHARACTERISTICS OF STUDIES . . . . .	60
DATA AND ANALYSES . . . . .	206
ADDITIONAL TABLES . . . . .	210

# Decision aids for people facing health treatment or screening decisions

Dawn Stacey<sup>1,2</sup>, France Légaré<sup>3</sup>, Krystina Lewis<sup>1</sup>, Michael J Barry<sup>4</sup>, Carol L Bennett<sup>5</sup>, Karen B Eden<sup>6</sup>, Margaret Holmes-Rovner<sup>7</sup>, Hilary Llewellyn-Thomas<sup>8</sup>, Anne Lyddiatt<sup>9</sup>, Richard Thomson<sup>10</sup>, Lyndal Trevena<sup>11</sup>

<sup>1</sup>School of Nursing, University of Ottawa, Ottawa, Canada. <sup>2</sup>Centre for Practice Changing Research, Ottawa Hospital Research Institute, Ottawa, Canada. <sup>3</sup>Population Health and Optimal Health Practices Research Axis, CHU de Québec Research Center, Université Laval, Québec City, Canada. <sup>4</sup>Informed Medical Decisions Foundation, Boston, MA, USA. <sup>5</sup>Clinical Epidemiology Program, Ottawa Hospital Research Institute, Ottawa, Canada. <sup>6</sup>Department of Medical Informatics and Clinical Epidemiology, Oregon Health Sciences University, Portland, Oregon, USA. <sup>7</sup>Center for Ethics and Humanities in the Life Sciences, Michigan State University College of Human Medicine, East Lansing, Michigan, USA. <sup>8</sup>The Dartmouth Center for Health Policy & Clinical Practice, The Geisel School of Medicine at Dartmouth, Dartmouth College, Hanover, New Hampshire, USA. <sup>9</sup>No affiliation, Ingersoll, Canada. <sup>10</sup>Institute of Health and Society, Newcastle University, Newcastle upon Tyne, UK. <sup>11</sup>The University of Sydney, Sydney, Australia

Contact address: Dawn Stacey, School of Nursing, University of Ottawa, 451 Smyth Road, Ottawa, ON, Canada. [dstacey@uottawa.ca](mailto:dstacey@uottawa.ca).

**Editorial group:** Cochrane Consumers and Communication Group.

**Publication status and date:** New search for studies and content updated (conclusions changed), published in Issue 4, 2017.

**Citation:** Stacey D, Légaré F, Lewis K, Barry MJ, Bennett CL, Eden KB, Holmes-Rovner M, Llewellyn-Thomas H, Lyddiatt A, Thomson R, Trevena L. Decision aids for people facing health treatment or screening decisions. *Cochrane Database of Systematic Reviews* 2017, Issue 4. Art. No.: CD001431. DOI: 10.1002/14651858.CD001431.pub5.

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

## ABSTRACT

### Background

Decision aids are interventions that support patients by making their decisions explicit, providing information about options and associated benefits/harms, and helping clarify congruence between decisions and personal values.

### Objectives

To assess the effects of decision aids in people facing treatment or screening decisions.

### Search methods

Updated search (2012 to April 2015) in CENTRAL; MEDLINE; Embase; PsycINFO; and grey literature; includes CINAHL to September 2008.

### Selection criteria

We included published randomized controlled trials comparing decision aids to usual care and/or alternative interventions. For this update, we excluded studies comparing detailed versus simple decision aids.

### Data collection and analysis

Two reviewers independently screened citations for inclusion, extracted data, and assessed risk of bias. Primary outcomes, based on the International Patient Decision Aid Standards (IPDAS), were attributes related to the choice made and the decision-making process.

Secondary outcomes were behavioural, health, and health system effects.

We pooled results using mean differences (MDs) and risk ratios (RRs), applying a random-effects model. We conducted a subgroup analysis of studies that used the patient decision aid to prepare for the consultation and of those that used it in the consultation. We used GRADE to assess the strength of the evidence.

## Main results

We included 105 studies involving 31,043 participants. This update added 18 studies and removed 28 previously included studies comparing detailed versus simple decision aids. During the 'Risk of bias' assessment, we rated two items (selective reporting and blinding of participants/personnel) as mostly unclear due to inadequate reporting. Twelve of 105 studies were at high risk of bias.

With regard to the attributes of the choice made, decision aids increased participants' knowledge (MD 13.27/100; 95% confidence interval (CI) 11.32 to 15.23; 52 studies; N = 13,316; high-quality evidence), accuracy of risk perceptions (RR 2.10; 95% CI 1.66 to 2.66; 17 studies; N = 5096; moderate-quality evidence), and congruency between informed values and care choices (RR 2.06; 95% CI 1.46 to 2.91; 10 studies; N = 4626; low-quality evidence) compared to usual care.

Regarding attributes related to the decision-making process and compared to usual care, decision aids decreased decisional conflict related to feeling uninformed (MD -9.28/100; 95% CI -12.20 to -6.36; 27 studies; N = 5707; high-quality evidence), indecision about personal values (MD -8.81/100; 95% CI -11.99 to -5.63; 23 studies; N = 5068; high-quality evidence), and the proportion of people who were passive in decision making (RR 0.68; 95% CI 0.55 to 0.83; 16 studies; N = 3180; moderate-quality evidence).

Decision aids reduced the proportion of undecided participants and appeared to have a positive effect on patient-clinician communication. Moreover, those exposed to a decision aid were either equally or more satisfied with their decision, the decision-making process, and/or the preparation for decision making compared to usual care.

Decision aids also reduced the number of people choosing major elective invasive surgery in favour of more conservative options (RR 0.86; 95% CI 0.75 to 1.00; 18 studies; N = 3844), but this reduction reached statistical significance only after removing the study on prophylactic mastectomy for breast cancer gene carriers (RR 0.84; 95% CI 0.73 to 0.97; 17 studies; N = 3108). Compared to usual care, decision aids reduced the number of people choosing prostate-specific antigen screening (RR 0.88; 95% CI 0.80 to 0.98; 10 studies; N = 3996) and increased those choosing to start new medications for diabetes (RR 1.65; 95% CI 1.06 to 2.56; 4 studies; N = 447). For other testing and screening choices, mostly there were no differences between decision aids and usual care.

The median effect of decision aids on length of consultation was 2.6 minutes longer (24 versus 21; 7.5% increase). The costs of the decision aid group were lower in two studies and similar to usual care in four studies. People receiving decision aids do not appear to differ from those receiving usual care in terms of anxiety, general health outcomes, and condition-specific health outcomes. Studies did not report adverse events associated with the use of decision aids.

In subgroup analysis, we compared results for decision aids used in preparation for the consultation versus during the consultation, finding similar improvements in pooled analysis for knowledge and accurate risk perception. For other outcomes, we could not conduct formal subgroup analyses because there were too few studies in each subgroup.

## Authors' conclusions

Compared to usual care across a wide variety of decision contexts, people exposed to decision aids feel more knowledgeable, better informed, and clearer about their values, and they probably have a more active role in decision making and more accurate risk perceptions. There is growing evidence that decision aids may improve values-congruent choices. There are no adverse effects on health outcomes or satisfaction. New for this update is evidence indicating improved knowledge and accurate risk perceptions when decision aids are used either within or in preparation for the consultation. Further research is needed on the effects on adherence with the chosen option, cost-effectiveness, and use with lower literacy populations.

## PLAIN LANGUAGE SUMMARY

### Decision aids to help people who are facing health treatment or screening decisions

#### Review question

We reviewed the effects of decision aids on people facing health treatment or screening decisions. In this update, we added 18 new studies for a total of 105.

#### Background

Making a decision about the best treatment or screening option can be hard. People can use decision aids when there is more than one option and neither is clearly better, or when options have benefits and harms that people value differently. Decision aids may be

pamphlets, videos, or web-based tools. They state the decision, describe the options, and help people think about the options from a personal view (e.g. how important are possible benefits and harms).

### **Study characteristics**

For research published up to April 2015, there were 105 studies involving 31,043 people. The decision aids focused on 50 different decisions. The common decisions were about: surgery, screening (e.g. prostate cancer, colon cancer, prenatal), genetic testing, and medication treatments (e.g. diabetes, atrial fibrillation). The decision aids were compared to usual care that may have included general information or no intervention. In the 105 studies, 89 evaluated a patient decision aid used by people in preparation for the visit with the clinician, and 16 evaluated its use during the visit with the clinician.

### **Key results with quality of the evidence**

When people use decision aids, they improve their knowledge of the options (high-quality evidence) and feel better informed and more clear about what matters most to them (high-quality evidence). They probably have more accurate expectations of benefits and harms of options (moderate-quality evidence) and probably participate more in decision making (moderate-quality evidence). People who use decision aids may achieve decisions that are consistent with their informed values (evidence is not as strong; more research could change results). People and their clinicians were more likely to talk about the decision when using a decision aid. Decision aids have a variable effect on the option chosen, depending on the choice being considered. Decision aids do not worsen health outcomes, and people using them are not less satisfied. More research is needed to assess if people continue with the option they chose and also to assess what impact decision aids have on healthcare systems.

## SUMMARY OF FINDINGS FOR THE MAIN COMPARISON [\[Explanation\]](#)

Patient decision aids compared with usual care for adults considering treatment or screening decisions						
<b>Patient or population:</b> adults considering treatment or screening decisions <b>Settings:</b> all settings <b>Intervention:</b> patient decision aid <b>Comparison:</b> usual care						
Outcomes	Illustrative comparative benefits* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed benefit	Corresponding benefit				
	Usual care	Patient decision aid				
<b>Knowledge - all studies</b> Standardized on score from 0 (no knowledge) to 100 (perfect knowledge), soon after exposure to the decision aid	The mean knowledge score was 56.9% across control groups, ranging from 27.0% to 85.2%	The mean knowledge score in the intervention groups was 13.27 higher (11.32 to 15.23 higher)	-	13,316 (52 studies)	⊕⊕⊕⊕ <b>High</b> <sup>a,b</sup>	Higher scores indicate better knowledge. 46 out of 52 studies showed a statistically significant improvement in knowledge
<b>Accurate risk perceptions - all studies</b> Assessed soon after exposure to the decision aid	269 per 1000 <sup>c</sup>	565 per 1000 (447 to 716 per 1000)	RR 2.10 (1.66 to 2.66)	5096 (17 studies)	⊕⊕⊕○ <b>Moderate</b> <sup>a,d</sup>	-
<b>Congruence between the chosen option and informed values - all studies</b> Assessed soon after exposure to the decision aid	289 per 1000 <sup>c</sup>	595 per 1000 (422 to 841 per 1000)	RR 2.06 (1.46 to 2.91)	4626 (10 studies)	⊕⊕○○ <b>Low</b> <sup>a,d,e,f</sup>	-

<b>Decisional conflict: uninformed subscale - all studies</b> Standardized on score from 0 (not uninformed) to 100 (uninformed) Assessed soon after exposure to the decision aid	The mean for outcome 'feeling uninformed' ranged across control groups from 11.1 to 61.1 Scores ≤ 25 associated with following through on decisions. Scores > 38 associated with delay in decision making	The mean feeling uninformed in the intervention groups was 9.28 lower (12.20 to 6.36 lower)	-	5707 (27 studies)	⊕⊕⊕⊕ <b>High</b> <sup>a,b</sup>	Lower scores indicate feeling more informed
<b>Decisional conflict: unclear about personal values subscale - all studies</b> Standardized on score from 0 (not unclear) to 100 (unclear) Assessed soon after exposure to the decision aid	The mean for outcome 'feeling unclear about personal values' ranged across control groups from 15.5 to 53.2 Scores ≤ 25 associated with follow-through with decisions. Scores > 38 associated with delay in decision making	The mean feeling unclear values in the intervention groups was 8.81 lower (11.99 to 5.63 lower)	-	5068 (23 studies)	⊕⊕⊕⊕ <b>High</b> <sup>a,b</sup>	Lower scores indicate feeling clearer about values
<b>Participation in decision making: clinician-controlled decision making - all studies</b> Assessed soon after consultation with clinician	228 per 1000 <sup>c</sup>	155 per 1000 (125 to 189 per 1000)	RR 0.68 (0.55 to 0.83)	3180 (16 studies)	⊕⊕⊕○ <b>Moderate</b> <sup>a,e</sup>	Patient decision aids aim to increase patient involvement in making decisions; lower proportion of clinician-controlled decision making is better
<b>Adverse events</b>	There were no adverse effects on health outcomes or satisfaction, and no other adverse effects reported					

\*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; RR: risk ratio

GRADE Working Group grades of evidence

**High quality:** further research is very unlikely to change our confidence in the estimate of effect.

**Moderate quality:** further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

**Low quality:** further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

**Very low quality:** we are very uncertain about the estimate.

<sup>a</sup>The vast majority of studies measuring this outcome were not at high risk of bias.

<sup>b</sup>The GRADE ratings for these outcomes were not downgraded for heterogeneity given the generally consistent direction of effects across studies for the decision aid compared to usual care groups.

<sup>c</sup>The data source for the assumed risk was the mean control event rate.

<sup>d</sup>The GRADE rating was downgraded given the lack of precision.

<sup>e</sup>The GRADE rating was downgraded given the lack of consistency.

<sup>f</sup>The GRADE rating was downgraded given the lack of directness. As well, the outcome was measured using various approaches with no gold standard approach.



## BACKGROUND

Many health treatment and screening decisions have no single 'best' choice. These types of decisions are considered 'preference-sensitive' because there is insufficient evidence about outcomes or there is a need to trade off known benefits and harms. *Clinical Evidence* analyzed 3000 treatments, classifying 50% as having insufficient evidence, 24% as likely to be beneficial, 7% as requiring trade-offs between benefits and harms, 5% as unlikely to be beneficial, 3% as likely to be ineffective or harmful, and only 11% as being clearly beneficial (Clinical Evidence 2013). Not only does one have to take into account the strength of the evidence, but even for the 11% of treatments that show beneficial effects for populations, physicians need to translate the probabilistic nature of the evidence for individual patients to help them reach a decision based on informed values. Patient decision aids are an intervention that can be used to present such evidence (Brouwers 2010). This review is an update of the review last published in 2014 of the comparisons between patient decision aids and usual care (Stacey 2014b). To provide a more focused review, we removed 28 studies that compared detailed versus simple decision aids.

### Description of the intervention

According to the International Patient Decision Aids Standards (IPDAS) Collaboration (Elwyn 2006; IPDAS 2005a; Joseph-Williams 2013), decision aids are evidence-based tools designed to help patients make specific and deliberated choices among healthcare options. Patient decision aids supplement (rather than replace) clinicians' counselling about options. The specific aims of decision aids and the type of decision support they provide may vary slightly, but in general they:

1. explicitly state the decision that needs to be considered;
2. provide evidence-based information about a health condition, the options, associated benefits, harms, probabilities, and scientific uncertainties;
3. help patients to recognize the values-sensitive nature of the decision and to clarify, either implicitly or explicitly, the value they place on the benefits and harms. (To accomplish this, patient decision aids may describe the options in enough detail that clients can imagine what it is like to experience the physical, emotional, and social effects, or they may guide clients to consider which benefits and harms are most important to them.) Decision aids differ from usual health education materials. Decision aids make the decision being considered explicit, providing a detailed, specific, and personalized focus on options and outcomes for the purpose of preparing people for decision making. In contrast, health education materials help people to understand their diagnosis, treatment, and management in general terms, but given their broader perspective, these materials are not focused on decision points and thus do not necessarily help them to participate in decision making. Many decision aids are based on a con-

ceptual model or theoretical framework (Durand 2008; Mulley 1995; O'Connor 1998b; Rothert 1987).

In response to concerns about variability in the quality of patient decision aids, the IPDAS Collaboration reached agreement on criteria for judging their quality (Elwyn 2006). More than 100 researchers, clinicians, patients, and policymakers from 14 countries participated. Participants addressed three domains of quality: clinical content, development process, and evaluation of a patient decision aid's effectiveness. A series of background papers informing the original IPDAS criteria were updated in 2013 (IPDAS 2013). Subsequently, an international team of researchers reached consensus on a shorter set of qualifying and certifying criteria (Joseph-Williams 2013). Informed by IPDAS, the Washington State Health Authority launched the first programme for certifying patient decision aids in 2016 (Washington State 2016).

### How the intervention might work

Decision aids can be used before, during, or after a clinical encounter to enable patients to become active, informed participants. Providing the patient decision aid in preparation for the consultation allows people more time to digest the information and be ready to discuss the decision, but this may not be feasible in some health decisions (e.g. antibiotics for upper respiratory infections). Decision aids can also facilitate shared decision making. Shared decision making is defined as a process through which clinicians and patients make healthcare choices together (Charles 1997; Makoul 2006), representing the crux of people-centred care (Weston 2001). However, the way the clinician provides information may strongly affect people's preferences (Hibbard 1997), prompting the need for standardized information such as patient decision aids. Patients who are more active in making decisions about their health have better health outcomes and healthcare experiences (Hibbard 2013; Kiesler 2006). In summary, patient decision aids may help clinicians and patients come to quality decisions, grounded in patients' values and taking into account the potential trade-offs in benefits and risks of different options.

### Why it is important to do this review

Given the broad range of stakeholders interested in patient decision aids and the rapidly expanding field of research, there was a need to update this review to identify studies on new decisions or conducted in new countries and to strengthen the synthesized evidence supporting use of patient decision aids for outcomes that do not yet have high-quality evidence. In fact, the 2014 publication was the most cited Cochrane Review in 2015 based on 1888 reviews published in 2013 and 2014. With growing development of patient decision aids for use in the consultation, we wanted to conduct a subgroup analysis of patient decision aids used in preparation for versus within the consultation.

Results from previous reviews were used to inform clinical practice guidelines such as Patient Experience in Adult NHS Services (NCGC/NICE 2012) and Decision Support for Adults Living with Chronic Kidney Disease (RNAO 2009). Subgroup analyses of included studies have focused on anxiety (Bekker 2003), adherence (Trenaman 2016), values congruence (Munro 2016), participant trial identity (Brown 2015), and heterogeneity (Gentles 2013). Other systematic reviews have been conducted on the use of patient decision aids as one type of intervention to facilitate shared decision making in clinical practice (Coyne 2013; Duncan 2010; Elwyn 2013; Legare 2010; Legare 2014).

## OBJECTIVES

To assess the effects of decision aids in people facing treatment or screening decisions.

## METHODS

### Criteria for considering studies for this review

#### Types of studies

We included all published studies that used a randomized controlled trial (RCT) design evaluating patient decision aids.

#### Types of participants

We included studies involving adults aged 18 years or older who were making decisions about screening or treatment options for themselves, a child, or an incapacitated significant other. We excluded studies in which participants were making hypothetical choices.

#### Types of interventions

We included studies that evaluated a patient decision aid as part of the intervention. Decision aids were defined as interventions designed to help people make specific and deliberated choices among options (including the status quo), by making the decision explicit and by providing (at the minimum) information on the options and outcomes relevant to a person's health status as well as implicit methods to clarify values. The aid also may have included: information on the disease/condition; costs associated with options; probabilities of outcomes tailored to personal health risk factors; an explicit values clarification exercise; information on others' opinions; a personalized recommendation on the basis of clinical characteristics and expressed preferences; and guidance

or coaching in the steps of making and communicating decisions with others.

We excluded studies if interventions focused on: decisions about lifestyle changes, clinical trial entry, or general advance directives (e.g. do not resuscitate); education programmes not geared to a specific decision; and interventions designed to promote adherence or elicit informed consent regarding a recommended option. We also excluded studies when the relevant decision aid(s) were not available to us and not adequately described in the article(s), because we could not determine the aids' characteristics and whether or not they met the minimum criteria to qualify as patient decision aids.

#### Types of comparisons

We included studies that compared patients exposed to a patient decision aid to patients in comparison groups that were exposed to usual care, general information, clinical practice guideline, placebo intervention, or no intervention. For the purposes of this review, we refer to all such control comparisons as 'usual care'.

We excluded studies that compared two different types of patient decision aids.

#### Types of outcome measures

To ascertain whether the decision aids achieved their objectives, we examined a broad range of outcomes. Although the decision aids focused on diverse clinical decisions, many had similar objectives such as improving knowledge scores, the accuracy of risk perceptions, and participation in decision making. Many of these evaluation criteria mapped onto the International Patient Decision Aids Standards (IPDAS) criteria for evaluating the effectiveness of decision aids (Elwyn 2006; IPDAS 2005b; Sepucha 2013). The IPDAS criteria were attributes related to the choice (e.g. match between the chosen option and the features that matter most to the informed patient) and to the decision-making process (e.g. helps patients to recognize that a decision needs to be made; know the options and their features; understand that values affect the decision; be clear about the features that matter most; discuss values with their clinician; and become involved in their preferred ways). A complete list of outcomes, specified in advance of the review, included primary and secondary outcomes.

#### Primary outcomes

##### Evaluation criteria that map onto the IPDAS criteria

- Attributes of the choice made: does the patient decision aid improve the match between the chosen option and the features that matter most to the informed patient (demonstrated by outcomes such as knowledge, accurate risk perceptions, values-choice congruence)?

- Attributes of the decision-making process: does the patient decision aid help patients to recognize that a decision needs to be made, feel informed about the options and their features, be clear about the option features that matter most, discuss values with their clinician, and become involved in decision making?

### Other decision-making process variables

- Decisional conflict
- Patient-clinician communication
- Participation in decision making
- Proportion undecided
- Satisfaction with the choice, with the process of decision making, and with the preparation for decision making

### Secondary outcomes

#### Behaviour

- Choice (the actual choice implemented; if not reported, the participants' preferred option was used as a surrogate measure)
- Adherence to chosen option

#### Health outcomes

- Health status and quality of life (generic and condition-specific)
- Anxiety, depression, emotional distress, regret, confidence

#### Healthcare system

- Costs, cost-effectiveness
- Consultation length
- Litigation rates

### Search methods for identification of studies

Our search strategy for the review included:

1. searching electronic medical and social science databases; and
2. searching other resources.

### Electronic searches

For this update, we used the same search strategy that was revised by the Trials Search Coordinator at the Cochrane Consumers and Communication Group in the last update (Stacey 2014b).

Therefore, the cumulative search of electronic databases is as follows.

- Cochrane Central Register of Controlled Trials (CENTRAL; 2015, Issue 6) in the Cochrane Library (searched to 24 April 2015).
- MEDLINE Ovid (1966 to 24 April 2015).
- Embase Ovid (1980 to 24 April 2015).
- PsycINFO Ovid (1806 to 24 April 2015).
- CINAHL Ovid (1982 to September 2008), then in Ebsco (to 24 April 2015).

We present the search strategies in Appendix 1 and Appendix 2.

### Searching other resources

On 18 December 2015 we also searched trial registries (World Health Organization, ClinicalTrials.gov), the Internet using Google and Google Scholar, and the Decision Aid Library Inventory ([decisionaid.ohri.ca](http://decisionaid.ohri.ca)). Finally, reference lists of all newly included trials were searched.


### Data collection and analysis

For this current update, we focused only on new publications that had appeared since the previous publication (Stacey 2014b), and we limited the inclusion to patient decision aids versus usual care. As such, we removed studies from the previous reviews that compared detailed versus simple patient decision aids to provide a more focused review.

### Selection of studies

Pairs of eight review authors (CB, DS, RT, MB, MHR, KE, NC, DR) screened all identified citations. We retrieved the full text of any papers identified as potentially relevant by at least one author, listing all papers excluded from the review at this stage, with reasons, in the 'Characteristics of excluded studies' table. We also provided citation details and any available information about ongoing studies, and we collated and reported details of additional publications, so that each study (rather than each report) was the unit of interest. We report the screening and selection process in Figure 1.

Figure 1. Study flow diagram.



## Data extraction and management

Two research assistants extracted data independently (KL, IS). We compared findings and resolved inconsistencies through discussion with the principal investigator (DS) and, when necessary, with a co-author (CB). No review authors extracted data for their own studies in this update nor in any other versions of this review.

One review author entered all extracted data into Review Manager 5 (RevMan 5), and a second one worked independently to check for accuracy against the data extraction sheets (RevMan 2014).

## Assessment of risk of bias in included studies

Two research assistants independently appraised studies using the Cochrane 'Risk of bias' tool (current update: KL, IS) (Higgins 2011, Chapter 8). We judged each item as conferring high, low, or unclear risk of bias as set out in the criteria provided by Higgins 2011, and we provided a quote from the study report and a justification for our judgement for each item in the 'Risk of bias' table. For the item on 'other' potential sources of bias, the assessment included: whether the same clinician provided consultation to both the intervention and usual care groups with measures taken post-consultation, whether clustering was accounted for in the analysis; and potential sources of bias reported by the authors in the study limitations.

We resolved inconsistencies by discussion with the principal investigator (DS) and, when necessary, with a co-author (CB). No review authors appraised risk of bias for their own studies in this update nor in any other versions of this review.

Studies were deemed to be at the highest risk of bias if they were scored as at high risk on any of the items of the risk of bias tool (Higgins 2011).

## Measures of treatment effect

For dichotomous outcomes, we analyzed data based on the number of events and the number of people assessed in the intervention and comparison groups. We will use these to calculate the risk ratio (RR) and 95% confidence interval (CI). For continuous measures, we analyzed data based on the mean, standard deviation (SD) and number of people assessed for both the intervention and comparison groups to calculate mean difference (MD) and 95% CI.

First, we described study characteristics individually. The a priori comparison was usual care versus decision aids. For studies in which there were more than one intervention group, we extracted data from the groups that provided the strongest contrast between the intervention and control groups. We pooled results across studies in cases where investigators used similar outcome measures and

the effects were expected to be independent of the type of decision studied. For example, we expected decision aids to improve knowledge and create accurate perceptions of options, benefits, and harms; to reduce decisional conflict; and to enhance active participation in decision making. Therefore, we pooled data from included RCTs for these outcomes if trials used comparable measures. To facilitate pooling of data for some outcomes (e.g. knowledge, decisional conflict), we standardized the scores to range from 0 to 100 points. When analysing the effects of decision aids on choices, we pooled outcomes on more homogeneous subgroups of decisions (choice of major surgery versus conservative options; screening test or not, etc.).

## Unit of analysis issues

We checked for unit-of-analysis errors. Where we found errors and sufficient information was available, we re-analyzed the data using the appropriate unit of analysis by taking account of the intraclass correlation (ICC). We obtained estimates of the ICC by contacting authors of included studies, or we imputed them using estimates from external sources. For two studies (Kupke 2013; Lewis 2010), it was not possible to obtain sufficient information to re-analyze the data, and we reported these studies as being at high risk for 'other' bias based on these unit-of-analysis errors. We made no adjustments to the data based on these two studies that were included in meta-analysis for knowledge only.

## Dealing with missing data

We contacted authors to obtain missing data. Where possible, we conducted analysis on an intention-to-treat basis; otherwise, we analyzed data as reported. We reported on the levels of loss to follow-up and assessed this as a source of potential bias.

## Assessment of heterogeneity

For this update and in previous versions of the review, we grouped studies together across populations and settings. The aim was to enable an assessment of the effectiveness of decision aids across conditions, rather than to focus on disease-specific contexts. Given that decision aids are a well-defined and clearly delineated type of intervention, we decided that this approach was defensible. On the basis of grouping studies across populations and decision aid elements, we anticipated that there would be a substantial degree of heterogeneity in our pooled effect estimates. However, we decided that we would consider the direction of effects and variability in these rather than variability in the size of effects, as the major basis for our interpretation of heterogeneity. This meant that for those pooled effect estimates where the direction of effect was

consistent across studies, we did not downgrade for inconsistency, despite some variability in the size of effects across individual studies. We did downgrade for inconsistency for one outcome: congruence between the chosen option and informed values. This was because there is no accepted gold standard measure for assessing this outcome, and we considered that variability in measurement by the included studies added further uncertainty about the effects of decision aids for this outcome.

Where heterogeneity was present in pooled effect estimates, we explored possible reasons for variability by conducting subgroup analysis in the 2009 update (O'Connor 2009b). The post hoc analysis included the IPDAS effectiveness criteria to explore heterogeneity according to the following factors: the type of decision (treatment versus screening), the type of media of the decision aid (video/computer versus audio booklet/pamphlet), and the possibility of a ceiling effect based on usual-care scores (resulting in the removal of studies with lower scores for knowledge and accurate risk perception and higher scores for decisional conflict using the subscales measuring levels of informedness and clarity of values). We analyzed the effect of removing the biggest outlier(s) (defined by visual inspection of forest plots). Given that the post hoc analysis did not alter the findings from the 2009 update, we did not re-conduct the post hoc analysis for the IPDAS effectiveness criteria.

### Assessment of reporting biases

We used funnel plots to assess publication bias.

### Data synthesis

We used RevMan 5 software to estimate a weighted intervention effect with 95% confidence intervals (RevMan 2014). For continuous measures, we used mean differences (MD); for dichotomous outcomes, we calculated pooled relative risks (RR). We analyzed all data with a random-effects model because of the diverse nature of the studies being combined and then anticipated variability in the populations and interventions of the included studies. We summarized all of the results for the primary outcomes and rated the strength of evidence using GRADE (Andrews 2013), presenting these in a 'Summary of findings' table (Higgins 2011).

### Subgroup analysis and investigation of heterogeneity

For this update, we conducted a subgroup analysis to compare the effects of the intervention when used in preparation for the consultation with the effects of those used during the consultation to usual care.

### Sensitivity analysis

We performed post hoc sensitivity analyses to examine the effect of excluding studies of lower methodological quality. The analysis

excluded studies that were at high risk of bias for any of the categories in the 'Risk of bias' assessment (Higgins 2011).

### 'Summary of findings' table

We prepared a 'Summary of findings' table to present the results of meta-analysis, based on the methods described in Chapter 11 of the *Cochrane Handbook for Systematic Reviews of Interventions* (Schünemann 2011). We presented the results of meta-analysis for the major comparison of the review for each of the key outcomes. We provided a source and rationale for each assumed risk cited in the table and used the GRADE criteria to rank the quality of the evidence for each outcome on each of the following domains: risk of bias, inconsistency, imprecision, indirectness, and publication bias. Two authors independently assessed the quality of the evidence using the GRADEprofiler (GRADEpro) software (GRADEpro GDT).

## RESULTS

### Description of studies

The current version of our review updates our 2014 version, Stacey 2014b, with 18 new studies (Bozic 2013; Brazell 2014; Chabrera 2015; Fraenkel 2012; Knops 2014; Köpke 2014; Kuppermann 2014; Lam 2013; LeBlanc 2015; Legare 2012; Lepore 2012; Mathers 2012; Mott 2014; Sawka 2012; Shourie 2013; Stacey 2014a; Taylor 2006; Williams 2013). For this update, we excluded 28 previously included studies due to the comparisons being limited to detailed versus simple patient decision aids (Deschamps 2004; Deyo 2000; Dodin 2001; Goel 2001; Green 2004; Hunter 2005; Kuppermann 2009; Labrecque 2010; Lalonde 2006; Legare 2003; Leung 2004; Myers 2005a; Myers 2011; O'Connor 1998a; O'Connor 1999a; Raynes-Greenow 2010; Rostom 2002; Rothert 1997; Schapira 2000; Schapira 2007; Solberg 2010; Street 1995; Tiller 2006; Van Roosmalen 2004; Volk 2008; Wakefield 2008a; Wakefield 2008b; Wakefield 2008c).

### Results of the search

In total, we identified 46,054 citations from the electronic database searches and 258 citations from other sources. Of these, we assessed 504 citations for eligibility using the full text (see Figure 1).

### Included studies

The remaining 151 citations provided data on 105 studies that met our inclusion criteria, 18 of which are new for this update. The 105 RCTs, involving 31,043 participants, presented results from 10 countries: Australia (10 studies), Canada (15 studies), China (1 study), Finland (2 studies), Germany (6 studies), Netherlands (2



studies), Spain (1 study), Sweden (1 study), the UK (16 studies), the USA (50 studies), and Australia plus Canada (1 study). We present study details below and in the [Characteristics of included studies](#) table.

### Unit of randomization

Ninety studies randomized individual patients, and 15 randomized clusters. For cluster trials, [Allen 2010](#) randomized 12 company worksites; [Fraenkel 2012](#), 2 groups of primary care physicians; [Hamann 2006](#), 12 inpatient psychiatric units; [Kupke 2013](#), 49 dental students; [Legare 2011](#), 4 family medicine group practices; [Legare 2012](#), 12 family medicine group practices; [Lewis 2010](#), 32 family medicine group practices; [Loh 2007](#), 30 general practitioners; [Mathers 2012](#), 49 general medicine practices; [McAlister 2005](#), 102 primary care practices; [Mullan 2009](#), 40 clinicians; [Nagle 2008](#), 60 general practitioners; [Shourie 2013](#), 50 general medicine practices; [Weymiller 2007](#), 21 endocrinologists; and [Whelan 2004](#), 27 surgeons.

For 10 studies ([Allen 2010](#); [Legare 2011](#); [Legare 2012](#); [Loh 2007](#); [Mathers 2012](#); [Mullan 2009](#); [Nagle 2008](#); [Shourie 2013](#); [Weymiller 2007](#); [Whelan 2004](#)), the cluster effect was taken into account in the published outcome data, and the meta-analysis used published results. Although [Hamann 2006](#) did not account for the cluster effect in the published outcome data, the way this study was reported did not allow us to include it in the meta-analysis, so we did not re-analyze the data and report the study separately. For [McAlister 2005](#), meta-analysis was done applying the design effect (based on the published intracluster correlation coefficient (ICC)). For [Fraenkel 2012](#), the authors stated that adding a random effect for physician clusters did not contribute to better-fitting regression models, and we removed it from the analysis. The analysis by [Kupke 2013](#) and [Lewis 2010](#) did not account for clustering.

### Decision aids and comparisons

The 105 included studies evaluated decision aids that focused on 50 different decisions ([Table 1](#)). The most common decisions were about prostate cancer screening (14 studies), colon cancer screening (10 studies), medication for diabetes (4 studies), breast cancer genetic testing (4 studies), prenatal screening (4 studies), medication for atrial fibrillation (4 studies), and surgery (mastectomy for breast cancer, 4 studies; hysterectomy, 3 studies; prostate cancer treatment, 4 studies). New decision topics added in this update included surgery for prolapsed pelvic organs (1 study) and asymptomatic aortic abdominal aneurysm (1 study); restoration

for tooth decay (1 study); measles, mumps, and rubella vaccine for infants (1 study); treatment of post-traumatic stress disorder (1 study); and radioactive iodine treatment for thyroid cancer (1 study).

The decision aids used different formats and were compared to a variety of control interventions (e.g. usual care, general information, no intervention, guideline, placebo intervention). We noted the nature of usual care when reported (see [Characteristics of included studies](#) table). For this review, we have grouped control interventions and refer to them as 'usual care'.

According to the definition of a patient decision aid, all of the studies evaluated patient decision aids that included information about the options and outcomes and provided at least implicit clarification of values. Most patient decision aids included information on the clinical problem (90.5%) as well as outcome probabilities (89.5%). Fewer patient decision aids provided guidance in the steps of decision making (65.7%), explicit methods to clarify values (57.1%), and/or examples of others' experiences (41.0%) (see [table Characteristics of included studies](#)).

### Excluded studies


We excluded 302 studies upon close perusal of the relevant papers (see [Characteristics of excluded studies](#)). The reasons for exclusion were: the study was not a randomized controlled trial; the decision was hypothetical, with participants not actually at a point of decision making; the intervention was not focused on making a choice; the intervention offered no decision support in the form of a decision aid or did not provide enough information about the decision aid; no comparison outcome data were provided; the study did not evaluate the decision aid; the study was a protocol; the decision aid was about clinical trial entry, lifestyle choice, or advanced care planning; the study involved testing the presentation of the decision aid, but with no difference in the content of the decision aid between study groups; or the study compared a detailed versus simple decision aid.

We also identified 61 ongoing RCTs through trial registration databases, personal contact, and published protocols in the electronic database searches (see references to [Ongoing studies](#) and [table Characteristics of ongoing studies](#)).

### Risk of bias in included studies


Details on the ratings and rationale for risk of bias are in the [Characteristics of included studies](#) table and displayed in [Figure 2](#) and [Figure 3](#).

**Figure 2. Risk of bias summary as percentages across all included studies.**





**Figure 3. Risk of bias summary for each included study.**



## Allocation

When assessing risk of selection bias, we rated all 105 studies as being at low or unclear risk of bias. Allocation concealment methods prompted a rating of low or unclear risk of bias in 104 studies and high risk of bias in 1 study (Kupke 2013).

## Blinding

We judged 102 studies to be at low or unclear risk of performance and detection bias for the blinding of participants and personnel, while 3 (2.9%) studies were at high risk of bias. High risk of bias was due to lack of blinding of physicians to the status of patients randomized to the patient decision aid and alternative interventions (Auvinen 2004; Krist 2007; Man-Son-Hing 1999). We rated the blinding of outcome assessment as leading to low or unclear risk of bias in all 105 studies.

## Incomplete outcome data

For 103 studies, aspects related to incomplete outcome data conferred low or unclear risk of bias. In two (1.9%) studies (Chambers 2012; Mott 2014), there was high risk of bias due to high attrition rates.

## Selective reporting

We rated all 105 studies as being at either low risk of bias because the protocol was registered publicly or at unclear risk of bias because we could not assess the extent or the impact of any reporting bias.

## Other potential sources of bias

Of 105 studies, we rated 98 as being at low or unclear risk of other potential sources of bias. The other seven (6.7%) discussed other potential risks of bias (Brazell 2014; Clancy 1988; Hamann 2006; Knops 2014; Kupke 2013; LeBlanc 2015; Lewis 2010). We rated Brazell 2014 and LeBlanc 2015 as being at high risk of bias given that the same physicians provided consultation to both intervention and control groups, and measures were taken after physician consultation. Clancy 1988 describes a potential for selection bias because non-randomized medical residents were added to the decision aid group, and there was a low response rate among those offered decision aid. We rated Knops 2014 as being at

high risk of bias given that a large number of potential participants did not participate in the study. Hamann 2006, Kupke 2013, and Lewis 2010 did not account for clustering in their analyses.

## Effects of interventions

See: [Summary of findings for the main comparison](#)

In addition to [Summary of findings for the main comparison](#), see the [Data and analyses](#) figures for pooled data and Additional tables for outcome data that we did not pool. This section presents the attributes of the choice made, the attributes of the decision process, and secondary outcomes.

## Primary outcomes


### Attributes of the choice made: does the patient decision aid improve the match between the chosen option and the features that matter most to the informed patient?

The randomized controlled trials used three measures that correspond to this outcome: knowledge scores, accuracy of risk perceptions, and congruence between the chosen option and the patient's values.

## Knowledge

Seventy-one of the 105 studies (67.6%) assessed the effects of decision aids on knowledge. The studies' knowledge tests were based on information contained in the decision aid. The proportion of accurate responses was transformed to a percentage scale ranging from 0% (no correct responses) to 100% (fully correct responses). There is high-quality evidence that patient decision aids were more effective than usual care (52 studies) on knowledge scores (MD 13.27, 95% CI 11.32 to 15.23; Analysis 1.1). In absolute terms the group receiving usual care had, on average, 57 of 100 answers correct. Those in the decision aid group scored better, with 70 of 100 answers correct on average (from 68 to 72 correct). Nineteen additional studies presented knowledge scores that could not be included in the pooled outcome (see [Table 2](#)). Most of these other studies reported statistically-significantly higher knowledge scores for those exposed to the decision aid compared to usual care. The funnel plot for knowledge as an outcome in studies comparing decision aid to usual care shows that these studies are at low risk for publication bias ([Figure 4](#)).

**Figure 4. Funnel plot of comparison: I Knowledge, outcome: I.I Knowledge - all studies.**



#### **Accurate risk perceptions (i.e. perceived probabilities of outcomes)**

Of 105 studies, 25 (23.8%) examined the effects of patient decision aids on the accuracy of patients' perceived probabilities of outcomes (see Analysis 2.1; [Table 3](#)). We classified the accuracy of perceived outcome probabilities according to the percentage of individuals whose judgments corresponded to the scientific evidence about the chances of an outcome for similar people. For studies that elicited risk perceptions using multiple items, we averaged the proportion of accurate risk perceptions.

There is moderate-quality evidence that patient decision aids were more effective than usual care for transmitting accurate risk perceptions (risk ratio (RR) 2.10, 95% CI 1.66 to 2.66, 17 studies; Analysis 2.1). This means that for every 1000 people receiving usual care, 269 were likely to accurately interpret risk, whereas far more people (565 people per 1000; from 447 to 716) accurately interpreted risk after using a decision aid.

Eight studies reported results that were not amenable to pooling (see [Table 3](#)). [Fraenkel 2012](#); [Hanson 2011](#); [Kuppermann 2014](#); [Mathieu 2010](#); and [Smith 2010](#) reported a statistically significant improvement in accurate perceptions of outcomes for the decision aid group compared to usual care, and [Miller 2005](#) reported no effect on risk perception. In another study, [Weymiller 2007](#) reported

participants allocated to the decision aid had a significantly more accurate perception of their estimated cardiovascular risk without statin therapy compared to the usual care group; this effect was greater when the clinician used the decision aid during the consultation rather than when the researcher used the decision aid in preparation for the consultation ( $P_{interaction} = 0.03$ ). For the final study by [Mann E 2010](#), three of eight knowledge test items measured accurate risk perceptions, but results were presented for total knowledge and not individual items. The funnel plot for accurate risk perception as an outcome in studies comparing decision aid to usual care shows low risk for publication bias.

#### **Congruence between chosen option and values**

Of 105 studies, 16 (15.3%) measured congruence between the chosen options and the patients' values. Six measured values-choice congruence without considering knowledge ([Arterburn 2011](#); [Berry 2013](#); [Frosch 2008a](#); [Legare 2008a](#); [Lerman 1997](#); [Vandemheen 2009](#)). Of 10 studies that measured informed values-choice congruence, eight used the Multi-Dimensional Measure of Informed Choice ([Bjorklund 2012](#); [Fagerlin 2011](#); [Mathieu 2007](#); [Mathieu 2010](#); [Nagle 2008](#); [Smith 2010](#); [Steckelberg 2011](#); [Trevena 2008](#)), which assesses the extent to which the choice is

based on relevant knowledge, is consistent with a person's values/attitudes, and is behaviourally implemented (Michie 2002). These studies operationalized the measure in terms of knowledge scores higher than the mid-point of the scale, attitude scale scores higher than the mid-point, and choice being congruent with attitude. Two other studies measured informed values-based choice: Schwalm 2012 assessed the extent to which the choice was based on knowledge score  $\geq 60\%$  and a score for three values-importance ratings that matched the choice; and Stacey 2014a assessed the extent to which the choice was based on knowledge score  $\geq 66\%$  and measured values-choice congruence using a logistic regression model. For the 10 studies that measured informed values-choice congruence, two used preferred choice (Mathieu 2010; Trevena 2008), and the other eight used actual choice.

There is low quality evidence that patient decision aids were more effective than usual care for selecting an option that was congruent with their informed values (RR 2.06, 95% CI 1.46 to 2.91, 10 studies; Analysis 3.1). Of the 10 studies, 8 individually showed statistically higher congruence scores for the patient decision aid compared to usual care, and 2 showed no difference (Bjorklund 2012; Mathieu 2010). Repeating this analysis using the studies that measured actual choice and not preferred choice revealed a pooled RR of 2.13 (95% CI 1.44 to 3.14; 8 studies). A sub-analysis of studies using the Multi-Dimensional Measure of Informed Choice revealed a pooled RR of 2.08 (95% CI 1.40 to 3.08, 8 studies; Analysis 3.3).

There was no difference between patient decision aid and usual care for the six studies that measured values-choice congruence without considering knowledge scores (Arterburn 2011; Berry 2013; Frosch 2008a; Legare 2008a; Lerman 1997; Vandemheen 2009; see Table 4). We did not pool these studies because of how they reported results. Arterburn 2011 reported that, compared to the control group, those exposed to the decision aid experienced a more rapid early improvement of value-choice concordance immediately after exposure. Legare 2008a reported that women's valuing of the non-chemical aspect of natural health products was positively associated with their choice of natural health products in managing menopausal symptoms ( $P = 0.006$ ). The other four studies reported no differences between groups. However, Frosch 2008a observed that men exposed to the decision aid who chose not to have a prostate-specific antigen (PSA) test rated their concern about prostate cancer lower than men who requested a PSA test, while men assigned to the usual care group provided similar ratings of concern regardless of their PSA choice.

**Attributes of the decision process: does the decision aid help patients to recognize that a decision needs to be made, know the options and their features, understand that values affect the decision, be clear about the features that matter most to them, discuss values with their clinician, and become involved in their preferred ways?**

In relation to the International Patient Decision Aids Standards (IPDAS) decision process criteria, no studies evaluated the extent to which patient decision aids helped participants to recognize that a decision needed to be made or understand that values affect the decision. Some studies measured participants' self-reports about feeling informed and clear about personal values. The measures used to evaluate these criteria were two subscales of the previously validated Decisional Conflict Scale (DCS) (O'Connor 1995).

### Decisional conflict


Of 105 studies, 63 (60.0%) evaluated decisional conflict using the DCS (O'Connor 1995). The DCS is reliable, discriminates between those who make or delay decisions, is sensitive to change, and discriminates between different decision support interventions (Morgan 2000; O'Connor 1995; O'Connor 1998b). The scale measures the constructs of overall decisional conflict and the particular factors contributing to uncertainty (e.g. feeling uncertain, uninformed, unclear about values, and unsupported in decision making). A final subscale measures perceived effective decision making. The scores were standardized to range from 0 (no decisional conflict) to 100 points (extreme decisional conflict). Scores of 25 or lower are associated with follow-through with decisions, whereas scores that exceed 38 are associated with delay in decision making (O'Connor 1998b). When decision aids are compared to usual care, a negative score indicates a reduction in decisional conflict, favouring the decision aid.

Analysis 4.1.1 summarizes the decisional conflict results for the 42 studies that compared decision aids to usual care. We report on 21 studies that were not amenable to pooling in Table 5 (original DCS), Table 6 (low literacy version), and Table 7 (SURE test version).

The mean difference (MD) for total DCS scores was  $-7.22$  points out of 100, favouring the patient decision aid over usual care groups (95% CI  $-9.12$  to  $-5.31$ ; see Analysis 4.1.1). Sixteen studies that could not be pooled (Table 5) reported mixed results on the original DCS. Of four studies that used the low literacy version (Fraenkel 2012; Smith 2010; Taylor 2006; Williams 2013), all reported statistically significant improvement (i.e. reduced) in total (or subscale) decisional conflict scores in the decision aid group, compared to usual care (Table 6). Stacey 2014a reported no difference between groups using the SURE test version.


The 'feeling uninformed' subscale of the DCS measures self-reported comfort with knowledge, not actual knowledge. We elected to consider this as a process measure and to reserve the gold standard of objective knowledge tests for assessing decision quality. There was high-quality evidence that patient decision aids were more effective than usual care in reducing patients' 'feeling uninformed' about options, benefits, and harms (MD  $-9.28$ , 95% CI  $-12.20$  to  $-6.36$ ; 27 studies; Analysis 4.1.2). The funnel plot for 'feeling uninformed' as an outcome in studies comparing decision aid to usual care shows low risk for publication bias (Figure 5).

**Figure 5. Funnel plot of comparison: 4.1 Decisional conflict: DA vs usual care - all studies, outcome: 4.1.2 Uninformed subscale**



There was high-quality evidence that patient decision aids were more effective than usual care for reducing patients' 'feeling unclear about values' subscale of the DCS (MD  $-8.81$ ; 95% CI  $-11.99$  to  $-5.63$ ; 23 studies; Analysis 4.1.3). The funnel plot for using 'feeling unclear about values' as an outcome in studies comparing decision aid to usual care shows low risk for publication bias (Figure 6).

**Figure 6. Funnel plot of comparison: 4.1 Decision conflict: DA vs usual care - all studies, outcome: 4.1.3 Unclear subscale**



#### Patient-clinician communication

Of 105 studies, 10 (9.5%) measured the effect of decision aids on patient-clinician communication. Of these 10 studies, 5 evaluated a patient decision aid used primarily within the consultation with the clinician, and 5 evaluated a patient decision aid used in preparation for the consultation.

Five studies compared the effect of usual care versus a decision aid used within the clinical encounter (or, in [Weymiller 2007](#), half the decision aid participants were exposed just prior to the encounter), evaluating the extent of shared decision making communication by analyzing the audio recordings using the OPTION scale ([Hess 2012](#); [LeBlanc 2015](#); [Montori 2011](#); [Mullan 2009](#); [Weymiller 2007](#)). The OPTION scale measures the extent to which health-care providers use behaviours that involve patients in decision making ([Elwyn 2005](#)). All five studies reported statistically higher mean OPTION scores in the patient decision aid group compared to usual care (see [Table 8](#)).

Four of five studies reported that compared to those in the usual care group, significantly higher proportions of participants exposed to the patient decision aid in preparation for the consultation reported that they discussed the decision with their clinician ([Fraenkel 2012](#); [Hanson 2011](#); [Lepore 2012](#); [Sheridan 2011](#); see

[Table 8](#)). The fifth study showed no between-group difference in discussion of cardiovascular disease with the clinician ([Sheridan 2006](#); see [Table 8](#)).

#### Participation in decision making

Of 105 studies, 24 (22.9%) measured the effect of decision aids on patients' perceived participation in decision making (Analysis 5.1; [Table 9](#)). [Davison 1997](#) used the Control Preferences Scale ([Degner 1992](#)). This scale uses five response statements to measure the role in decision making: two represent an active or patient-controlled role; one a shared or collaborative role; and two response statements represent a passive or clinician-controlled role. Most other studies used comparable response statements that could be classified within each of the three groupings of the Control Preferences Scale, except for [Hamann 2006](#), which used the COMRADE instrument to measure patient perception of involvement, and two others that used other measures of perceived involvement ([Hanson 2011](#); [Loh 2007](#); see [Table 9](#)).

Using the groupings of the Control Preferences Scale, 16 of 24 studies reported on clinician-controlled decision making. Consistent with the hypothesis that patient decision aids increase patient participation in decision making, there was moderate-quality ev-

idence that patient decision aids were more effective than usual care for reducing clinician-controlled decision making (RR 0.68; 95% CI 0.55 to 0.83; Analysis 5.1.1). In this field, there is no consensus on the hypothesized effects of decision aids on measures of patient-controlled decision making or shared decision making. Of 24 studies, 15 reported on participants assuming an active (patient-controlled) role in decision making and were pooled for analysis. Compared to usual care, decision aid use increased patient-controlled decision making (RR 1.28, 95% CI 1.05 to 1.55; Analysis 5.1.2). The 15 studies that reported on a shared decision-making role showed no difference between decision aid and usual care (RR 0.95; 95% CI 0.83 to 1.10; Analysis 5.1.3).

Of eight studies that could not be pooled, [Allen 2010](#), [Leighl 2011](#), [Rubel 2010](#), and [Van Peperstraten 2010](#) reported no between-group differences in these roles (Table 9). Three studies reported that a statistically significant proportion of patients exposed to the decision aid either participated ([Sheridan 2011](#)) - or at least felt involved - in decision making ([Hamann 2006](#); [Loh 2007](#)). However, [Hamann 2006](#) did not analyze results accounting for the use of design clusters. [Hanson 2011](#) reported that a higher proportion described feeling involved (83% vs. 77%), but the difference between groups was not statistically significant.

### Proportion undecided

Of 105 studies, 24 (22.9%) measured the proportion of participants remaining undecided: of these, 22 studies could be pooled. A significantly lower proportion of people remained undecided after exposure to a decision aid (RR 0.64; 95% CI 0.52 to 0.79; Analysis 6.1).

[Kasper 2008](#) measured progress in decision making using a single item ranging from '0 = completely undecided' to '100 = made my decision'. Given the difference in the measure Kasper used, these results were not included in the meta-analysis. In this study, both the patients exposed to a decision aid and the usual care group progressed in their decision making, with no difference between the groups (Table 10). [Sawka 2012](#) reported that 10.8% in the patient decision aid group versus 21.6% in the usual care group reported not knowing if they preferred taking adjuvant radioactive iodine.

### Satisfaction

Nineteen included studies (18.1%) measured satisfaction as it relates to the choice and the preparation for and the process of decision making. When possible, we standardized the scores to a 0 to 100 point scale, with higher scores reflecting greater satisfaction. Nineteen studies (18.1%) measured satisfaction with the choice. Of these 19 studies, 4 reported that people exposed to the decision aid had higher satisfaction with their choice compared to usual care, and the other 15 reported no statistically significant differences ([Chabrera 2015](#); [Heller 2008](#); [Laupacis 2006](#); [Montgomery](#)

[2007](#); see Analysis 7.1 and Table 11). For results that used a similar measure (Analysis 7.1), there was high satisfaction for all participants, with a median score of 82.5% for the decision aid and 80.0% for the usual care groups.

Of 105 total studies, 11 (10.5%) measured satisfaction with the decision, 11 (10.5%) measured satisfaction with the decision-making process (see Analysis 7.6; plus [Hess 2012](#) and [Vodermaier 2009](#) in Table 12), 4 measured satisfaction with information provided ([LeBlanc 2015](#); [Laupacis 2006](#); [Montori 2011](#); [Oakley 2006](#)), 3 measured satisfaction with the clinician ([Laupacis 2006](#); [Miller 2005](#); [Vodermaier 2009](#)), and 1 measured satisfaction with participating in decision making ([Kennedy 2002](#)). There were mixed results, but no studies reported that those exposed to patient decision aids were significantly less satisfied compared to usual care. For results that used a similar measure of satisfaction with the decision-making process (Analysis 7.4), there was high satisfaction for all participants, with median scores of 83.8% for the decision aid and 77.8% for the usual care groups. Although there were no differences between participant groups in satisfaction with the information in the [Montori 2011](#), clinicians using the decision aid had higher satisfaction.

Three studies (2.9%) measured satisfaction with preparation for decision making using the Preparation for Decision Making Scale ([Bennett 2010](#)) (Table 13). Compared to usual care, two studies reported significant improvements in people's satisfaction with their preparation for making decisions: in [Fraenkel 2007](#) after using decision aids about management of knee osteoarthritis, and in [Vandemheen 2009](#) regarding referral to a lung transplant centre. The third study found no statistically significant difference on this subscale's four items ([Stacey 2014a](#)).

## Secondary outcomes

### Behaviour

#### Choice

Choice was defined as the actual choice implemented. However, when studies did not report the actual choice, we used the patients' preferred option as a surrogate measure. Actual choices or preferences were reported as the percentage of individuals actually implementing or stating a preference for the most intensive or most invasive option.

In summary, patient decision aids decreased the number of patients choosing elective surgical procedures (excluding prophylactic mastectomy) and PSA testing in multiple studies. Single studies showed that decision aids increased the number of people choosing hepatitis B vaccination, psycho-educational therapies for mental health conditions, and medication for cardiovascular disease prevention. In contrast, decision aids decreased the rate of cardiac stress testing, the number of embryos being transplanted,



and the rate of antibiotic use for upper respiratory infections. The effect on patients' choice in other situations was more variable. There were mixed results for the choice of colon cancer screening, genetic testing, prenatal testing, anti-thrombosis therapy, breast screening, and diabetes medications. There was no difference between groups for choices about natural health products, hypertension therapy, breast cancer chemotherapy, schizophrenia medication, immunotherapy for multiple sclerosis, vaccines (for flu or measles, mumps, rubella), diabetes screening, birth control, osteoporosis treatment, chemotherapy for advanced cancer, chemopreventive medications, use of blood transfusions, childbirth procedures, treatment of prolapsed pelvic organs, or radioactive iodine treatment for thyroid cancer.

### *Choice for major elective surgery*

Eighteen studies (17.1%) focused on choices regarding major elective surgery (Analysis 8.1).

Using intention-to-treat analysis, there was a non-significant reduction in the number of patients choosing major elective surgery in the group receiving the decision aid compared to usual care (RR 0.86; 95% CI 0.75 to 1.00, 18 studies; Analysis 8.1.2). [Schwartz 2009a](#) reported a statistically significant uptake of prophylactic mastectomy for women who are BRCA1/2 gene carriers (114%). And after removing this study from the pooled results, there was a statistically significant reduction in the number of patients choosing major elective surgery (RR 0.84 95% CI 0.73 to 0.97; 17 studies; Analysis 8.1.3).

Four other studies showed statistically significant reductions in surgery rates: -29% for cardiac revascularization and bariatric surgery ([Arterburn 2011](#); [Morgan 2000](#)), -33% for orchiectomy ([Auvinen 2004](#)), and -74% for mastectomy ([Whelan 2004](#)). The other 15 studies showed no difference between the decision aid or usual care groups.

### *Choice for other elective surgery*

Two studies evaluated the effect of decision aids versus usual care on other elective surgical decisions. Decision aids did not significantly influence surgical abortion rates in [Wong 2006](#) or feeding tube insertions in [Hanson 2011](#) (Table 14).

### *Choice for prostate-specific antigen screening*

The effects of decision aids on prostate-specific antigen (PSA) screening decisions were variable in 13 studies (12.4%) that compared decision aids to usual care. The pooled RR for 10 studies was 0.88 (95% CI 0.80 to 0.98; Analysis 8.2.1); [Frosch 2008a](#), [Lepore 2012](#), and [Williams 2013](#) could not be included in the

pooled data (Table 14). [Frosch](#) reported a reduction in screening rates and the other two reported no difference.

### *Choice for colon cancer screening*

Of 10 studies (9.5%) on colon cancer screening, 3 reported statistically significant differences in choices, and 7 showed no difference. Two studies reported that compared to usual care, the decision aid significantly increased the screening rates by 64% and 70% ([Pignone 2000](#); [Ruffin 2007](#)). The other study reported a statistically significant reduction of 21% for screening ([Smith 2010](#)). There was an increase in screening rates in five studies, by 6% to 39%, but the difference was not statistically significant ([Lewis 2010](#); [Miller 2011](#); [Schroy 2011](#); [Steckelberg 2011](#); [Wolf 2000](#)). In two studies ([Dolan 2002](#); [Trevena 2008](#)), there was a 73% and 4% decrease in screening rates that was not statistically significant. The pooled RR was 1.12 (95% CI 0.95 to 1.31, 10 studies; Analysis 8.2.2).

### *Choice for cancer genetic screening*

Four studies reported preferences or uptake rates for breast cancer genetic screening (3.8%). The decision aid did not significantly affect preferences for breast cancer genetic screening when compared to usual care. The pooled RR was 0.99 (95% CI 0.71 to 1.38, 3 studies; Analysis 8.2.3). One study reported an increase in screening rates by 14% ([Lerman 1997](#)), a second study reported an increase of 18% ([Green 2001](#)), and a third study reported a decrease of 29% ([Schwartz 2001](#)). [Miller 2005](#) reported that women exposed to the decision aid who were at higher risk of breast cancer increased their intention to obtain genetic testing, while those at average risk decreased their intention (Table 14).

### *Choice for breast screening*

There were lower mammography screening rates among women aged 38 to 45 years of age ([Mathieu 2010](#)), but no between-group difference in women aged 70 or older who were exposed to a decision aid versus usual care ([Mathieu 2007](#); Table 14).

### *Choice for prenatal screening*

In all four studies focusing on decisions around prenatal screening, prenatal testing rates were not affected by a decision aid compared to usual care ([Bekker 2004](#); [Bjorklund 2012](#); [Kuppermann 2014](#); [Nagle 2008](#)). Meta-analysis included two studies, showing no effect (RR 0.99, 95% CI 0.91 to 1.09, 2 studies; [Bjorklund 2012](#); [Kuppermann 2014](#); Analysis 8.2.4).



### ***Choice for stress test for chest pain***

Compared to usual care, adults presenting with chest pain in the emergency department who received the decision aid had significantly lower rates of stress testing (58% versus 77%) (Hess 2012; Table 14).

### ***Choice for screening for diabetes***

Compared to usual care, there was no difference in diabetes screening rates in Marteau 2010 or preferences for screening in Mann E 2010 in adults exposed to a decision aid (Table 14).

### ***Choice to take antibiotics for upper respiratory infection***

Compared to usual care, using a decision aid in the consultation decreased prescriptions for antibiotics for upper respiratory infections in Legare 2012, although this difference was not statistically significant in Legare 2011 (Table 14).

### ***Choice for atrial fibrillation treatment***

Three studies evaluated the effect of a decision aid on the use of anti-thrombotic therapy for atrial fibrillation versus usual care (Table 14). One study demonstrated a non-significant reduction in warfarin use of 25% (Man-Son-Hing 1999). The second study evaluated the proportions of patients choosing the option that was appropriate relative to their level of risk, and found no significant difference between the groups (McAlister 2005). Thomson 2007 reported that patients in the usual care group (guided by practice recommendations) were much more likely to start warfarin (15/16; 93.8%) compared to the decision aid group (4/16; 25%; RR 0.27; 95% CI: 0.11 to 0.63).

### ***Choice to take breast cancer prevention medication***

There was no difference in medication use among women at risk of breast cancer who were exposed to the decision aid versus usual care (Fagerlin 2011; Table 14).

### ***Choice for cardiovascular disease prevention***

There was an increase in patient preferences for any effective cardiovascular disease risk-reducing strategy (including medication) when using a decision aid versus usual care (63% versus 42%) (Sheridan 2011; Table 14).

### ***Choice for chemotherapy for cancer***

There was no statistically significant difference in the rates of chemotherapy for adults with advanced colorectal cancer (77% versus 71%) (Leighl 2011; Table 14). Whelan 2003 also found no significant effect on preferences for adjuvant chemotherapy versus no chemotherapy for early stage breast cancer.

### ***Choice for diabetes treatment with new medications***

Four studies evaluated patient decision aids compared to usual care on decisions about starting new medications for diabetes (Mann D 2010; Mathers 2012; Mullan 2009; Weymiller 2007). Although there was no statistically significant difference between groups for individual studies, pooled results indicated a significant increase in starting new medications (RR 1.65, 95% CI 1.06 to 2.56; Analysis 8.3).

### ***Choice to take hypertension medication***

Montgomery 2003 found no significant effect for decision aids over usual care on the initiation of medication for hypertension (Table 14).

### ***Choice for menopausal symptom treatment***

In a study comparing a decision aid to usual care (Murray 2001b), there was a non-significant decrease of 8% in hormone therapy (Table 14). Preferences for natural health products in women experiencing menopausal symptoms were no different for women exposed to the decision aid compared to women exposed to the usual education materials (Legare 2008a).

### ***Choice for multiple sclerosis immunotherapy***

Kasper 2008 reported no difference in the uptake of immunotherapy in people with multiple sclerosis who were exposed to a decision aid compared to usual care based on practice guidelines (Table 14).

### ***Choice to take osteoporosis treatment***

There was no difference in prescriptions for bisphosphonates for osteoporosis treatment (LeBlanc 2015; Table 14). Montori 2011 found no significant effect of decision aids over usual care on the uptake of medication for osteoporosis treatment.

## Mental health

[Hamann 2006](#) found no difference in prescription rates for antipsychotic medications but reported a statistically significant increase in the uptake in psycho-education ( $P = 0.003$ ) in people with schizophrenia exposed to the decision aid compared to usual care ([Table 14](#)). [Mott 2014](#) reported that a higher proportion of participants in the decision aid group with post-traumatic stress disorder completed psychotherapy sessions (4 of 9) compared to usual care (1 of 11).

## Obstetrical choices

### *Childbirth procedures*

Three studies focused on childbirth issues, using a decision aid compared to usual care. There was no difference in preference for vaginal birth in [Shorten 2005](#) or actual vaginal mode of delivery in [Montgomery 2007](#) following a previous cesarean section. Another study found no difference in actual choice to undergo external cephalic version for women with breech presentation ([Nassar 2007](#)).

### *Birth control approaches*

There was no difference in the birth control methods chosen for those in the decision aid versus usual care groups ([Langston 2010](#)).

### *Embryo transplantation*

Compared to usual care, those in the decision aid group were significantly more likely to choose a single embryo transplant (43% versus 32%) ([Van Peperstraten 2010](#)).

## Vaccines

Compared to usual care, there was a non-significant increase in intentions to get the flu vaccine in those exposed to the decision aid (46% versus 27%) ([Chambers 2012](#)), a statistically significant increase in uptake of hepatitis B vaccination with decision aids ([Clancy 1988](#)), and no difference in uptake of measles, mumps, rubella vaccine in infants ([Shourie 2013](#)).

## Other choices

### *Blood transfusions*

There was no difference in the uptake of preoperative autologous blood donation when a decision aid was compared to usual care ([Laupacis 2006](#)).

### *Lung transplant referral*

There was no difference in referral rates for consideration of lung transplant in people with advanced cystic fibrosis exposed to a decision aid versus usual care ([Vandemheen 2009](#)).

### *Pelvic organ prolapse treatment*

There was no difference in treatment rates for prolapsed pelvic organs ([Brazell 2014](#)).

### *Thyroid cancer radioactive iodine treatment*

There was no difference in the rates of adjuvant radioactive iodine treatment for thyroid cancer ([Sawka 2012](#)).

## Adherence (continuance/compliance) with chosen option

Of 105 studies, 16 (15.2%) measured adherence using various approaches ([Table 15](#)).

Based on the measurement framework by [Trenaman 2016](#), we grouped adherence according to adherence to the baseline choice and adherence to the treatment. Six studies measured only adherence to the baseline choice ([Langston 2010](#); [Legare 2012](#); [Lepore 2012](#); [Man-Son-Hing 1999](#); [Mathers 2012](#); [Trevena 2008](#)), 6 studies measured only adherence to treatment ([Loh 2007](#); [Mann D 2010](#); [Mott 2014](#); [Mullan 2009](#); [Oakley 2006](#); [Sheridan 2011](#)), and 4 studies measured both ([LeBlanc 2015](#); [Montgomery 2003](#); [Montori 2011](#); [Weymiller 2007](#)).

For the 10 studies that measured adherence to choice, two studies reported that patients exposed to decision aids had higher adherence compared to usual care ([Mathers 2012](#); [Montori 2011](#)), and 8 reported no difference between groups. For example, [Mathers 2012](#) asked participants, 6 months after their decision, whether or not they had changed their initial choice about starting insulin for type II diabetes (decision aid 68.1% versus 56.3% usual care;  $P = 0.041$ ). Montori used pharmacy records to determine if participants who chose bisphosphonates actually took their medication on more than 80% of the days for which it was prescribed (100% decision aid versus 74% usual care;  $P = 0.009$ ).

For the 10 studies that measured adherence to treatment, 2 studies reported that patients exposed to decision aids had higher adherence compared to usual care ([Mott 2014](#); [Sheridan 2011](#)), 1 study reported that patients exposed to decision aids had lower adherence ([Mullan 2009](#)), and 7 reported no difference. Mott reported the percentage of participants at four months who engaged in nine

or more psychotherapy sessions (4 of 4 decision aid group participants versus 1 of 5 usual care). Sheridan measured the percentage of participants who, 3 months after initiating therapy, were continuing (59% decision aid versus 34% usual care;  $P < 0.01$ ). Mullan used pharmacy records to determine the days covered by medication use (97.5% decision aid versus 100% usual care).

## Health outcomes

### General health outcomes

Eleven studies (10.5%) compared a decision aid to usual care in terms of general health outcomes (Table 16). Ten of these used either the previously validated Medical Outcomes Study 36-item Short-Form Health Survey (SF-36) or the 12-item Short-Form Health Survey (SF-12) (Stewart 1992), while Vuorma 2003 used the RAND-36 (Hays 1993). As shown in Table 16, there were no significant differences for mental health function or social function in any of the seven studies. In one study (Barry 1997), general health and physical function outcome scores were significantly better in the decision aid group compared to usual care for men considering treatments for benign prostatic disease. Of the two studies evaluating the effect of a decision aid for women considering treatment for abnormal uterine bleeding, Kennedy 2002 found a statistically significant improvement in role physical function, and Vuorma 2003 found a statistically significant improvement in emotional role functioning for women.

In two studies measuring health utilities using the Euroqol EQ-5D (Murray 2001a; Murray 2001b), there was no difference between the decision aid and usual care groups. There was also no between-group difference in the LeBlanc 2015 study, which used the Euroqol 5D health thermometer.

### Condition-specific health outcomes

Seven studies (6.7%) used various measures to assess condition-specific health outcomes (Table 17). Outcomes included urinary symptoms (Barry 1997; Murray 2001a), angina (Bernstein 1998), functional assessment of cancer therapy (Leighl 2011), menopausal symptoms (Murray 2001b), and menstrual symptoms (Protheroe 2007; Vuorma 2003). Five studies found no significant effects on condition-specific health outcomes (Bernstein 1998; Leighl 2011; Murray 2001a; Murray 2001b; Vuorma 2003). Protheroe 2007 reported significantly higher menorrhagia-related quality of life scores in women exposed to the decision aid compared to usual care. Barry 1997 showed an improvement in urinary symptoms in favour of the decision aid group, but it was not statistically significant.

### Other health outcomes

Seven studies (6.7%) reported on other health outcomes (Table 18), including death (Auvinen 2004; Knops 2014), glycated haemoglobin (Mathers 2012), angina (Morgan 2000), stroke (Thomson 2007), successful pregnancy (Van Peperstraten 2010), and pain (Vuorma 2003). There were no statistically significant differences between groups.

### Preference-linked health outcomes

None of the 105 studies measured preference-linked health outcomes - that is, whether the patients experienced the outcomes they preferred and avoided the outcomes they wanted to avoid.

### Anxiety

Of 105 studies, 31 (29.5%) measured anxiety, with 24 using the previously validated State Trait Anxiety Inventory (Spielberger 1970), 2 using the anxiety subscale of the Hospital Anxiety and Depression Scale (Knops 2014; Lam 2013), 2 using questions about worry (Fraenkel 2012; Smith 2010), 2 measuring intrusive thoughts (Lewis 2010; McCaffery 2010), and 1 using a single question on a seven-point Likert scale (Johnson 2006; see Table 19). Of 18 studies that used the State Trait Anxiety inventory within 1 month postintervention, 2 (11.1%) reported that the decision aid group had significantly lower anxiety scores for people considering birthing options after a previous caesarean (Montgomery 2007) and for women considering options for the treatment of menorrhagia (Protheroe 2007). None of the studies demonstrated significant differences in effects on people's state anxiety at one month (2 studies), three months (6 studies), six months (4 studies), or one year (2 studies). There was no significant difference between groups for the other instruments that measured anxiety.

### Depression

Of 105 studies, 6 (5.7%) measured the effect of decision aids on depression using various instruments (Table 20). None of the studies reported a statistically significant difference between groups for decisions about cancer treatment (Davison 1997; Whelan 2004), depression (Loh 2007), prenatal genetic testing (Nagle 2008), or for women considering the number of embryos to transplant (Van Peperstraten 2010). At 10 months' postintervention, there were lower levels of depression in women deciding about breast cancer surgery who were exposed to the patient decision aid versus the usual care, but no differences at 1 week, 1 month, or 4 months postintervention (Lam 2013).

### Regret

Of 105 studies, 7 (6.7%) measured the effect of decision aids on decision regret, using the five-item Decisional Regret scale (Brehaut

2003; see Table 21). At 4 and 10 months postintervention, women with breast cancer who were considering surgery and used a decision aid reported lower regret scores compared to women receiving usual care (Lam 2013). There was no statistically significant between-group difference in the other six studies.

### Confidence

Of 105 studies, 8 (7.8%) measured the effect of decision aids on confidence levels (see Table 22). Four of these studies used the Decisional Self-efficacy Scale (Allen 2010; Arterburn 2011; Fraenkel 2007; Smith 2010). Four studies reported a statistically significant improvement in confidence or self-efficacy with decision making in the decision aid compared to the usual care groups (Chambers 2012; Fraenkel 2007; Gattellari 2003; McBride 2002), and the other studies reported no difference between groups.

### Healthcare system effects

#### Cost and resource use

Of eight studies (7.6%) examining cost and resource use, one conducted a cost-effectiveness analysis (Kennedy 2002), five evaluated the effect of decision aids compared to usual care on total costs (Montgomery 2007; Murray 2001a; Murray 2001b; Van Peperstraten 2010; Vuorma 2003), and two measured resource use (Legare 2012; Thomson 2007) (see Table 23).

The cost-effectiveness analysis (Kennedy 2002) was conducted from the healthcare system perspective, using USD values from 1999 to 2000 and calculating costs over two years. The decision aid with nurse coaching demonstrated the lowest mean cost (USD 1566) compared to decision aid alone (USD 2026) or usual care (USD 2751).

Of the five studies that evaluated total costs, two reported no statistically significant difference in the patient decision aid compared to usual care (Montgomery 2007; Vuorma 2003). Two studies reported higher costs for the patient decision aid group when including the cost of the interactive video disc equipment (USD 216 at 1999 prices) and no statistically significant difference between groups when removing this cost (Murray 2001a; Murray 2001b). The fifth study reported that the mean total savings in the decision aid group versus usual care was EUR 169.75 per couple (Van Peperstraten 2010).

For healthcare resource use in upper respiratory infection, Legare 2012 reported no difference in the rates of repeat consultations for the same reason, and Thomson 2007 reported no difference in the rates of general clinician consultations in the three months following the intervention. Both studies used the patient decision aid in the consultation.

### Consultation length

Of 105 studies, 10 (9.5%) evaluated the effect of a decision aid compared to usual care on consultation length (see Table 23). The median consultation length was 24 minutes (range 3.8 to 68.3) for patient decision aid compared to 21 minutes (range 4.2 to 65.7) for usual care. The difference was 2.6 minutes longer (7.5% increase) than usual care consultations (range 0.4 minutes shorter to 23 minutes longer). The length of consultation was significantly longer for the patient decision aid group in two studies (Bekker 2004; Thomson 2007), and eight studies reported no difference. Bekker 2004 reported that consultations about prenatal diagnostic testing were 5.9 minutes longer, and Thomson 2007 reported consultations about treatment for atrial fibrillation were 23 minutes longer when using a computerized decision aid with standard gamble method within the consultation.

### Litigation rates

None of the 105 studies examined the effect of decision aids on litigation.

### Adverse events

There were no adverse effects on health outcomes or satisfaction, and no other adverse events reported.

### Subgroup analysis - in preparation for versus during the consultation

Of 105 studies, 89 (84.8%) primarily evaluated the patient decision aid when used by the patient in preparation for the consultation, and 16 (15.2%) primarily evaluated the patient decision aid when used within the consultation. The patient decision aids used during the consultation focused on prenatal screening (Bekker 2004); cardiac stress testing (Hess 2012); dental surgery (Johnson 2006); restoration of tooth decay (Kupke 2013); antibiotics for upper respiratory infection (Legare 2011; Legare 2012); medication use for depression (Loh 2007), diabetes (Mann D 2010; Mullan 2009; Weymiller 2007), osteoporosis (LeBlanc 2015; Montori 2011), prevention of breast cancer (Ozanne 2007), and atrial fibrillation (Thomson 2007); surgery for breast cancer (Whelan 2004); and chemotherapy for breast cancer (Whelan 2003).

### Knowledge

When considered separately by subgroups, there was no difference between knowledge scores for those exposed to the decision aid in preparation for the consultation compared to those used in the consultation itself (Analysis 1.2: MD 13.77% versus 10.57%, test for subgroup difference  $P = 0.31$ ,  $I^2 = 3\%$ ). Weymiller 2007 reported a higher mean difference when the decision aid was administered during the consultation but not if it was administered

by research staff in preparation for the consultation. For the studies evaluating decision aids used in the consultation not included in the pooled outcome, two showed a statistically significant improvement in knowledge (LeBlanc 2015; Ozanne 2007), and two showed no difference (Mann D 2010; Thomson 2007).

### Accurate risk perceptions

When analyzing pre-consultation and in-consultation decision aids further, accurate risk perceptions were not different between studies that used the decision aid in preparation for the consultation and those where the intervention occurred during the consultation (Analysis 2.2: RR 2.25 versus RR 1.79, test for subgroup differences:  $P = 0.33$ ,  $I^2 = 0\%$ ). The only study evaluating a decision aid within the consultation that was not included in the meta-analysis, Weymiller 2007, reported a higher proportion with accurate risk perception when the decision aid was administered during the consultation, but found no difference between groups when administered by research staff in preparation for the consultation.

### Decisional conflict uninformed subscale

Too few studies measured the uninformed subscale in those exposed to decision aid within the consultation to be able to compare with those who used decision aids in preparation for the consultation. Weymiller 2007 reported that participants felt less uninformed when the decision aid was administered during the consultation, but not if it was administered by research staff in preparation for the consultation.

### Decisional conflict unclear values subscale

Too few studies measured the unclear values subscale in those exposed to decision aid within the consultation to be able to compare with those who used decision aids in preparation for the consultation. Weymiller 2007 reported that participants felt less unclear about values when the decision aid was administered during the consultation, but not if it was administered by research staff in preparation for the consultation.

### Patient-clinician communication

Due to variation in the reporting of data for this outcome, we were unable to investigate the effect of intervention timing on the variation in the effect on communication. Five studies evaluated a patient decision aid primarily used within the consultation with the clinician, and five evaluated a patient decision aid used in preparation for the consultation (see Table 8). All five studies that used the decision aid during consultations reported statistically higher mean OPTION scores in the patient decision aid group compared to usual care (Hess 2012; LeBlanc 2015; Montori 2011; Mullan 2009; Weymiller 2007). Four of five studies assessing the effects

of pre-consultation decision aid delivery (Fraenkel 2012; Hanson 2011; Lepore 2012; Sheridan 2011) reported that, compared to those in the usual care group, significantly higher proportions of participants exposed to the patient decision aid in preparation for the consultation reported that they discussed the decision with their clinician, and the fifth study showed no between-group difference (Sheridan 2006).

### Participation in decision making

There were too few studies on decision aids used during the consultation to interpret findings from the subgroup analysis (Analysis 5.2; Analysis 5.3).

### Length of the consultation

Due to variation in the reporting of data for this outcome, we were unable to investigate the effects of intervention timing on the length of consultation. Of seven studies that evaluated decision aids used within the consultation (Bekker 2004; LeBlanc 2015; Loh 2007; Ozanne 2007; Thomson 2007; Weymiller 2007; Whelan 2003), two reported that the length of the consultation was significantly longer for the patient decision aid group (Bekker 2004; Thomson 2007). There was no difference for the other studies. The three studies that evaluated decision aids used in preparation for the consultation reported no between group difference in the length of the consultation (Bozic 2013; Krist 2007; Vodermaier 2009).

### Other outcomes

For values-choice congruence and proportion undecided, none of the studies of patient decision aids used during the consultation measured these outcomes. For satisfaction, there were a range of different approaches to measuring this outcome with mixed results and too few studies to make any descriptive comparisons. For choice, there were too few studies to conduct a subgroup analysis of pooled comparisons.

### Post hoc analysis

#### Effects of study quality

To examine the potential bias arising from including studies of low methodological quality, we excluded 12 studies with a high risk of bias for any of the seven risk of bias criteria from the analysis (Auvinen 2004; Brazell 2014; Chambers 2012; Clancy 1988; Hamann 2006; Knops 2014; Krist 2007; Kupke 2013; LeBlanc 2015; Lewis 2010; Man-Son-Hing 1999; Mott 2014; see Figure 3). Overall, the results remained the same (Table 24; Analysis 1.3; Analysis 2.3; Analysis 3.5; Analysis 4.4).



## Heterogeneity

When comparing patient decision aids to usual care, there was statistically significant heterogeneity in five of six of the IPDAS effectiveness criteria: knowledge scores, accurate risk perceptions, congruence between values and choice; feeling uninformed, and feeling unclear regarding personal values. There was no statistically significant heterogeneity for participation in decision making. It should be noted that the heterogeneity of the effect was not manifested in its direction but only in its size. For the 2009 update (O'Connor 2009b), we explored the potential factors contributing to heterogeneity (Table 25). Overall, regardless of the subgroup analyses conducted, scores for outcomes were similar to the overall effect, as indicated by overlapping confidence intervals.

## DISCUSSION

### Summary of main results

In this updated review, we added 18 new studies for a total of 105 studies comparing patient decision aids to usual care. This update also removed 28 studies that compared detailed versus simple patient decision aids that were included in the previous update. Based on the GRADE assessment (Summary of findings for the main comparison), there is high-quality evidence that compared to usual care, decision aids improve people's knowledge regarding options and reduce the decisional conflict stemming from feeling uninformed and unclear about their personal values. There is moderate-quality evidence that decision aids stimulate people to take a more active role in decision making and increase the accuracy of their risk perceptions. There is lower-quality evidence that decision aids improve congruence between the chosen option and personal values. This outcome is measured using a variety of different approaches, and the evidence could be strengthened by more standardized measurement. Moreover, decision aids decreased the proportion of people remaining undecided.

Although not a primary outcome of the review, the effect of decision aids on patients' choosing particular options continues to be variable. The numbers of patients choosing to have major elective surgery continues to decrease in favour of more conservative options, except when the baseline rates are low (e.g. surgery for benign prostate hyperplasia, prophylactic mastectomy for women who are carriers of the BRCA gene). The numbers of men choosing prostate-specific antigen (PSA) testing were fewer after exposure to decision aids.

Decision aids do no better than usual care in terms of their effects on people's satisfaction with decision making or health outcomes such as general quality of life or condition-specific quality of life. However, no studies measured preference-linked health outcomes, nor were adverse events reported. There was also no difference in anxiety. For length of consultation, eight studies found no difference, while two studies found a median increase of 2.6 minutes

(7.5%) in the decision aid group compared to usual care consultations. There continue to be too few studies to determine the effects of decision aids on costs/resource use (Trenaman 2014). Although there may be additional costs involved in delivering decision aids, an independent review of decision aid studies with economic outcomes concluded that "this was likely to be small relative to the benefit to patients in terms of improved decision quality when effective decision aids are used" (NCGC/NICE 2012). Given the variability in measurement strategies, it difficult to determine the effect of patient decision aids on adherence to the chosen option or treatment.

New for this update, we analyzed the pooled data for decision aids used in preparation for the consultation separately from decision aids used in the consultation, and we found that there were similar improvements in knowledge, accurate risk perceptions, and patient-clinician communication.

### Overall completeness and applicability of evidence

#### Main effects of decision aids

The largest and most consistent benefits of decision aids, relative to usual care, are better knowledge of options and outcomes, and more accurate perceptions of outcome probabilities. These observations are clinically important because the usual care groups' scores for knowledge and perception of outcome probabilities were lower than the intervention groups'; both knowledge and perception of outcome probabilities are important for ensuring informed decision making. These effects suggest that current 'usual care' may not be good enough when informing people about these complex, values-sensitive decisions. People need to comprehend the options and outcome probabilities in order to consider and communicate to their clinicians the personal value they place on the benefits versus the harms. Likewise, pooling results from additional studies in this update shows a significant increase in informed values-based choice when decision aids were compared to usual care, and the results appear to be similar across subgroup analyses of studies that used the same composite measure.

Decision aids also help people feel more comfortable with their choices than usual care. This is revealed by the reduced scores for overall decisional conflict and for the decisional conflict subscales. People who use decision aids generally feel more informed about options and clearer regarding their personal values.

Compared to usual care strategies, decision aids improve individuals' perception of involvement in decision making. This observation suggests that the International Patient Decision Aids Standards criterion of helping patients participate 'in ways that they prefer' needs to be assessed after a patient has adequate information about what involvement means using interventions such as patient decision aids. People may have a mistaken preference for

passivity because they believe that the best choice relies on the expertise of the clinician (which option is medically reasonable?) rather than understand the importance of their own preferences for outcomes of options (which outcomes matter most to me?). Evidence continues to build that decision aids have a positive effect on the patient-clinician consultation (in 9 of the 10 studies that assessed this effect). Of the studies that measured patient-clinician communication, five involved using decision aids within the consultation and five in preparation for the consultation. At the same time, evidence on length of consultation indicates either no difference (8 studies) or slightly longer (2 studies) consultations in the decision aid group compared to usual care consultations. However, few studies have reported on the impact of the context in which the patient decision aids are used. A previous subgroup analysis of 29 studies evaluating patient decision aids for treatment decisions reported greater improvement in knowledge scores ( $P = 0.03$ ) when the patient decision aid was evaluated within the clinical pathway of care, compared to when patients volunteered to participate in the study independent of their clinician (Brown 2015).

### Variable effects of decision aids

There may be several reasons for the variable effect of decision aids on the outcome of choices. First, most studies were under-powered to detect important differences in the outcome of choices. Second, not enough is known about baseline rates for optimal use of specific options. Third, in the studies reporting the outcome 'choices' at baseline and postdecision aid, some options may have been under-used and others over-used, relative to the choices individuals would make if they were more fully informed. Under these circumstances, one could expect to observe directional effects on choices once people become better informed and more involved in decision making.

Relatively under-used options at baseline were prostate surgery for benign prostatic hyperplasia and prophylactic mastectomy for breast cancer gene carriers. In this prostate-related example, there was a shortage of urologists and low referral rates for benign prostatic hyperplasia, whereas the breast-related example reflects the growing number of women who test gene positive and become aware of their options for preventing breast cancer. Hence, under-use of an option may be corrected with exposure to a decision aid. In the other surgical decision aid studies, there were higher numbers of people choosing surgery in the control group (e.g. cardiac revascularization, back surgery, hysterectomy, orchiectomy, mastectomy). The procedure may have been chosen due to people's inflated perceptions of the probabilities of benefits, lack of appreciation of the probabilities of harms, and lack of awareness of alternatives (Hoffman 2015). Exposure to the decision aid reduced the number of people choosing elective surgery in favour of more conservative alternatives.

### Limited effects of decision aids

The limited effects of decision aids on reported satisfaction with the decision-making process and with the actual choice made may indicate that decision aids have a limited effect on satisfaction. The null effects may also be due to measurement insensitivity. This is especially likely when satisfaction with usual care is already quite high (e.g. ceiling effects) and when choices are inherently difficult to make because of competing benefits and harms. Furthermore, once the decision is made, people may find it psychologically more comforting to say that they are satisfied rather than entertain doubts about what they have chosen (Gruppen 1994). There is a need to establish the 'essential ingredients' in decision aids and to identify the people who are most likely to benefit from them. As the body of available research grows, it will become easier and more important to assess the usefulness of different components of decision support for different clinical contexts, decision problems, and groups of people. For example, an analysis of decision aids used in higher versus lower socioeconomic groups indicated greater improvements for those of lower socioeconomic status (Durand 2014). Recently, the IPDAS Collaboration completed a set of evidence reviews underlying the IPDAS checklist (IPDAS 2013), proposing criteria for defining the intervention as a patient decision aid and minimal certifying criteria (Joseph-Williams 2013). These are being used to inform the certification of patient decision aids in the USA, England, and Norway.

It is not surprising that decision aids had limited effects on health outcomes. One reason for using a decision aid is that there is often no option with a clear health outcome advantage. For example, when men with localized prostate cancer consider active treatment options, their health outcomes can be different, depending on whether they choose surgery with higher risks of impotence or radiation therapy with higher risks of longer term bowel irritation. Therefore, if health outcomes are used in future investigations of decision aids in situations in which there is clearly no health outcome advantage, the key question to pose is: do patients experience the health outcomes they prefer and avoid the outcomes to which they are averse?

More recently, decision aids are being used in situations in which there may be a longer-term health advantage, for example, in preventive decisions about the management of type II diabetes and/or hypertension, when the longer-term health outcome may be to avoid stroke (Mann D 2010; Mathers 2012; Montgomery 2003; Mullan 2009; Weymiller 2007). Interestingly, the pooled results showed a statistically significant increase in medication initiation when participants were exposed to the decision aid compared to usual care.

### Unknown effects of decision aids

The effect of patient decision aids on adherence to the chosen option is an area of uncertainty. The adherence results are difficult to interpret due to incomplete data, primarily self-reported data,

varying length of follow-ups, and small sample sizes. Moreover, studies reporting this outcome such as [Man-Son-Hing 1999](#) had very little variation in choice (over 90% of long-term aspirin users decided to stay on aspirin). When examining adherence, it would be important to do so in the early phase, when presumably the issue is actually decisional in nature (e.g. filling the prescription, picking up the prescription, refilling the prescription) rather than involving the management of side effects and in a manner that separates those choosing to change versus those remaining with the status quo.

Despite the positive effects of decision aids on patient-clinician communication, some authors are concerned about the potential negative influence that decision aids may have on the relational aspects of the decision-making process; this concern highlights the need for further evaluation when decision aids are implemented as part of the routine process of care ([Charles 2010](#); [LeBlanc 2010](#)). In the context of decision aid use, cost-effectiveness and health utilities are other secondary outcome measures about which little is known and further evaluation is required ([Trenaman 2014](#)). We also need to establish ways of measuring preference-linked health outcomes to better determine the effect on quality of life. It is unlikely that we will observe the effect of decision aids on litigation rates in studies of decision aids, given the time delay to litigation and the rarity of this type of event. There do not appear to be any adverse events from using decision aids, but this could be more clearly examined in future studies. In fact, a mock trial that used a patient decision aid for prostate-specific antigen testing found that the majority of jurors (94%) would indicate that the standard of care had been met ([Barry 2008](#)). A recent systematic review concluded that there was insufficient evidence to determine if patient decision aids could reduce medical malpractice litigation ([Durand 2014](#)).

## Quality of the evidence

Risk of bias ratings reveal between-study variability. We rated few studies as being at low risk of bias for blinding of participants and personnel and most studies as being at unclear risk of bias. Likewise, the majority of studies were rated as being at unclear risk of bias for selective reporting. When we conducted a post hoc analysis that involved removing studies at high risk of bias from the meta-analysis, there was no effect on the results. The conclusions of this review are limited by inadequate power to detect important between-subgroup differences in effectiveness and by the wide variability in the decision contexts, the elements within the patient decision aids, the type of comparison delivered (collectively referred to as usual care here), the targeted outcomes, and the evaluation procedures. The small number of studies for most outcomes did not allow for analysis of publication bias due to failure to publish negative studies. Moreover, most studies were at unclear risk of selective outcome reporting, indicating that there

may have been bias arising from a failure to report all negative findings.

We rated the six primary outcomes in the 'Summary of findings' table using GRADE and assessed outcomes as high quality (knowledge, feeling uninformed, feeling unclear values), moderate quality (accurate risk perception, clinician-controlled role in decision making), and low quality (values-choice congruence). For values-choice congruence, the GRADE rating was downgraded for lack of consistency, directness, and precision. More specifically, congruence was measured using various approaches, as there is no gold standard measurement approach ([Munro 2016](#)). Several of the outcomes demonstrated statistically significant levels of heterogeneity. For the outcome of knowledge, for example, heterogeneity would be expected, given that the knowledge tests themselves were not standardized. However, we did not downgrade the ratings for knowledge, feeling uninformed, and feeling unclear values based on heterogeneity given the consistent direction of findings across studies. Moreover, the heterogeneity found in the various outcomes reflects differences across clinically diverse studies; therefore, the pooled effect size and confidence intervals should be interpreted as a range across conditions, which may not be applicable to a specific condition.

## Potential biases in the review process

The strength of this systematic review is that patient decision aids improve several key primary outcomes across a wide variety of populations and decision contexts. The potential biases in the review process are due to limitations associated with having inadequate power to detect potentially important differences in effectiveness between subgroups, to differentiate between the most effective elements within the patient decision aid, and to investigate any differences associated with the type of comparison interventions used in studies. Several of the outcomes demonstrated statistically significant heterogeneity. This reflects differences across clinically diverse studies; therefore, the pooled effect size and confidence intervals should be interpreted as a range across conditions, which may not be applicable to a specific condition. In the [Gentles 2013](#) subgroup analysis exploring three potential sources of heterogeneity (e.g. type of control intervention, decision aid IPDAS quality score, participants' baseline accurate risk perception), participants' baseline accurate risk perception was an important variable for explaining heterogeneity. Authors reported that when participants' baseline scores for accurate risk perception were lower, decision aids led to great improvement. Furthermore, we limited the extracted study data to only two comparison groups (e.g. most intensive intervention including a patient decision aid and usual care); therefore, we did not investigate the possibility of intermediate effects with less intensive decision aid interventions.



## Agreements and disagreements with other studies or reviews

Our results confirm many of the observations reported in the previous versions of our review and in a comparative effectiveness review that focused on studies evaluating oncology-specific patient decision aids (Trikalinos 2014). We published the first systematic review of 17 randomized trials of decision aids in 1999 (O'Connor 1999b; O'Connor 2001), followed by updates in 2003 with a total of 35 studies (O'Connor 2003), in 2009 with a total of 55 studies (O'Connor 2009b), in 2011 with a total of 86 studies (Stacey 2011), and 2014 with a total of 115 studies (Stacey 2014b).

## AUTHORS' CONCLUSIONS

### Implications for practice

The positive effects of decision aids on improving people's knowledge of risks and benefits, feeling informed, and feeling clear about their values across a wide variety of decision contexts provides sufficient evidence for using them in clinical practice. They probably also facilitate accurate risk perception and active participation in decision making. However, several conditions may be necessary for successful implementation, including: good quality decision aids that meet the needs of the population; clinicians who are willing to use decision aids in their practice; effective systems for delivering decision support; and clinicians and healthcare consumers who are skilled in shared decision making. Although there have been some strides in achieving these conditions (Elwyn 2013; O'Connor 2007), the use of patient decision aids will not occur without adequate attention to implementation barriers to implementing and maintaining their use in routine clinical practice (Elwyn 2013; Gravel 2006; Legare 2008b; Legare 2010; Legare 2014).

New in this update was a subgroup analysis of the findings based on timing of decision aid used either before or during a consultation. Although knowledge scores and accurate risk perceptions were significantly higher in the decision aid group compared to the usual care, there was no difference in these outcomes when comparing decision aids used in preparation for versus during the consultation.

## Implications for research

Studies are needed to deepen our understanding of interactions between patient decision aid use and the patterns of patient-clinician communication; format issues such as the web-based delivery of patient decision aids; and downstream effects on cost, resource use, and adherence. Although this update shows new studies conducted in Spain and China, most studies have taken place in North America, the UK, Europe, and Australia. There were far fewer studies of patient decision aids used within the consultation than those delivered pre-consultation, and this is an area of further research given the important issue of implementation.

With the addition of more studies in the systematic review, it may be possible to tease out the reasons for heterogeneity of results, including variability in: study quality; comparison intervention; elements within patient decision aids; decision type; setting where it was used; and format of decision aid (e.g. video, Internet, booklet). Research should also explore the degree of detail in patient decision aids that is required for positive effects according to the IPDAS criteria. In particular, evaluation is needed to compare the effect of those decision aids that meet the minimal IPDAS criteria for certification versus those that meet the full roster of IPDAS quality criteria (Joseph-Williams 2013).

## ACKNOWLEDGEMENTS

The Cochrane Consumers and Communication Group (editors, academic and consumer referees) provided peer review and advice regarding the review and checked extracted data for newly included studies. We thank John Kis-Rigo at La Trobe University who revised the search strategy used since the 2014 update. David Rovner and Nananda Col helped with screening studies for inclusion, and Intissar Souli assisted with data extraction and 'Risk of bias' assessment. Anton Saarimaki set up and managed a web-based title and abstract screening application that facilitated independent screening of citations by the review authors, and he also verified references for patient decision aids in Table 1. Alain Mayhew provided guidance in the interpretation of the 'Summary of findings' table. Dean Fergusson provided consultation on the statistical analysis.

## REFERENCES

### References to studies included in this review

#### Allen 2010 *{published data only}*

Allen JD, Othus MK, Hart A Jr, Tom L, Li Y, Berry D, et al. A randomized trial of a computer-tailored decision aid to improve prostate cancer screening decisions: results from the take the wheel trial. *Cancer Epidemiology, Biomarkers and Prevention* 2010;**19**(9):2172–86.

Allen JD, Othus MKD, Hart A Jr, Mohllajee AP, Bowen D. Do men make informed decisions about prostate cancer screening? Baseline results from the “Take the Wheel” Trial. *Medical Decision Making* 2011;**31**:108–120.

#### Arterburn 2011 *{published data only}*

Arterburn D, Westbrook E, Bogart T, Sepucha K, Bock S, Weppner W. Randomized trial of a video-based patient decision aid for bariatric surgery. *Obesity* 2011;**19**(8):1669–75.

#### Auvinen 2004 *{published data only}*

\* Auvinen A, Hakama M, Ala-Opas M, Vornanen T, Leppilähti M, Salminen P, et al. A randomized trial of choice of treatment in prostate cancer: the effect of intervention on the treatment chosen. *BJU International* 2004;**93**(1):52–6.  
Auvinen A, Vornanen T, Tammela TL, Ala-Opas M, Leppilähti M, Salminen P, et al. A randomized trial of the choice of treatment in prostate cancer: design and baseline characteristics. *BJU International* 2001;**88**(7):708–15.  
Huang RC, Auvinen A, Hakama M, Tammela TLJ, Ala-Opas M, Leppilähti M, et al. Effect of intervention on decision making of treatment for disease progression, prostate-specific antigen biochemical failure and prostate cancer death. *Health Expectations* 2014;**17**(6):776–83.

#### Barry 1997 *{published and unpublished data}*

\* Barry MJ, Cherkin DC, Chang Y, Fowler FJ, Skates S. A randomized trial of a multimedia shared decision-making program for men facing a treatment decision for benign prostatic hyperplasia. *Disease Management and Clinical Outcomes* 1997;**1**(1):5–14.  
Rovner DR, Wills CE, Bonham V, Williams G, Lillie J, Kelly-Blake K, et al. Decision aids for benign prostatic hyperplasia: applicability across race and education. *Medical Decision Making* 2004;**24**(4):359–66.

#### Bekker 2004 *{published data only}*

\* Bekker HL, Hewison J, Thornton JG. Applying decision analysis to facilitate informed decision making about prenatal diagnosis for Down syndrome: a randomised controlled trial. *Prenatal Diagnosis* 2004;**24**(4):265–75.  
Bekker HL, Hewison J, Thornton JG. Understanding why decision aids work: linking process with outcome. *Patient Education and Counseling* 2003;**50**(3):323–9.

#### Bernstein 1998 *{published and unpublished data}*

Bernstein SJ, Skarupski KA, Grayson CE, Starling MR, Bates ER, Eagle KA. A randomized controlled trial of information-giving to patients referred for coronary angiography: effects on outcomes of care. *Health Expectations* 1998;**1**(1):50–61.

#### Berry 2013 *{published data only}*

\* Berry DL, Halpenny B, Hong F, Wolpin S, Lober WB, Russell KJ, et al. The personal patient profile-prostate decision support for men with localized prostate cancer: a multi-center randomized trial. *Urologic Oncology* 2013;**31**(7):1012–21.

Berry DL, Wang Q, Halpenny B, Hong F. Decision preparation, satisfaction and regret in a multi-center sample of men with newly diagnosed localized prostate cancer.

*Patient Education and Counseling* 2012;**88**(2):262–7.

Bosco JLF, Halpenny B, Berry DL. Personal preferences and discordant prostate cancer treatment choice in an intervention trial of men newly diagnosed with localized prostate cancer. *Health and Quality of Life Outcomes* 2012;**10**(123):1–8.

Underhill ML, Hong F, Berry DL. When study site contributes to outcomes in a multi-center randomized trial: a secondary analysis of decisional conflict in men with localized prostate cancer. *Health and Quality of Life Outcomes* 2014;**12**:159.

#### Bjorklund 2012 *{published data only}*

Bjorklund U, Marsk A, Levin C, Ohman SG. Audiovisual information affects informed choice and experience of information in antenatal Down syndrome screening—a randomized controlled trial. *Patient Education and Counseling* 2012;**86**(3):390–5.

Öhman SG, Björklund U, Marsk A. Does an informational film increase women's possibility to make an informed choice about second trimester ultrasound?. *Prenatal Diagnosis* 2012;**32**(9):833–9.

#### Bozic 2013 *{published data only}*

Bozic KJ, Belkora J, Chan V, Youm J, Zhou T, Dupax J, et al. Shared decision making in patients with osteoarthritis of the hip and knee: results of a randomized controlled trial. *Journal of Bone and Joint Surgery: American Volume* 2013;**95**(18):1633–9.

Bozic KJ, Chenok KE, Schindel J, Chan V, Huddleston JI, Braddock C, Belkora J. Patient, surgeon, and healthcare purchaser views on the use of decision and communication aids in orthopaedic surgery: a mixed methods study. *BMC Health Services Research* 2014;**14**(366):1–10.

Youm J, Chan V, Belkora J, Bozic KJ. Impact of socioeconomic factors on informed decision making and treatment choice in patients with hip and knee OA. *The Journal of Arthroplasty* 2015;**30**(2):171–5.

#### Brazell 2014 *{published data only}*

Brazell HD, O'Sullivan DM, Forrest A, Greene JF. Effect of a decision aid on decision making for the treatment of pelvic organ prolapse. *Female Pelvic Medicine & Reconstructive Surgery* 2014;**21**(4):231–5.

#### Chabrera 2015 *{published data only}*

Chabrera C, Zabalegui A, Bonet M, Caro M, Areal J, González JR, Font A. A decision aid to support informed

- choices for patients recently diagnosed with prostate cancer. *Cancer Nursing* 2015;**38**(3):E42–E50.
- Chambers 2012** *{published data only}*  
Chambers LW, Wilson K, Hawken S, Puxty J, Crowe L, Lam PP, et al. Impact of the Ottawa influenza decision aid on healthcare personnel's influenza immunization decision: a randomized trial. *Journal of Hospital Infection* 2012;**82**(3):194–202.
- Clancy 1988** *{published data only}*  
Clancy CM, Cebul RD, Williams SV. Guiding individual decisions: a randomized, controlled trial of decision analysis. *American Journal of Medicine* 1988;**84**(2):283–8.
- Davison 1997** *{published data only}*  
Davison BJ, Degner LF. Empowerment of men newly diagnosed with prostate cancer. *Cancer Nursing* 1997;**20**(3):187–96.
- De Achaval 2012** *{published data only}*  
De Achaval S, Fraenkel L, Volk R, Cox V, Suarez-Almazor M. Impact of educational and patient decision aids on decisional conflict associated with total knee arthroplasty. *Arthritis Care & Research* 2012;**64**(2):229–37.
- Dolan 2002** *{published data only}*  
Dolan JG, Frisina S. Randomized controlled trial of a patient decision aid for colorectal cancer screening. *Medical Decision Making* 2002;**22**(2):125–39.
- Evans 2010** *{published data only}*  
Evans R, Joseph-Williams N, Edwards A, Newcombe R, Wright P, Kinnersley P, et al. Supporting informed decision making for prostate specific antigen (PSA) testing on the web: an online randomized controlled trial. *Journal of Medical Internet Research* 2010;**12**(3):e27.
- Fagerlin 2011** *{published data only}*  
Banegas MP, McClure JB, Barlow WE, Ubel PA, Smith DM, Zikmund-Fisher BJ, et al. Results from a randomized trial of a web-based, tailored decision aid for women at high risk for breast cancer. *Patient Education and Counseling* 2013;**91**:364–71.  
Fagerlin A. Randomization for Guide to Decide phase II. Word document provided by the authors.  
\* Fagerlin A, Dillard AJ, Smith DM, Zikmund-Fisher BJ, Pitsch R, McClure JB, et al. Women's interest in taking tamoxifen and raloxifene for breast cancer prevention: response to a tailored decision aid. *Breast Cancer Research and Treatment* 2011;**127**(3):681–8.  
Korfage IJ, Fuhrel-Forbis A, Ubel PA, Zikmund-Fisher BJ, Greene SM, McClure JB, et al. Informed choice about breast cancer prevention: randomized controlled trial of an online decision aid intervention. *Breast Cancer Research* 2013;**15**(R74):1–9.
- Fraenkel 2007** *{published data only}*  
Fraenkel L, Rabidou N, Wittink D, Fried T. Improving informed decision-making for patients with knee pain. *Journal of Rheumatology* 2007;**34**(9):1894–8.
- Fraenkel 2012** *{published data only}*  
Fraenkel L, Street RL Jr, Towle V, O'Leary JR, Iannone L, Van Ness PH, Fried TR. A pilot randomized controlled trial of a decision support tool to improve the quality of communication and decision-making in individuals with atrial fibrillation. *Journal of the American Geriatrics Society* 2012;**60**(8):1434–41.
- Frosch 2008a** *{published data only}*  
Frosch DL, Bhatnagar V, Tally S, Hamori CJ, Kaplan RM. Internet patient decision support: a randomized controlled trial comparing alternative approaches for men considering prostate cancer screening. *Archives of Internal Medicine* 2008;**168**(4):363–9.
- Gattellari 2003** *{published data only}*  
Gattellari M, Ward JE. Does evidence-based information about screening for prostate cancer enhance consumer decision-making? A randomised controlled trial. *Journal of Medical Screening* 2003;**10**(1):27–39.
- Gattellari 2005** *{published data only}*  
Gattellari M, Ward JE. A community-based randomised controlled trial of three different educational resources for men about prostate cancer screening. *Patient Education and Counseling* 2005;**57**(2):168–82.
- Green 2001** *{published data only}*  
Green MJ, Biesecker BB, McInerney AM, Mauger D, Fost N. An interactive computer program can effectively educate patients about genetic testing for breast cancer susceptibility. *American Journal of Medical Genetics* 2001;**103**(1):16–23.
- Hamann 2006** *{published data only}*  
Hamann J, Cohen R, Leucht S, Busch R, Kissling W. Shared decision making and long-term outcome in schizophrenia treatment. *Journal of Clinical Psychiatry* 2007;**68**(7):992–7.  
\* Hamann J, Langer B, Winkler V, Busch R, Cohen R, Leucht S, et al. Shared decision making for in-patients with schizophrenia. *Acta Psychiatrica Scandinavica* 2006;**114**(4):265–73.
- Hanson 2011** *{published data only}*  
Ersek M, Sefcik JS, Feng-Chang L, Lee TJ, Gilliam R, Hanson LC. Provider staffing effect on a decision aid intervention. *Clinical Nursing Research* 2014;**23**:36–53.  
Hanson L, Carey T, Caprio A, Joon Lee T, Ersek M, Garrett J, et al. Improving decision making for feeding options in advanced dementia: a randomized, controlled trial. *Journal of the American Geriatrics Society* 2011;**59**(11):2009–16.  
Snyder EA, Caprio AJ, Wessell K, Lin FC, Hanson LC. Impact of a decision aid on surrogate decision-makers' perceptions of feeding options for patients with dementia. *American Medical Directors Association* 2013;**14**(2):114–8.
- Heller 2008** *{published data only}*  
Heller L, Parker PA, Youssef A, Miller MJ. Interactive digital education aid in breast reconstruction. *Plastic & Reconstructive Surgery* 2008;**122**(3):717–24.
- Hess 2012** *{published data only}*  
Hess EP, Knoedler MA, Shah ND, Kline JA, Breslin M, Branda ME, et al. The chest pain choice decision aid: a randomized trial. *Circulation: Cardiovascular Quality and Outcomes* 2012;**5**(3):251–9.

**Jibaja-Weiss 2011** {published data only}

Jibaja-Weiss M, Volk R, Granchi T, Neff N, Robinson E, Spann S, et al. Entertainment education for breast cancer surgery decisions: a randomized trial among patients with low health literacy. *Patient Education and Counseling* 2011; **84**(1):41–8.

**Johnson 2006** {published data only}

Johnson BR, Schwartz A, Goldberg J, Koerber A. A chairside aid for shared decision making in dentistry: a randomized controlled trial. *Journal of Dental Education* 2006; **70**(2):133–41.

**Kasper 2008** {published data only}

Kasper J, Kopke S, Muhlhauser I, Nubling M, Heesen C. Informed shared decision making about immunotherapy for patients with multiple sclerosis (ISDIMS): A randomized controlled trial. *European Journal of Neurology* 2008; **15**(12): 1345–52.

**Kennedy 2002** {published data only}

Kennedy AD, Sculpher MJ, Coulter A, Dwyer N, Rees M, Abrams KR, et al. Effects of decision aids for menorrhagia on treatment choices, health outcomes, and costs: a randomized controlled trial. *JAMA* 2002; **288**(21):2701–8.

**Knops 2014** {published data only}

Knops AM, Goossens A, Ubbink DT, Balm R, Koelemay MJ, Vahl AC, et al. DECAID Trial Group. A decision aid regarding treatment options for patients with an asymptomatic abdominal aortic aneurysm: a randomised clinical trial. *European Journal of Vascular and Endovascular Surgery* 2014; **48**(3):276–283.

**Krist 2007** {published data only}

Krist AH, Woolf SH, Johnson RE, Kerns JW. Patient education on prostate cancer screening and involvement in decision making. *Annals of Family Medicine* 2007; **5**(2): 112–9.

**Kupke 2013** {published data only (unpublished sought but not used)}

Kupke J, Wicht MJ, Stützer H, Derman SH, Lichtenstein NV, Noack MJ. Does the use of a visualised decision board by undergraduate students during shared decision-making enhance patients' knowledge and satisfaction? A randomised controlled trial. *European Journal of Dental Education* 2013; **17**(1):19–25.

**Kuppermann 2014** {published data only}

Kuppermann M, Pena S, Bishop JT, Nakagawa S, Gregorich SE, Sit A, et al. Effect of enhanced information, values clarification, and removal of financial barriers on use of prenatal genetic testing: a randomized clinical trial. *Journal of the American Medical Association* 2014; **312**(12):1210–7.

**Lam 2013** {published data only}

Lam WW, Chan M, Or A, Kwong A, Suen D, Fielding R. Reducing treatment decision conflict difficulties in breast cancer surgery: a randomized controlled trial. *Journal of Clinical Oncology* 2013; **31**(23):2879–85.

**Langston 2010** {published data only}

Langston A, Rosario L, Westhoff C. Structured contraceptive counseling: a randomised controlled trial. *Patient Education and Counseling* 2010; **81**(3):362–7.

**Laupacis 2006** {published data only}

Laupacis A, O'Connor AM, Drake ER, Rubens FD, Robblee JA, Grant FC, et al. A decision aid for autologous pre-donation in cardiac surgery - a randomized trial. *Patient Education and Counseling* 2006; **61**(3):458–66.

**LeBlanc 2015** {published data only}

LeBlanc A, Wang AT, Wyatt K, Branda ME, Shah ND, Van Houten H, et al. Encounter decision aid vs. clinical decision support or usual care to support patient-centered treatment decisions in osteoporosis: the osteoporosis choice randomized trial II. *PLOS ONE* 2015; **10**(5):1–13.

**Legare 2008a** {published data only}

Legare F, Dodin S, Stacey D, Leblanc A, Tapp S. Patient decision aid on natural health products for menopausal symptoms: randomized controlled trial. *Menopause International* 2008; **14**(3):105–10.

**Legare 2011** {published data only}

Legare F, Labrecque M, LeBlanc A, Njoya M, Laurier C, Cote L, et al. Training family physicians in shared decision making for the use of antibiotics for acute respiratory infections: a pilot clustered randomized controlled trial. *Health Expectations* 2011; **14**:96–110.

**Legare 2012** {published and unpublished data}

Legare F, Labrecque M, Cauchon M, Castel J, Turcotte S, Grimshaw J. Training family physicians in shared decision-making to reduce the overuse of antibiotics in acute respiratory infections: a cluster randomized trial. *Canadian Medical Association Journal* 2012; **184**(13):E726–34.

**Leighl 2011** {published data only}

Leighl NB, Shepherd HL, Butow PN, Clarke SJ, McJannett M, Beale PJ, et al. Supporting treatment decision making in advanced cancer: a randomized trial of a decision aid for patients with advanced colorectal cancer considering chemotherapy. *Journal of Clinical Oncology* 2011; **29**(15): 2077–84.

**Lepore 2012** {published data only}

Lepore SJ, Wolf RL, Basch CE, Godfrey M, McGinty E, Shmukler C, et al. Informed decision making about prostate cancer testing in predominantly immigrant black men: a randomized controlled trial. *Annals of Behavioral Medicine* 2012; **44**(3):320–30.

**Lerman 1997** {published data only}

Lerman C, Biesecker B, Benkendorf JL, Kerner J, Gomez-Caminero A, Hughes C, et al. Controlled trial of pretest education approaches to enhance informed decision-making for BRCA1 gene testing. *Journal of the National Cancer Institute* 1997; **89**(2):148–57.

**Lewis 2010** {published data only}

\* Lewis C, Pignone M, Schild L, Scott T, Winquist A, Rimer B, et al. Effectiveness of a patient and practice-level colorectal cancer screening intervention in health plan members: design and baseline findings of the CHOICE trial. *Cancer* 2010; **116**(7):1664–73.  
Pignone M, Winquist A, Schild L, Lewis C, Scott T, Hawley J, et al. Effectiveness of a patient and practice-

- level colorectal cancer screening intervention in health plan members. *Cancer* 2011;**117**(15):3252–62.
- Pignone MP, Brenner AT, Hawley S, Sheridan SL, Lewis CL, Jonas DE, et al. Conjoint analysis versus rating and ranking for values elicitation and clarification in colorectal cancer screening. *Journal of General Internal Medicine* 2011; **27**(1):45–50.
- Loh 2007** *{published data only}*  
Loh A, Simon D, Harter M. Effects of shared decision making in primary care of depressive patients - better compliance and treatment effects. *Klinikerarzt* 2007;**36**(1): 38–41.  
\* Loh A, Simon D, Wills CE, Kriston L, Niebling W, Harter M. The effects of a shared decision-making intervention in primary care of depression: a cluster-randomized controlled trial. *Patient Education and Counseling* 2007;**67**(3):324–32.
- Mann D 2010** *{published data only}*  
Mann DM, Ponienman D, Montori VM, Arciniega J, McGinn T. The statin choice decision aid in primary care: a randomized trial. *Patient Education and Counseling* 2010; **80**(1):138–40.
- Mann E 2010** *{published data only}*  
Mann E, Kellar I, Sutton S, Kinmonth AL, Hankins M, Griffin S, et al. Impact of informed-choice invitations on diabetes screening knowledge, attitude and intentions: an analogue study. *BMC Public Health* 2010;**10**:768.
- Man-Son-Hing 1999** *{published and unpublished data}*  
Man-Son-Hing M, Laupacis A, O'Connor AM, Biggs J, Drake E, Yetisir E, et al. A patient decision aid regarding antithrombotic therapy for stroke prevention in atrial fibrillation: a randomized controlled trial. *JAMA* 1999;**282** (8):737–43.
- Marteau 2010** *{published data only}*  
Kellar I, Mann E, Kinmonth AL, Prevost AT, Sutton S, Marteau TM. Can informed choice invitations lead to inequities in intentions to make lifestyle changes among participants in a primary care diabetes screening programme? Evidence from a randomized trial. *Public Health* 2011;**125**(9):645–52.  
\* Marteau TM, Mann E, Prevost AT, Vasconcelos JC, Kellar I, Sanderson S, et al. Impact of an informed choice invitation on uptake of screening for diabetes in primary care (DICISION): randomised trial. *BMJ* 2010;**340**:c2138.
- Mathers 2012** *{published data only}*  
Brown I, Bradley A, Ng CJ, Colwell B, Mathers N. Investigating active ingredients in a complex intervention: a nested study within the Patient and Decision Aids (PANDAs) randomised controlled trial for people with type 2 diabetes. *BMC Research Notes* 2014;**7**:347.  
Mathers N, Ng CJ, Campbell MJ, Colwell B, Brown I, Bradley A. Clinical effectiveness of a patient decision aid to improve decision quality and glycaemic control in people with diabetes making treatment choices: a cluster randomised controlled trial (PANDAs) in general practice. *BMJ Open* 2012;**2**(6):1–12.
- Mathieu 2007** *{published data only}*  
Mathieu E, Barratt A, Davey HM, McGeechan K, Howard K, Houssami N. Informed choice in mammography screening: a randomized trial of a decision aid for 70-year-old women. *Archives of Internal Medicine* 2007;**167**(19): 2039–46.
- Mathieu 2010** *{published data only}*  
Mathieu E, Barratt AL, McGeechan K, Davey HM, Howard K, Houssami N. Helping women make choices about mammography screening: an online randomized trial of a decision aid for 40-year-old women. *Patient Education and Counseling* 2010;**81**(1):63–72.
- McAlister 2005** *{published data only}*  
McAlister FA, Man-Son-Hing M, Straus SE, Ghali WA, Anderson D, Majumdar SR, et al. Impact of a patient decision aid on care among patients with nonvalvular atrial fibrillation: a cluster randomized trial. *CMAJ* 2005;**173**(5): 496–501.
- McBride 2002** *{published data only}*  
Bastian LA, McBride CM, Fish L, Lyna P, Farrell D, Lipkus IM, et al. Evaluating participants' use of a hormone replacement therapy decision-making intervention. *Patient Education and Counseling* 2002;**48**(3):283–91.  
\* McBride CM, Bastian LA, Halabi S, Fish L, Lipkus IM, Bosworth HB, et al. A tailored intervention to aid decision making about hormone replacement therapy. *American Journal of Public Health* 2002;**92**(7):1112–4.
- McCaffery 2010** *{published data only}*  
McCaffery KJ, Irwig L, Turner R, Chan SF, Macaskill P, Lewicka M, et al. Psychosocial outcomes of three triage methods for the management of borderline abnormal cervical smears: an open randomised trial. *BMJ* 2010;**340**: b4491.
- Miller 2005** *{published data only}*  
Miller SM, Fleisher L, Roussi P, Buzaglo JS, Schnoll R, Slater E, et al. Facilitating informed decision making about breast cancer risk and genetic counseling among women calling the NCI's Cancer Information Service. *Journal of Health Communication* 2005;**10**(Suppl 1):119–36.
- Miller 2011** *{published data only}*  
Duren-Winfield V, Onsomu EO, Case DL, Pignone M, Miller D. Health literacy and computer-assisted instruction: usability and patient preference. *Journal of Health Communication* 2015;**20**:491–8.  
Miller D, Spangler J, Case D, Goff D, Singh S, Pignone M. Effectiveness of a web-based colorectal cancer screening patient decision aid: a randomized controlled trial in a mixed-literacy population. *American Journal of Preventive Medicine* 2011;**40**(6):608–15.
- Montgomery 2003** *{published and unpublished data}*  
Emmett CL, Montgomery AA, Peters TJ, Fahey T. Three-year follow-up of a factorial randomised controlled trial of two decision aids for newly diagnosed hypertensive patients. *British Journal of General Practice* 2005;**55**(516):551–3.  
\* Montgomery AA, Fahey T, Peters TJ. A factorial randomised controlled trial of decision analysis and

- an information video plus leaflet for newly diagnosed hypertensive patients. *British Journal of General Practice* 2003;**53**(491):446–53.
- Montgomery 2007** {published data only}  
Frost J, Shaw. Women's views on the use of decision aids for decision making about the method of delivery following a previous caesarean section: qualitative interview study. *British Journal of Obstetrics and Gynecology* 2009;**116**(7): 896–905.  
Hollinghurst S, Emmett C, Peters TJ, Watson H, Fahey T, Murphy DJ, et al. Economic evaluation of the DIAMOND randomized trial: cost and outcomes of 2 decision aids for mode of delivery among women with previous caesarian section. *BMJ* 2010;**30**:453–63.  
\* Montgomery AA, Emmett CL, Fahey T, Jones C, Ricketts I, Patel RR, et al. Two decision aids for mode of delivery among women with previous caesarean section: randomised controlled trial. *BMJ* 2007;**334**(7607):1305.
- Montori 2011** {published data only}  
\* Montori VM, Shah ND, Pencille LJ, Branda ME, Van Houten HK, Swiglo BA. Use of a decision aid to improve treatment decisions in osteoporosis: the osteoporosis choice randomized trial. *American Journal of Medicine* 2011;**124**(6):549–56.  
Pencille LJ, Campbell ME, Van Houten HK, Shah ND, Mullan RJ, Swiglo BA, et al. Protocol for the Osteoporosis Choice trial. A pilot randomized trial of a decision aid in primary care practice. *Trials* 2009;**10**:113.
- Morgan 2000** {published and unpublished data}  
Morgan MW. *A Randomized Trial of the Ischemic Heart Disease Shared Decision Making Program: An Evaluation of a Decision Aid [Masters Thesis]*. Toronto: University of Toronto, 1997.  
\* Morgan MW, Deber RB, Llewellyn-Thomas HA, Gladstone P, Cusimano RJ, O'Rourke K, et al. Randomized, controlled trial of an interactive videodisc decision aid for patients with ischemic heart disease. *Journal of General Internal Medicine* 2000;**15**(10):685–93.
- Mott 2014** {published data only}  
Mott JM, Stanley MA, Street RL Jr, Grady RH, Teng EJ. Increasing engagement in evidence-based PTSD treatment through shared decision-making: a pilot study. *Military Medicine* 2014;**179**(2):143–9.
- Mullan 2009** {published data only}  
Mullan RJ, Montori VM, Shah ND, Christianson TJ, Bryant SC, Guyatt GH, et al. The diabetes mellitus medication choice decision aid: a randomized trial. *Archives of Internal Medicine* 2009;**169**(17):1560–8.
- Murray 2001a** {published and unpublished data}  
Murray E, Davis H, Tai SS, Coulter A, Gray A, Haines A. Randomised controlled trial of an interactive multimedia decision aid on benign prostatic hypertrophy in primary care. *BMJ* 2001;**323**(7311):493–6.
- Murray 2001b** {published and unpublished data}  
Murray E, Davis H, Tai SS, Coulter A, Gray A, Haines A. Randomized controlled trial of an interactive multimedia decision aid on hormone replacement therapy in primary care. *BMJ* 2001;**323**(7311):490–3.
- Nagle 2008** {published data only}  
\* Nagle C, Gunn J, Bell R, Lewis S, Meiser B, Metcalfe S, et al. Use of a decision aid for prenatal testing of fetal abnormalities to improve women's informed decision making: a cluster randomised controlled trial. *BJOG: An International Journal of Obstetrics and Gynaecology* 2008;**115**(3):339–47.  
Nagle C, Lewis S, Meiser B, Metcalfe S, Carlin JB, Bell R, et al. Evaluation of a decision aid for prenatal testing of fetal abnormalities: a cluster randomised trial [ISRCTN22532458]. *BMC Public Health* 2006;**6**:96.
- Nassar 2007** {published data only}  
Nassar N, Roberts CL, Raynes-Greenow CH, Barratt A, Pear B, Decision Aid for Breech Presentation Trial Collaborators. Evaluation of a decision aid for women with breech presentation at term: a randomised controlled trial [ISRCTN14570598]. *BJOG: An International Journal of Obstetrics & Gynaecology* 2007;**114**(3):325–33.
- Oakley 2006** {published data only}  
Oakley S, Walley T. A pilot study assessing the effectiveness of a decision aid on patient adherence with oral bisphosphonate medication. *Pharmaceutical Journal* 2006;**276**(7399):536–8.
- Ozanne 2007** {published data only}  
Ozanne EM, Annis C, Adduci K, Showstack J, Esserman L. Pilot trial of a computerized decision aid for breast cancer prevention. *Breast Journal* 2007;**13**(2):147–54.
- Partin 2004** {published and unpublished data}  
Partin MR, Nelson D, Flood AB, Friedemann-Sanchez G, Wilt TJ. Who uses decision aids? Subgroup analyses from a randomized controlled effectiveness trial of two prostate cancer screening decision support interventions. *Health Expectations* 2006;**9**(3):285–95.  
\* Partin MR, Nelson D, Radosevich D, Nugent S, Flood AB, Dillon N, et al. Randomized trial examining the effect of two prostate cancer screening educational interventions on patient knowledge, preferences, and behaviors. *Journal of General Internal Medicine* 2004;**19**(8):835–42.
- Pignone 2000** {published data only}  
Pignone M, Harris R, Kinsinger L. Videotape-based decision aid for colon cancer screening. A randomized, controlled trial. *Annals of Internal Medicine* 2000;**133**(10): 761–9.
- Protheroe 2007** {published data only}  
Patel S, Ngunjiri A, Hee SW, Yang Y, Brown S, Friede T, et al. Primum non nocere: shared informed decision making in low back pain - a pilot cluster randomised trial. *BMC Musculoskeletal Disorders* 2014;**15**:282.  
Protheroe J, Bower P, Chew-Graham C. The use of mixed methodology in evaluating complex interventions: identifying patient factors that moderate the effects of a decision aid. *Family Practice* 2008;**24**(6):594–600.  
\* Protheroe J, Bower P, Chew-Graham C, Peters TJ, Fahey T. Effectiveness of a computerized decision aid in primary

- care on decision making and quality of life in menorrhagia: results of the MENTIP randomized controlled trial. *Medical Decision Making* 2007;**27**(5):575–84.
- Rubel 2010** {published data only}  
Rubel SK, Miller JW, Stephens RL, Xu Y, Scholl LE, Holden EW, et al. Testing the effects of a decision aid for prostate cancer screening. *Journal of Health Communication* 2010;**15**(3):307–21.
- Ruffin 2007** {published data only}  
Ruffin MT, Fetter MD, Jimbo M. Preference-based electronic decision aid to promote colorectal cancer screening: results of a randomized controlled trial. *Preventive Medicine* 2007;**45**(4):267–73.
- Sawka 2012** {published and unpublished data}  
Sawka AM, Straus S, Rotstein L, Brierley JD, Tsang RW, Asa S, et al. Randomized controlled trial of a computerized decision aid on adjuvant radioactive iodine treatment for patients with early-stage papillary thyroid cancer. *Journal of Clinical Oncology* 2012;**30**(23):2906–11.
- Schroy 2011** {published data only}  
\* Schroy PC 3rd, Emmons K, Peters E, Glick JT, Robinson PA, Lydotes MA, et al. The impact of a novel computer-based decision aid on shared decision making for colorectal cancer screening: a randomized trial. *Medical Decision Making* 2011;**31**(1):93–107.  
Schroy PC 3rd, Emmons KM, Peters E, Glick JT, Robinson PA, Lydotes MA, et al. Aid-assisted decision making and colorectal cancer screening: a randomized controlled trial. *American Journal of Preventive Medicine* 2012;**43**(6):573–83.
- Schwalm 2012** {published data only}  
Schwalm JD, Stacey D, Pericak D, Natarajan MK. Radial artery versus femoral artery access options in coronary angiogram procedures: randomized controlled trial of a patient-decision aid. *Circulation: Cardiovascular Quality and Outcomes* 2012;**5**(3):260–6.
- Schwartz 2001** {published data only}  
Schwartz MD, Benkendorf J, Lerman C, Isaacs C, Ryan-Robertson A, Johnson L. Impact of educational print materials on knowledge, attitudes, and interest in BRCA1/BRCA2: testing among Ashkenazi Jewish women. *Cancer* 2001;**92**(4):932–40.
- Schwartz 2009a** {published data only}  
Hooker GW, Leventhal KG, DeMarco T, Peshkin BN, Finch C, Wahl E, et al. Longitudinal changes in patient distress following interactive decision aid use among BRCA1/2 carriers: a randomized trial. *Medical Decision Making* 2011;**31**(3):412–21.  
\* Schwartz MD, Valdimarsdottir HB, DeMarco TA, Peshkin BN, Lawrence W, Rispoli J, et al. Randomized trial of a decision aid for BRCA1/BRCA2 mutation carriers: impact on measures of decision making and satisfaction. *Health Psychology* 2009;**28**(1):11–9.
- Sheridan 2006** {published data only}  
Sheridan SL, Shadle J, Simpson RJ Jr, Pignone MP. The impact of a decision aid about heart disease prevention on patients' discussions with their doctor and their plans for prevention: a pilot randomized trial. *BMC Health Services Research* 2006;**6**:121.
- Sheridan 2011** {published data only}  
Sheridan SL, Draeger LB, Pignone MP, Keyserling TC, Simpson RJ Jr, Rimer B, et al. A randomized trial of an intervention to improve use and adherence to effective coronary heart disease prevention strategies. *BMC Health Services Research* 2011;**11**:331.  
Sheridan SL, Draeger LB, Pignone MP, Rimer B, Bangdiwala SI, Cai J, Gizlice Z, Keyserling TC, Simpson RJ. The effect of a decision aid intervention on decision making about coronary heart disease risk reduction: secondary analyses of a randomized trial. *BMC Medical Informatics and Decision Making* 2014;**14**(14):1–11.
- Shorten 2005** {published and unpublished data}  
Shorten A, Shorten B, Keogh J, West S, Morris J. Making choices for childbirth: a randomized controlled trial of a decision-aid for informed birth after cesarean. *Birth* 2005;**32**(4):252–61.
- Shourie 2013** {published data only}  
Shourie S, Jackson C, Cheater FM, Bekker HL, Edlin R, Tubeuf S, et al. A cluster randomised controlled trial of a web based decision aid to support parents' decisions about their child's Measles Mumps and Rubella (MMR) vaccination. *Vaccine* 2013;**31**(50):6003–10.
- Smith 2010** {published data only}  
Smith SK, Barratt A, Trevena L, Simpson JM, Jansen J, McCaffery KJ. A theoretical framework for measuring knowledge in screening decision aid trials. *Patient Education and Counseling* 2012;**89**:330–6.  
Smith SK, Kearney P, Trevena L, Barratt A, Nutbeam D, McCaffery KJ. Informed choice in bowel cancer screening: a qualitative study to explore how adults with lower education use decision aids. *Health Expectations* 2012;**17**:511–22.  
Smith SK, Simpson JM, Trevena LJ, McCaffery KJ. Factors associated with informed decisions and participation in bowel cancer screening among adults with lower education and literacy. *Medical Decision Making* 2014;**34**(6):756–72.  
Smith SK, Trevena L, Simpson JM, Barratt A, Nutbeam D, McCaffery KJ. A decision aid to support informed choices about bowel cancer screening among adults with low education: randomised controlled trial. *BMJ* 2010;**341**:c5370.
- Stacey 2014a** {published and unpublished data}  
Stacey D, Hawker G, Dervin G, Tugwell P, Bolland L, Pomey MP, et al. Decision aid for patients considering total knee arthroplasty with preference report for surgeons: A pilot randomized controlled trial. *BMC Musculoskeletal Disorders* 2014;**15**:54.
- Steckelberg 2011** {published data only}  
Steckelberg A, Hulfenhaus C, Haastert B, Muhlhauser I. Effect of evidence based risk information on "informed choice" in colorectal cancer screening: randomised controlled trial. *BMJ* 2011;**342**:d3193.

**Taylor 2006** {published data only}

Taylor KL, Davis JL 3rd, Turner RO, Johnson L, Schwartz MD, Kerner JF, et al. Educating African American men about the prostate cancer screening dilemma: a randomized intervention. *Cancer Epidemiology, Biomarkers & Prevention* 2006;**15**(11):2179–88.

**Thomson 2007** {published data only}

Kaner E, Heaven B, Rapley T, Murtagh M, Graham R, Thomson R, et al. Medical communication and technology: a video-based process study of the use of decision aids in primary care consultations. *BMC Medical Informatics and Decision Making* 2007;**7**(2):1–11.

\* Thomson RG, Eccles MP, Steen IN, Greenaway J, Stobbart L, Murtagh MJ, et al. A patient decision aid to support shared decision-making on anti-thrombotic treatment of patients with atrial fibrillation: randomised controlled trial. *Quality & Safety in Health Care* 2007;**16**(3):216–23.

**Trevena 2008** {published data only}

Trevena LJ, Irwig L, Barratt A. Randomized trial of a self-administered decision aid for colorectal cancer screening. *Journal of Medical Screening* 2008;**15**(2):76–82.

**Vandemheen 2009** {published data only}

Vandemheen KL, O'Connor A, Bell SC, Freitag A, Bye P, Jeanneret A, et al. Randomized trial of a decision aid for patients with cystic fibrosis considering lung transplantation. *American Journal of Respiratory & Critical Care Medicine* 2009;**180**(8):761–8.

**Van Peperstraten 2010** {published data only}

Kreuwel I, van Peperstraten A, Hulscher M, Kremer J, Grol R, Nelen W, Hermens R. Evaluation of an effective multifaceted implementation strategy for elective single-embryo transfer after in vitro fertilization. *Human Reproduction* 2013;**28**(2):336–42.

Van Peperstraten A, Nelen W, Grol R, Zielhuis G, Adang E, Stalmeier P, et al. The effect of a multifaceted empowerment strategy on decision making about the number of embryos transferred in in vitro fertilisation: randomised controlled trial. *BMJ* 2010;**341**:c2501.

**Vodermaier 2009** {published data only}

Vodermaier A, Caspari C, Koehm J, Kahlert S, Ditsch N, Untch M. Contextual factors in shared decision making: a randomised controlled trial in women with a strong suspicion of breast cancer. *British Journal of Cancer* 2009;**100**(4):590–7.

**Volk 1999** {published and unpublished data}

\* Volk RJ, Cass AR, Spann SJ. A randomized controlled trial of shared decision making for prostate cancer screening. *Archives of Family Medicine* 1999;**8**(4):333–40.

Volk RJ, Spann SJ, Cass AR, Hawley ST. Patient education for informed decision making about prostate cancer screening: a randomized controlled trial with 1-year follow-up. *Annals of Family Medicine* 2003;**1**(1):22–8.

**Vuorma 2003** {published data only}

\* Vuorma S, Rissanen P, Aalto AM, Hurskainen R, Kujansuu E, Teperi J. Impact of patient information booklet on

treatment decision - a randomized trial among women with heavy menstruation. *Health Expectations* 2003;**6**(4):290–7.

Vuorma S, Teperi J, Aalto AM, Hurskainen R, Kujansuu E, Rissanen P. A randomized trial among women with heavy menstruation - impact of a decision aid on treatment outcomes and costs. *Health Expectations* 2004;**7**(4):327–37.

**Watson 2006** {published data only}

Watson E, Hewitson P, Brett J, Bukach C, Evans R, Edwards A, et al. Informed decision making and prostate specific antigen (PSA) testing for prostate cancer: a randomised controlled trial exploring the impact of a brief patient decision aid on men's knowledge, attitudes and intention to be tested. *Patient Education and Counseling* 2006;**63**(3):367–79.

**Weymiller 2007** {published data only}

Jones LA, Weymiller AJ, Shah N, Bryant SC, Christianson TJH, Guyatt GH, et al. Should clinicians deliver decision aids? further exploration of the statin choice randomized trial results. *Medical Decision Making* 2009;**29**(4):468–74.

Nannenga MR, Montori VM, Weymiller AJ, Smith SA, Christianson TJ, Bryant SC, et al. A treatment decision aid may increase patient trust in the diabetes specialist. The Statin Choice randomized trial. *Health Expectations* 2009;**12**(1):38–44.

\* Weymiller AJ, Montori VM, Jones LA, Gafni A, Guyatt GH, Bryant SC, et al. Helping patients with type 2 diabetes mellitus make treatment decisions: statin choice randomized trial. *Archives of Internal Medicine* 2007;**167**(10):1076–82.

**Whelan 2003** {published and unpublished data}

Whelan T, Sawka C, Levine M, Gafni A, Reyno L, Willan A, et al. Helping patients make informed choices: a randomized trial of a decision aid for adjuvant chemotherapy in lymph node-negative breast cancer. *Journal of the National Cancer Institute* 2003;**95**(8):581–7.

**Whelan 2004** {published and unpublished data}

Whelan T, Levine M, Willan A, Gafni A, Sanders K, Mirsky D, et al. Effect of a decision aid on knowledge and treatment decision making for breast cancer surgery: a randomized trial. *JAMA* 2004;**292**(4):435–41.

**Williams 2013** {published and unpublished data}

Williams RM, Davis KM, Luta G, Edmond SN, Dorfman CS, Schwartz MD, et al. Fostering informed decisions: A randomized controlled trial assessing the impact of a decision aid among men registered to undergo mass screening for prostate cancer. *Patient Education and Counseling* 2013;**91**:329–36.

**Wolf 1996** {published data only}

\* Wolf AM, Nasser JF, Wolf AM, Schorling JB. The impact of informed consent on patient interest in prostate-specific antigen screening. *Archives of Internal Medicine* 1996;**156**(12):1333–6.

Wolf AM, Schorling JB. Preferences of elderly men for prostate-specific antigen screening and the impact of informed consent. *Journals of Gerontology Series A-Biological Sciences & Medical Sciences* 1998;**53**(3):M195–200.



**Wolf 2000 {published and unpublished data}**

Wolf AM, Schorling JB. Does informed consent alter elderly patients' preferences for colorectal cancer screening? Results of a randomized trial. *Journal of General Internal Medicine* 2000;**15**(1):24–30.

**Wong 2006 {published data only}**

Wong SS, Thornton JG, Gbolade B, Bekker HL. A randomised controlled trial of a decision-aid leaflet to facilitate women's choice between pregnancy termination methods. *BJOG: An International Journal of Obstetrics & Gynaecology* 2006;**113**(6):688–94.

**References to studies excluded from this review**

**Abadie 2009 {published data only}**

Abadie R, Weymiller AJ, Tilburt J, Shah ND, Charles C, Gafni A, et al. Clinician's use of the Statin Choice decision aid in patients with diabetes: a videographic study nested in a randomized trial. *Journal of Evaluation in Clinical Practice* 2009;**15**(3):492–7.

**Adab 2003 {published data only}**

Adab P, Marshall T, Rouse A, Randhawa B, Sangha H, Bhangoo N. Randomised controlled trial of the effect of evidence based information on women's willingness to participate in cervical cancer screening. *Journal of Epidemiology & Community Health* 2003;**57**(8):589–93.

**Alegría 2014 {published data only}**

Alegría M, Carson N, Flores M, Li X, Shi P, Lessios AS, et al. Activation, self-management, engagement, and retention in behavioral health care: a randomized clinical trial of the DECIDE intervention. *JAMA Psychiatry* 2014;**71**(5):557–65.

**Al Saffar 2008 {published data only}**

Al Saffar N, Abdulkareem A, Abdulhakeem A, Salah AQ, Heba M. Depressed patients' preferences for education about medications by pharmacists in Kuwait. *Patient Education and Counseling* 2008;**72**(1):94–101.

**Altiner 2007 {published data only}**

Altiner A, Brockmann S, Sielk M, Wilm S, Wegscheider K, Abholz HH. Reducing antibiotic prescriptions for acute cough by motivating GPs to change their attitudes to communication and empowering patients: a cluster-randomized intervention study. *Journal of Antimicrobial Chemotherapy* 2007;**60**(3):638–44.

**Anderson 2011 {published data only}**

Anderson C, Carter J, Nattress K, Beale P, Philp S, Harrison J, et al. "The booklet helped me not to panic": a pilot of a decision aid for asymptomatic women with ovarian cancer and with rising CA-125 levels. *International Journal of Gynecological Cancer* 2011;**21**(4):737–43.

**Arimori 2006 {published data only}**

Arimori N. Randomized controlled trial of decision aids for women considering prenatal testing: the effect of the Ottawa Personal Decision Guide on decisional conflict. *Japan Journal of Nursing Science* 2006;**3**(2):119–30.

**Armstrong 2005 {published data only}**

Armstrong K, Weber B, Ubel PA, Peters N, Holmes J, Schwartz JS. Individualized survival curves improve satisfaction with cancer risk management decisions in women with BRCA1/2 mutations. *Journal of Clinical Oncology* 2005;**23**(36):9319–28.

**Arterburn 2013 {published data only}**

Arterburn D, Flum DR, Westbrook EO, Fuller S, Shea M, Bock SN, Landers J, Kowalski K, Turnbull E, Cummings DE, CROSSROADS Study Team. A population-based, shared decision-making approach to recruit for a randomized trial of bariatric surgery versus lifestyle for type 2 diabetes. *Surgery for Obesity and Related Diseases* 2013;**9**(6):837–44.

**Au 2011 {published data only}**

Au AH, Lam WW, Chan MC, Or AY, Kwong A, Suen D, et al. Development and pilot-testing of a decision aid for use among Chinese women facing breast cancer surgery. *Health Expectations* 2011;**14**(4):405–16.

**Bakken 2014 {published data only}**

Bakken S, Jia H, Chen ES, Choi J, John RM, Lee NJ, et al. The effect of a mobile health decision support system on diagnosis and management of obesity, tobacco use, and depression in adults and children. *The Journal for Nurse Practitioners* 2014;**10**(10):774–80.

**Becker 2009 {published data only}**

Becker H, Stuijbergen AK, Dormire SL. The effects of hormone therapy decision support for women with mobility impairments. *Health Care for Women International* 2009;**30**(9):845–54.

**Belkora 2012 {published data only}**

Belkora J, Stupar L, O'Donnell S, Loucks A, Moore D, Jupiter C, et al. Decision support by telephone: randomized controlled trial in a rural community setting. *Patient Education and Counseling* 2012;**89**(1):134–42.

**Bellmunt 2010 {published data only}**

Bellmunt J, Eisen T, Szczylik C, Mulders P, Porta C. A new patient-focused approach to the treatment of metastatic renal cell carcinoma: establishing customized treatment options. *BJU International* 2010;**107**(8):1190–9.

**Bennett 2011 {published data only}**

Bennett PA. Making the choice: cesarean delivery by maternal request versus planned vaginal birth [PhD thesis]. University of Colorado at Denver. ProQuest. Ann Arbor: University of Colorado at Denver, 2011.

**Bieber 2006 {published data only}**

Bieber C, Muller KG, Blumenstiel K, Eich W. Participative decision-making as a measure to improve the doctor-patient interaction with fibromyalgia patients [Partizipative Entscheidungsfindung als Maßnahme zur Verbesserung der Arzt–Patient–Interaktion mit Fibromyalgie–Patientinnen]. *Zeitschrift für Medizinische Psychologie* 2006;**15**(2):53–60. Bieber C, Muller KG, Blumenstiel K, Hochlehnert A, Wilke S, Hartmann M, et al. A shared decision-making communication training program for physicians treating

- fibromyalgia patients: effects of a randomized controlled trial. *Journal of Psychosomatic Research* 2008;**64**(1):13–20.
- Branda 2013** *{published data only}*  
Branda ME, LeBlanc A, Shah ND, Tiedje K, Ruud K, Van Houten H, et al. Shared decision making for patients with type 2 diabetes: a randomized trial in primary care. *BMC Health Services Research* 2013;**13**(301):1–10.
- Brenner 2014** *{published data only}*  
Brenner A, Howard K, Lewis C, Sheridan S, Crutchfield T, Hawley S, et al. Comparing 3 values clarification methods for colorectal cancer screening decision-making: a randomized trial in the US and Australia. *Journal of General Internal Medicine* 2014;**29**(3):507–13.
- Breslin 2008** *{published data only}*  
Breslin M, Mullan RJ, Montori VM. The design of a decision aid about diabetes medications for use during the consultation with patients with type 2 diabetes. *Patient Education and Counseling* 2008;**73**(3):465–72.
- Brown 2004** *{published data only}*  
Brown RF, Butow PN, Sharrock MA, Henman M, Boyle F, Goldstein D, et al. Education and role modelling for clinical decisions with female cancer patients. *Health Expectations* 2004;**7**(4):303–16.
- Brundage 2001** *{published data only}*  
Brundage MD, Feldman-Stewart D, Cosby R, Gregg R, Dixon P, Youssef Y, et al. Phase I study of a decision aid for patients with locally advanced non-small-cell lung cancer. *Journal of Clinical Oncology* 2001;**19**(5):1326–35.
- Burton 2007** *{published data only}*  
Burton MJ. Booklet-based education in vestibular rehabilitation or symptom control improved subjective health in Meniere disease. *Evidence-Based Medicine* 2007;**12**(4):111.
- Buzhardt 2011** *{published data only}*  
Buzhardt J, Greenwood CR, Walker D, Anderson R, Howard W, Carta JJ. Effects of web-based support on early head start home visitors' use of evidence-based intervention decision making and growth in children's expressive communication. *NHSA Dialog* 2011;**14**(3):121–46.
- Campbell 2014** *{published data only}*  
Campbell SR, Holter MC, Manthey TJ, Rapp CA. The effect of CommonGround Software and Decision Support Center. *American Journal of Psychiatric Rehabilitation* 2014;**17**(2):166–80.
- Carling 2008** *{published data only}*  
Carling C, Kristoffersen DT, Herrin J, Treweek S, Oxman AD, Schunemann H, et al. How should the impact of different presentations of treatment effects on patient choice be evaluated? A pilot randomized trial. *PLOS ONE* 2008;**3**(11):e3693.
- Causarano 2015** *{published data only}*  
Causarano N, Platt J, Baxter NN, Bagher S, Jones JM, Metcalfe KA, Hofer SOP, O'Neill AC, Cheng T, Starenkyj E, Zhong T. Pre-consultation educational group intervention to improve shared decision-making for postmastectomy breast reconstruction: a pilot randomized controlled trial. *Support Cancer Care* 2015;**23**:1365–1375.
- Chadwick 1991** *{published data only}*  
Chadwick DJ, Gillatt DA, Gingell JC. Medical or surgical orchidectomy: the patients' choice. *BMJ* 1991;**302**(6776):572.
- Chan 2011** *{published data only}*  
Chan EC, McFall SL, Byrd TL, Mullen PD, Volk RJ, Ureda J, et al. A community-based intervention to promote informed decision making for prostate cancer screening among Hispanic American men changed knowledge and role preferences: a cluster RCT. *Patient Education and Counseling* 2011;**84**(2):e44–51.
- Chewning 1999** *{published data only}*  
Chewning B, Mosena P, Wilson D, Erdman H, Potthoff S, Murphy A, et al. Evaluation of a computerized contraceptive decision aid for adolescent patients. *Patient Education and Counseling* 1999;**38**(3):227–39.
- Chiew 2008** *{published data only}*  
Chiew KS, Shepherd H, Vardy J, Tattersall MHN, Butow PN, Leighl NB. Development and evaluation of a decision aid for patients considering first-line chemotherapy for metastatic breast cancer. *Health Expectations* 2008;**11**(1):35–45.
- Clouston 2014** *{published data only}*  
Clouston K, Katz A, Martens PJ, Sisler J, Turner D, Lobchuk M, et al. CIHR/CCMB Team in Primary Care Oncology (PCO-NET). Does access to a colorectal cancer screening website and/or a nurse-managed telephone help line provided to patients by their family physician increase fecal occult blood test uptake? Results from a pragmatic cluster randomized controlled trial. *BMC Cancer* 2014;**14**:263.
- Col 2007** *{published data only}*  
Col NF, Ngo L, Fortin JM, Goldberg RJ, O'Connor AM. Can computerized decision support help patients make complex treatment decisions? A randomized controlled trial of an individualized menopause decision aid. *Medical Decision Making* 2007;**27**(5):585–98.
- Colella 2004** *{published data only}*  
Colella KM, DeLuca G. Shared decision making in patients with newly diagnosed prostate cancer: a model for treatment education and support. *Urologic Nursing* 2004;**24**(3):187–91, 195–6.
- Costanza 2011** *{published data only}*  
Costanza ME, Luckmann RS, Rosal M, White MJ, LaPelle N, Partin M, et al. Helping men make an informed decision about prostate cancer screening: a pilot study of telephone counseling. *Patient Education and Counseling* 2011;**82**(2):193–200.
- Coulter 2003** *{published data only}*  
Coulter A. Patient information and shared decision-making in cancer care. *British Journal of Cancer* 2003;**89**(Suppl 1):S15–6.

- Cox 2012** *{published data only}*  
Cox CE, Lewis CL, Hanson LC, Hough CL, Kahn JM, White DB, et al. Development and pilot testing of a decision aid for surrogates of patients with prolonged mechanical ventilation. *Critical Care Medicine* 2012;**40**(8): 2327–34.
- Crang-Svalenius 1996** *{published data only}*  
Crang-Svalenius E, Dykes AK, Jorgensen C. Women's informed choice of prenatal diagnosis: early ultrasound examination-routine ultrasound examination-age-independent amniocentesis. *Fetal Diagnosis & Therapy* 1996;**11**(1):20–5.
- Davison 1999** *{published data only}*  
Davison BJ, Kirk P, Degner LE, Hassard TH. Information and patient participation in screening for prostate cancer. *Patient Education and Counseling* 1999;**37**(3):255–63.
- Davison 2007** *{published data only}*  
Davison BJ, Goldenberg SL, Wiens KP, Gleave ME. Comparing a generic and individualized information decision support intervention for men newly diagnosed with localized prostate cancer. *Cancer Nursing* 2007;**30**(5): E7–15.
- De Boer 2012** *{published data only}*  
De Boer JC, van Blijderveen G, van Dijk G, Duivenvoorden HJ, Williams M. Implementing structured, multiprofessional medical ethical decision-making in a neonatal intensive care unit. *Journal of Medical Ethics* 2012;**38**(10):596–601.
- Deen 2012** *{published data only}*  
Deen D, Lu WH, Weintraub MR, Maranda MJ, Elshafey S, Gold MR. The impact of different modalities for activating patients in a community health center setting. *Patient Education and Counseling* 2012;**89**(1):178–83.
- De Haan 2013** *{published data only}*  
De Haan MC, de Wijkerslooth TR, Stoop E, Bossuyt P, Fockens P, Thomeer M, et al. Informed decision-making in colorectal cancer screening using colonoscopy or CT-colonography. *Patient Education and Counseling* 2013;**91**(3):318–25.
- Deinzer 2009** *{published data only}*  
Deinzer A, Veelken R, Kohnen R, Schmieder RE. Is a shared decision-making approach effective in improving hypertension management?. *Journal of Clinical Hypertension* 2009;**11**(5):266–70.
- Denig 2014** *{published data only}*  
Denig P, Schuling J, Haaijer-Ruskamp F, Voorham J. Effects of a patient oriented decision aid for prioritising treatment goals in diabetes: pragmatic randomised controlled trial. *BMJ* 2014;**349**:g5651.
- Deschamps 2004** *{published and unpublished data}*  
Deschamps MA, Taylor JG, Neubauer SL, Whiting S, Green K. Impact of pharmacist consultation versus a decision aid on decision making regarding hormone replacement therapy. *International Journal of Pharmacy Practice* 2004;**12**(1):21–8.
- Deyo 2000** *{published and unpublished data}*  
\* Deyo RA, Cherkin DC, Weinstein J, Howe J, Ciol M, Mulley AG. Involving patients in clinical decisions: impact of an interactive video program on use of back surgery. *Medical Care* 2000;**38**(9):959–69.  
Phelan EA, Deyo RA, Cherkin DC, Weinstein JN, Ciol MA, Kreuter W, et al. Helping patients decide about back surgery: a randomized trial of an interactive video program. *Spine* 2001;**26**(2):206–12.
- Diefenbach 2012** *{published data only}*  
Diefenbach MA, Mohamed NE, Butz BP, Bar-Chama N, Stock R, Cesaretti J, et al. Acceptability and preliminary feasibility of an internet/CD-ROM-based education and decision program for early-stage prostate cancer patients: randomized pilot study. *Journal of Medical Internet Research* 2012;**14**(1):e6.
- Dobke 2008** *{published data only}*  
Dobke MK, Bhavsar D. Pilot trial of telemedicine as a decision aid for patients with chronic wounds. *Telemedicine Journal and e-health* 2008;**14**(3):245–9.
- Dodin 2001** *{published and unpublished data}*  
Dodin S, Legare F, Daudelin G, Tetroe J, O'Connor A. Making a decision about hormone replacement therapy. A randomized controlled trial [Prise de decision en matière d'hormonothérapie de remplacement]. *Canadian Family Physician* 2001;**47**:1586–93.
- Donovan 2012** *{published data only}*  
Donovan JL. Presenting treatment options to men with clinically localized prostate cancer: the acceptability of active surveillance/monitoring. *Journal of the National Cancer Institute. Monographs* 2012;**45**:191–6.
- Driscoll 2008** *{published data only}*  
Driscoll DL, Rupert DJ, Golin CE, McCormack LA, Sheridan SL, Welch BM. Promoting prostate-specific antigen informed decision-making. Evaluating two community-level interventions. *American Journal of Preventive Medicine* 2008;**35**(2):87–94.
- Dunn 1998** *{published and unpublished data}*  
Dunn RA, Shenouda PE, Martin DR, Schultz AJ. Videotape increases parent knowledge about poliovirus vaccines and choices of polio vaccination schedules. *Pediatrics* 1998;**102**(2):e26.
- Eaton 2011** *{published data only}*  
Eaton L, Cherry C, Cain D, Pope H. A novel approach to prevention for at-risk HIV negative men who have sex with men: creating a teachable moment to promote informed sexual decision making. *American Journal of Public Health* 2011;**101**(3):539–45.
- Eden 2009** *{published data only}*  
Eden KB, Dolan JG, Perrin NA, Kocaoglu D, Anderson N, Case J, et al. Patients were more consistent in randomized trial at prioritizing childbirth preferences using graphic-numeric than verbal formats. *Journal of Clinical Epidemiology* 2009;**62**(4):415–24.

**Eden 2014 {published data only}**

Eden KB, Perrin NA, Vesco KK, Guise JM. A randomized comparative trial of two decision tools for pregnant women with prior cesareans. *Journal of Obstetric, Gynecologic, & Neonatal Nursing* 2014;**43**:568–79.

**Eden 2015 {published data only}**

Eden KB, Perrin NA, Hanson GC, Messing JT, Bloom TL, Campbell JC, et al. Use of online safety decision aid by abused women. *American Journal of Preventive Medicine* 2015;**48**(4):372–83.

**Edwards 2012 {published data only}**

Edwards JA, Snyder FJ, Allen PM, Makinson KA, Hamby DM. Decision making for risk management: a comparison of graphical methods for presenting quantitative uncertainty. *Risk Analysis* 2012;**32**(12):2055–70.

**El-Jawahri 2010 {published data only}**

El-Jawahri A, Podgurski LM, Eichler AF, Plotkin SR, Temel JS, Mitchell SL. Use of video to facilitate end-of-life discussions with patients with cancer: a randomized controlled trial. *Journal of Clinical Oncology* 2010;**28**(2):305–10.

**Ellison 2008 {published data only}**

Ellison GL, Weinrich SP. A randomized trial comparing web-based decision aids on prostate cancer knowledge for African-American men. *Journal of the National Medical Association* 2008;**100**(10):1139–45.

**Elwyn 2004 {published data only}**

Elwyn G, Edwards A, Hood K, Robling M, Atwell C, Russell I, et al. Achieving involvement: process outcomes from a cluster randomized trial of shared decision making skill development and use of risk communication aids in general practice. *Family Practice* 2004;**21**(4):337–46.

**Emery 2007 {published data only}**

Emery J, Morris H, Goodchild R, Fanshawe T, Prevost AT, Bobrow M, et al. The GRAIDS Trial: a cluster randomised controlled trial of computer decision support for the management of familial cancer risk in primary care. *British Journal of Cancer* 2007;**97**(4):486–93.

**Emmett 2007 {published data only}**

Emmett CL, Murphy DJ, Patel RR, Fahey T, Jones C, Ricketts IW, et al. Decision-making about mode of delivery after previous caesarean section: development and piloting of two computer-based decision aids. *Health Expectations* 2007;**10**(2):161–72.

**Feldman-Stewart 2006 {published data only}**

Feldman-Stewart D, Brennenstuhl S, Brundage MD, Roques T. An explicit values clarification task: development and validation. *Patient Education and Counseling* 2006;**63**(3):350–6.

**Feldman-Stewart 2012 {published data only}**

Feldman-Stewart D, Tong C, Siemens R, Alibhai S, Pickles T, Robinson J, Brundage MD. The impact of explicit values clarification exercises in a patient decision aid emerges after the decision is actually made: evidence from a randomized controlled trial. *Medical Decision Making* 2012;**32**(4):616–26.

**Fiks 2013a {published data only}**

Fiks AG, Grundmeier RW, Mayne S, Song L, Feemster K, Karavite D, et al. Effectiveness of decision support for families, clinicians, or both on HPV vaccine receipt. *Pediatrics* 2013;**131**(6):1114–24.

**Flood 1996 {published data only}**

Flood AB, Wennberg JE, Nease RF Jr, Fowler FJ Jr, Ding J, Hynes LM. The importance of patient preference in the decision to screen for prostate cancer. Prostate Patient Outcomes Research Team. *Journal of General Internal Medicine* 1996;**11**(6):342–9.

**Francis 2009 {published data only}**

Francis NA, Butler CC, Hood K, Simpson S, Wood F, Nuttall J. Effect of using an interactive booklet about childhood respiratory tract infections in primary care consultations on reconsulting and antibiotic prescribing: a cluster randomised controlled trial. *BMJ* 2009;**339**:b2885.

**Fraval 2015 {published data only}**

Fraval A, Chandrananth J, Chong YM, Tran P, Coventry LS. Internet based patient education improves informed consent for elective orthopaedic surgery: a randomized controlled trial. *BMC Musculoskeletal Disorders* 2015;**16**:14.

**Frosch 2001 {published data only}**

Frosch DL, Kaplan RM, Felitti V. Evaluation of two methods to facilitate shared decision making for men considering the prostate-specific antigen test. *Journal of General Internal Medicine* 2001;**16**(6):391–8.

**Frosch 2003 {published data only}**

Frosch DL, Kaplan RM, Felitti VJ. A randomized controlled trial comparing internet and video to facilitate patient education for men considering the prostate specific antigen test. *Journal of General Internal Medicine* 2003;**18**(10):781–7.

**Frosch 2008b {published data only}**

Frosch DL, Legare F, Mangione CM. Using decision aids in community-based primary care: a theory-driven evaluation with ethnically diverse patients. *Patient Education and Counseling* 2008;**73**(3):490–6.

**Frosch 2011 {published data only}**

Frosch DL, Uy V, Ochoa S, Mangione CM. Evaluation of a behavior support intervention for patients with poorly controlled diabetes. *Archives of Internal Medicine* 2011;**171**(22):2011–7.

**Frost 2009 {published data only}**

Frost J, Shaw A, Montgomery A, Murphy DJ. Women's views on the use of decision aids for decision making about the method of delivery following a previous caesarean section: qualitative interview study. *BJOG: An International Journal of Obstetrics & Gynaecology* 2009;**116**(7):896–905.

**Fujiwara 2015 {published data only}**

Fujiwara H, Shimoda A, Ishikawa Y, Taneichi A, Ohashi M, Takahashi Y, et al. Effect of providing risk information on undergoing cervical cancer screening: a randomized controlled trial. *Archives of Public Health* 2015;**73**:7.

**Garvelink 2013 {published data only}**

Garvelink MM, ter Kuile MM, Fischer MJ, Louwé LA, Hilders CG, Kroep JR, et al. Development of a decision aid about fertility preservation for women with breast cancer in the Netherlands. *Journal of Psychosomatic Obstetrics and Gynecology* 2013;**34**(4):170–8.

**Genz 2012 {published data only}**

Genz J, Haastert B, Müller H, Verheyen F, Cole D, Rathmann W, et al. Blood glucose testing and primary prevention of Type 2 diabetes - evaluation of the effect of evidence-based patient information: a randomized controlled trial. *Diabetic Medicine* 2012;**29**(8):1011–20.

**Giordano 2014 {published data only}**

Giordano A, Lugaresi A, Confalonieri P, Granella F, Radice D, Trojano M, et al. Implementation of the “Sapere Migliora” information aid for newly diagnosed people with multiple sclerosis in routine clinical practice: a late-phase controlled trial. *Multiple Sclerosis Journal* 2014;**20**(9):1234–43.

**Goel 2001 {published and unpublished data}**

Goel V, Sawka CA, Thiel EC, Gort EH, O'Connor AM. Randomized trial of a patient decision aid for choice of surgical treatment for breast cancer. *Medical Decision Making* 2001;**21**(1):1–6.

**Graham 2000 {published data only}**

Graham W, Smith P, Kamal A, Fitzmaurice A, Smith N, Hamilton N. Randomised controlled trial comparing effectiveness of touch screen system with leaflet for providing women with information on prenatal tests. *BMJ* 2000;**320**(7228):155–60.

**Gray 2009 {published data only}**

Gray SW, O'Grady C, Karp L, Smith D, Schwartz JS, Hornik RC, et al. Risk information exposure and direct-to-consumer genetic testing for BRCA mutations among women with a personal or family history of breast or ovarian cancer. *Cancer Epidemiology, Biomarkers & Prevention* 2009;**18**(4):1303–11.

**Green 2001b {published data only}**

Green MJ, McInerney AM, Biesecker BB, Fost N. Education about genetic testing for breast cancer susceptibility: patient preferences for a computer program or genetic counselor. *American Journal of Medical Genetics* 2001;**103**(1):24–31.

**Green 2004 {published data only}**

Green MJ, Peterson SK, Baker MW, Friedman LC, Harper GR, Rubinstein WS, et al. Use of an educational computer program before genetic counseling for breast cancer susceptibility: effects on duration and content of counseling sessions. *Genetics in Medicine* 2005;**7**(4):221–9.

\* Green MJ, Peterson SK, Baker MW, Harper GR, Friedman LC, Rubinstein WS, et al. Effect of a computer-based decision aid on knowledge, perceptions, and intentions about genetic testing for breast cancer susceptibility: a randomized controlled trial. *JAMA* 2004;**292**(4):442–52.

**Greenfield 1985 {published data only}**

Greenfield S, Kaplan S, Ware JE Jr. Expanding patient involvement in care. Effects on patient outcomes. *Annals of Internal Medicine* 1985;**102**(4):520–8.

**Griffith 2008a {published data only}**

Griffith JM, Lewis CL, Brenner AR, Pignone MP. The effect of offering different numbers of colorectal cancer screening test options in a decision aid: a pilot randomized trial. *BMC Medical Informatics and Decision Making* 2008;**8**:4.

**Griffith 2008b {published data only}**

Griffith JM, Fichter M, Fowler FJ, Lewis C, Pignone MP. Should a colon cancer screening decision aid include the option of no testing? A comparative trial of two decision aids. *BMC Medical Informatics and Decision Making* 2008;**8**:10.

**Gruppen 1994 {published data only}**

Gruppen LD, Margolin J, Wisdom K, Grum CM. Outcome bias and cognitive dissonance in evaluating treatment decisions. *Academic Medicine* 1994;**69**(10 Suppl):S57–9.

**Gummersbach 2015 {published data only}**

Gummersbach E, in der Schmitt J, Mortsiefer A, Abholz HH, Wegscheider K, Pentzek M. Willingness to participate in mammography screening - a randomized controlled questionnaire study of responses to two patient information leaflets with different factual content. *Deutsches Ärzteblatt International* 2015;**112**(5):61–8.

**Hacking 2013 {published data only}**

Hacking B, Wallace L, Scott S, Kosmala-Anderson J, Belkora J, McNeill A. Testing the feasibility, acceptability and effectiveness of a 'decision navigation' intervention for early stage prostate cancer patients in Scotland - a randomised controlled trial. *Psycho-Oncology* 2013;**22**(5):1017–1024.

**Hall 2007 {published data only}**

Hall S, Chitty L, Dormandy E, Hollywood A, Wildschut HIJ, Fortuny A, et al. Undergoing prenatal screening for Down's syndrome: presentation of choice and information in Europe and Asia. *European Journal of Human Genetics* 2007;**15**(5):563–9.

**Hall 2011 {published data only}**

Hall MJ, Manne SL, Winkel G, Chung DS, Weinberg DS, Meropol NJ. Effects of a decision support intervention on decisional conflict associated with microsatellite instability testing. *Cancer Epidemiology, Biomarkers and Prevention* 2011;**20**(2):249–54.

**Hamann 2014 {published data only}**

Hamann J, Maris N, Iosifidou P, Mendel R, Cohen R, Wolf P, Kissling W. Effects of a question prompt sheet on active patient behaviour: a randomized controlled trial with depressed outpatients. *International Journal of Social Psychiatry* 2014;**60**(3):227–35.

**Harmsen 2014 {published data only}**

Harmsen CG, Kristiansen IS, Larsen PV, Nexøe J, Støvring H, Gyrd-Hansen D, et al. Communicating risk using absolute risk reduction or prolongation of life formats:

- cluster-randomised trial in general practice. *British Journal of General Practice* 2014;**64**(621):e199–207.
- Harwood 2011** *{published data only}*  
Harwood R, Douglas C, Clark D. Decision aids for breast and nodal surgery in patients with early breast cancer: development and a pilot study. *Asia-Pacific Journal of Clinical Oncology* 2011;**7**:114–22.
- Healton 1999** *{published data only}*  
Healton C, Taylor S, Messeri P, Weinberg G, Bamji M. Effects of ZDV-based patient education on intentions toward ZDV use, HIV testing and reproduction among a US cohort of women. *AIDS Care* 1999;**11**(6):675–86.
- Henderson 2013** *{published data only}*  
Henderson C, Brohan E, Clement S, Williams P, Lassman F, Schauman O, et al. Decision aid on disclosure of mental health status to an employer: feasibility and outcomes of a randomised controlled trial. *British Journal of Psychiatry* 2013;**203**(5):350–7.
- Herrera 1983** *{published data only}*  
Herrera AJ, Cochran B, Herrera A, Wallace B. Parental information and circumcision in highly motivated couples with higher education. *Pediatrics* 1983;**71**(2):233–4.
- Hess 2015** *{published data only}*  
Hess LM, Litwiller A, Byron J, Stutsman J, Kasper K, Learman LA. Preference elicitation tool for abnormal uterine bleeding treatment: a randomized controlled trial. *The Patient: Patient Centered Outcomes Research* 2015;**8**(2): 217–27.
- Hewison 2001** *{published data only}*  
Hewison J, Cuckle H, Baillie C, Sehmi I, Lindow S, Jackson F, et al. Use of videotapes for viewing at home to inform choice in Down syndrome screening: a randomised controlled trial. *Prenatal Diagnosis* 2001;**21**(2):146–9.
- Heyn 2013** *{published data only}*  
Heyn L, Finset A, Eide H, Ruland CM. Effects of an interactive tailored patient assessment on patient-clinician communication in cancer care. *Psycho-Oncology* 2013;**22** (1):89–96.
- Hickish 1995** *{published data only}*  
Hickish TF, Smith IE, Middleton G, Nicolson M. Patient preference for extended palliative chemotherapy for non-small cell lung cancer. *Lancet* 1995;**345**(8953):857–8.
- Hochlehnert 2006** *{published data only}*  
Hochlehnert A, Richter A, Bludau HB, Bieber C, Blumenstiel K, Mueller K, et al. A computer-based information-tool for chronic pain patients: computerized information to support the process of shared decision-making. *Patient Education and Counseling* 2006;**61**(1): 92–8.
- Hofbauer 2008** *{published data only}*  
Hofbauer GFL, Buhler RPN, French LE, Brockes M, Scheuer E. Patient-centered care in dermatology: an online system that provides accessible and appropriate information to guide patients' decision making. *Archives of Dermatology* 2008;**144**(9):1225–7.
- Hoffman 2009** *{published data only}*  
Hoffman RM, Walter LC. Colorectal cancer screening in the elderly: the need for informed decision making. *Journal of General Internal Medicine* 2009;**24**(12):1336–7.
- Holbrook 2007** *{published data only}*  
Holbrook A, Labiris R, Goldsmith CH, Ota K, Harb S, Sebaldt RJ. Influence of decision aids on patient preferences for anticoagulant therapy: a randomized trial. *CMAJ* 2007; **176**(11):1583–7.
- Hollen 2013** *{published data only}*  
Hollen PJ, Tyc VL, Donnangelo SF, Shannon SV, O'Laughlen MC, Hinton I, et al. A substance use decision aid for medically at-risk adolescents: results of a randomized controlled trial for cancer-surviving adolescents. *Cancer Nursing* 2013;**36**(5):355–67.
- Holloway 2003** *{published data only}*  
Holloway RM, Wilkinson C, Peters TJ, Russell I, Cohen D, Hale J, et al. Cluster-randomised trial of risk communication to enhance informed uptake of cervical screening. *British Journal of General Practice* 2003;**53**(493): 620–5.
- Holmes-Rovner 2011** *{published data only}*  
Holmes-Rovner M, Kelly-Blake K, Dwamena F, Dontje K, Henry R, Olomu A, et al. Shared decision making guidance reminders in practice (SDM-GRIP). *Patient Education and Counseling* 2011;**85**(2):219–24.
- Holt 2009** *{published data only}*  
Holt CL, Wynn TA, Litaker MS, Southward P, James S, Schulz E. A comparison of a spiritually based and non-spiritually based educational intervention for informed decision making for prostate cancer screening among church-attending African-American men. *Urologic Nursing* 2009;**29**(4):249–58.
- Hope 2010** *{published data only}*  
Hope N, Rombauts L. Can an educational DVD improve the acceptability of elective single embryo transfer? A randomized controlled study. *Fertility and Sterility* 2010;**94** (2):489–95.
- Huijbregts 2013** *{published data only}*  
Huijbregts KML, de Jong FJ, van Marwijk HWJ, Beekman ATF, Adèr HJ, Hakkaart-van Roijen L, et al. A target-driven collaborative care model for major depressive disorder is effective in primary care in the Netherlands. A randomized clinical trial from the depression initiative. *Journal of Affective Disorders* 2013;**146**:328–37.
- Hunt 2005** *{published data only}*  
Hunt LM, de Voogd KB, Castaneda H. The routine and the traumatic in prenatal genetic diagnosis: does clinical information inform patient decision-making?. *Patient Education and Counseling* 2005;**56**(3):302–12.
- Hunter 1999** *{published data only}*  
Hunter M, O'Dea I. An evaluation of a health education intervention for mid-aged women: five year follow-up of effects upon knowledge, impact of menopause and health. *Patient Education and Counseling* 1999;**38**(3):249–55.

**Hunter 2005 {published data only}**

Hunter AG, Cappelli M, Humphreys L, Allanson JE, Chiu TT, Peeters C, et al. A randomized trial comparing alternative approaches to prenatal diagnosis counseling in advanced maternal age patients. *Clinical Genetics* 2005;**67**(4):303–13.

**Huyghe 2009 {published data only}**

Huyghe E, Martinetti P, Sui D, Schover LR. Banking on Fatherhood: pilot studies of a computerized educational tool on sperm banking before cancer treatment. *Psycho-Oncology* 2009;**18**(9):1011–4.

**Ilic 2008 {published data only}**

Ilic D, Egberts K, McKenzie JE, Risgrider G, Green S. Informing men about prostate cancer screening: a randomized controlled trial of patient education materials. *Journal of General Internal Medicine* 2008;**23**(4):466–71.

**Isebaert 2007 {published data only}**

Isebaert S, Van Audenhove C, Haustermans K, DeRidder K, Junius S, Joniau S, et al. A decision aid for patients with localized prostate cancer: first results [Een beslissingshulp voor patienten met gelokaliseerde prostaatkanker: eerste resultaten]. *Tijdschrift voor Geneeskunde* 2007;**63**(1):15–21.

**Jackson 2011 {published data only}**

Jackson C, Cheater FM, Harrison W, Peacock R, Bekker H, West R, et al. Randomised cluster trial to support informed parental decision-making for the MMR vaccine. *BMC Public Health* 2011;**11**:475.

**Jerant 2007 {published data only}**

Jerant A, Kravitz RL, Rooney M, Amerson S, Kreuter M, Franks P. Effects of a tailored interactive multimedia computer program on determinants of colorectal cancer screening: a randomized controlled pilot study in physician offices. *Patient Education and Counseling* 2007;**66**(1):67–74.

**Jibaja-Weiss 2006 {published data only}**

Jibaja-Weiss ML, Volk RJ, Granchi TS, Neff NE, Spann SJ, Aoki N, et al. Entertainment education for informed breast cancer treatment decisions in low-literate women: development and initial evaluation of a patient decision aid. *Journal of Cancer Education* 2006;**21**(3):133–9.

**Joosten 2009 {published data only}**

Joosten EA, de Jong CA, de Weert-van Oene GH, Sensky T, van der Staak CP. Shared decision-making reduces drug use and psychiatric severity in substance-dependent patients. *Psychotherapy and Psychosomatics* 2009;**78**:245–53.

**Joosten 2011 {published data only}**

Joosten EA, De Jong CA, de Weert-van Oene GH, Sensky T, van der Staak CP. Shared decision-making: increases autonomy in substance-dependent patients. *Substance Use and Misuse* 2011;**48**:1037–48.

**Jorm 2003 {published data only}**

Jorm AF, Griffiths KM, Christensen H, Korten AE, Parslow RA, Rodgers B. Providing information about the effectiveness of treatment options to depressed people in the community: a randomized controlled trial of effects

on mental health literacy, help-seeking and symptoms. *Psychological Medicine* 2003;**33**(6):1071–9.

**Kakkilaya 2011 {published data only}**

Kakkilaya V, Groome L, Platt D, Kurepa D, Pramanik A, Caldito G, et al. Use of a visual aid to improve counseling at the threshold of viability. *Pediatrics* 2011;**128**(6):e1511–9.

**Kaplan 2014a {published data only}**

Kaplan CP, Livaudais-Toman J, Tice JA, Kerlikowske K, Gregorich SE, Pérez-Stable EJ, et al. A randomized, controlled trial to increase discussion of breast cancer in primary care. *Cancer Epidemiology, Biomarkers & Prevention* 2014;**23**(7):1245–53.

**Kaplan 2014b {published data only}**

Kaplan AL, Crespi CM, Saucedo JD, Connor SE, Litwin MS, Saigal CS. Decisional conflict in economically disadvantaged men with newly diagnosed prostate cancer. *Cancer* 2014;**120**(17):2721–7.

**Kassan 2012 {published data only}**

Kassan EC, Williams RM, Kelly SP, Barry SA, Penek S, Fishman MB, Cole CA, et al. Men's use of an internet-based decision aid for prostate cancer screening. *Journal of Health Communication* 2012;**17**(6):677–97.

**Kellar 2008 {published data only}**

Kellar I, Sutton S, Griffin S, Prevost AT, Kinmonth AL, Marteau TM. Evaluation of an informed choice invitation for type 2 diabetes screening. *Patient Education and Counseling* 2008;**72**(2):232–8.

**Kiatpongsan 2014 {published data only}**

Kiatpongsan S, Carlson K, Feibelmann S, Sepucha K. Decision aid reduces misperceptions about hormone therapy: a randomized controlled trial. *Menopause: The Journal of The North American Menopause Society* 2014;**21**(1):33–38.

**Kobelka 2009 {published data only}**

Kobelka C, Mattman A, Langlois S. An evaluation of the decision-making process regarding amniocentesis following a screen-positive maternal serum screen result. *Prenatal Diagnosis* 2009;**29**(5):514–9.

**Koelewijn-van Loon 2009 {published data only}**

Koelewijn-van Loon MS, van der Weijden T, van Steenkiste B, Ronda G, Winkens B, Severens JL, et al. Involving patients in cardiovascular risk management with nurse-led clinics: a cluster randomized controlled trial. *CMAJ* 2009;**181**(12):E267–74.

**Köpke 2009 {published data only}**

Köpke S, Kasper J, Mühlhauser I, Nübling M, Heesen C. Patient education program to enhance decision autonomy in multiple sclerosis relapse management: a randomized-controlled trial. *Multiple Sclerosis* 2009;**15**(1):96–104.

**Köpke 2014 {published data only}**

Köpke S, Kern S, Ziemssen T, Berghoff M, Kleiter I, Marziniak M, et al. Evidence-based patient information programme in early multiple sclerosis: a randomised controlled trial. *Journal of Neurology, Neurosurgery, and Psychiatry* 2014;**85**(4):411–18.

**Krawczyk 2012 {published data only}**

Krawczyk A. *Cancer Prevention and the Human Papillomavirus Vaccine: Psychosocial and Behavioural Factors Involved in Vaccination Decision-making [PhD thesis]*. Montreal: McGill Library, 2012.

**Kripalani 2007 {published data only}**

Kripalani S, Sharma J, Justice E, Justice J, Spiker C, Laufman LE, et al. Low-literacy interventions to promote discussion of prostate cancer: a randomized controlled trial. *American Journal of Preventive Medicine* 2007;**33**(2):83–90.

**Krones 2008 {published data only}**

Krones T, Keller H, Becker A, Sonnichsen A, Baum E, Donner-Banzhoff N. The theory of planned behaviour in a randomized trial of a decision aid on cardiovascular risk prevention. *Patient Education and Counseling* 2009;**78**(2): 169–76.

Krones T, Keller H, Sönnichsen A, Sadowski EM, Baum E, Wegscheider K, et al. Absolute cardiovascular disease risk and shared decision making in primary care: a randomized controlled trial. *Annals of Family Medicine* 2008;**6**(3): 218–27.

**Kuppermann 2009 {published data only}**

Kuppermann M, Norton ME, Gates E, Gregorich SE, Learman LA, Nakagawa S, et al. Computerized prenatal genetic testing decision-assisting tool: a randomized controlled trial. *Obstetrics & Gynecology* 2009;**113**(1): 53–63.

**Kurian 2009 {published data only}**

Kurian B, Trivedi M, Grannemann B, Claassen C, Daly E, Sunderajan P. A computerized decision support system for depression in primary care. *Primary Care Companion to the Journal of Clinical Psychiatry* 2009;**11**(4):140–6.

**Labrecque 2010 {published data only}**

Labrecque M, Paunescu C, Plesu I, Stacey D, Legare F. Evaluation of the effect of a patient decision aid about vasectomy on the decision-making process: a randomized trial. *Contraception* 2010;**82**(6):556–62.

**LaCroix 1999 {published data only}**

LaCroix AZ, Newton KM, Buist DSM, Curry SJ, Scholes D, Anderson LA, et al. Population-based strategy for improving informed decision making about hormone replacement therapy in managed care settings. *Women's Health Issues* 1999;**9**(6):306–18.

**Lairson 2011 {published data only}**

Lairson DR, Chan W, Chang YC, del Junco DJ, Vernon SW. Cost-effectiveness of targeted versus tailored interventions to promote mammography screening among women military veterans in the United States. *Evaluation and Program Planning* 2011;**34**(2):97–104.

**Lalonde 2006 {published data only}**

Lalonde L, O'Connor AM, Duguay P, Brassard J, Drake E, Grover SA. Evaluation of a decision aid and a personal risk profile in community pharmacy for patients considering options to improve cardiovascular health: the OPTIONS pilot study. *International Journal of Pharmacy Practice* 2006;**14**(1):51–62.

**Lancaster 2009 {published data only}**

Lancaster T. Physician training in the use of a decision aid increased patient participation in decision making for CVD prevention. *Evidence-Based Medicine* 2009;**14**(1):24.

**Landrey 2013 {published data only}**

Landrey AR, Matlock DD, Andrews L, Bronsert M, Denberg T. Shared decision making in prostate-specific antigen testing: the effect of a mailed patient flyer prior to an annual exam. *Journal of Primary Care & Community Health* 2013;**4**(1):67–74.

**Lazcano Ponce 2000 {published data only}**

Lazcano Ponce EC, Sloan NL, Winikoff B, Langer A, Coggins C, Heimbürger A, et al. The power of information and contraceptive choice in a family planning setting in Mexico. *Sexually Transmitted Infections* 2000;**76**(4):277–81.

**Legare 2003 {published data only}**

Legare F, O'Connor AM, Graham ID, Wells GA, Jacobsen MJ, Elmslie T, et al. The effect of decision aids on the agreement between women's and physicians' decisional conflict about hormone replacement therapy. *Patient Education and Counseling* 2003;**50**(2):211–21.

**Leung 2004 {published data only}**

Leung KY, Lee CP, Chan HY, Tang MH, Lam YH, Lee A. Randomised trial comparing an interactive multimedia decision aid with a leaflet and a video to give information about prenatal screening for Down syndrome. *Prenatal Diagnosis* 2004;**24**(8):613–8.

**Levin 2011 {published data only}**

Levin W, Campbell D, McGovern K, Gau J, Kosty D, Seeley J, Lewinsohn P. A computer-assisted depression intervention in primary care. *Psychological Medicine* 2011;**41**(7):1373–83.

**Lewis 2003 {published data only}**

Lewis CL, Pignone MP, Sheridan SL, Downs SM, Kinsinger LS. A randomized trial of three videos that differ in the framing of information about mammography in women 40 to 49 years old. *Journal of General Internal Medicine* 2003;**18**(11):875–83.

**Lewis 2012 {published data only}**

Lewis CL, Brenner AT, Griffith JM, Moore CG, Pignone MP. Two controlled trials to determine the effectiveness of a mailed intervention to increase colon cancer screening. *North Carolina Medical Journal* 2012;**73**(2):93–8.

**Lopez-Jornet 2012 {published data only}**

López-Jornet P, Camacho-Alonso F, Sanchez-Siles M. Patient information preferences and behaviour in relation to oral biopsies. *British Journal of Oral & Maxillofacial Surgery* 2012;**50**(8):e115–8.

**Lukens 2013 {published data only}**

Lukens JM, Solomon P, Sorenson SB. Shared decision-making for clients with mental illness: a randomized factorial survey. *Research on Social Work Practice* 2013;**23**(6):694–705.

**Lurie 2011 {published data only}**

Lurie J, Spratt K, Blood E, Tosteson T, Tosteson A, Weinstein J. Effects of viewing an evidence based video



- decision aid on patients' treatment preferences for spine surgery. *Spine* 2011;**36**(18):1501–4.
- Maisels 1983** *{published data only}*  
Maisels MJ, Hayes B, Conrad S, Chez RA. Circumcision: the effect of information on parental decision making. *Pediatrics* 1983;**71**(3):453–5.
- Mancini 2006** *{published data only}*  
Mancini J, Santin G, Chabal F, Julian-Reynier C. Cross-cultural validation of the Decisional Conflict Scale in a sample of French patients. *Quality of Life Research* 2006;**15**(6):1063–8.
- Manne 2009** *{published data only}*  
Manne SL, Coups EJ, Markowitz A, Meropol NJ, Haller D, Jacobsen PB, et al. A randomized trial of generic versus tailored interventions to increase colorectal cancer screening among intermediate risk siblings. *Annals of Behavioral Medicine* 2009;**37**(2):207–17.
- Manns 2005** *{published data only}*  
Manns B J, Taub K, Vanderstraeten C, Jones H, Mills C, Visser M, et al. The impact of education on chronic kidney disease patients' plans to initiate dialysis with self-care dialysis: a randomized trial. *Kidney International* 2005;**68**(4):1777–83.
- Markham 2003** *{published data only}*  
Markham R, Smith A. Limits to patient choice: example from anaesthesia. *BMJ* 2003;**326**(7394):863–4.
- Martin 2012** *{published data only}*  
Martin R, Brower M, Gerald A, Gallagher P, Tellinghuisen D. An experimental evaluation of patient decision aid design to communicate the effects of medications on the rate of progression of structural joint damage in rheumatoid arthritis. *Patient Education and Counseling* 2012;**86**(3):329–34.
- Maslin 1998** *{published data only}*  
\* Maslin AM, Baum M, Walker JS, A'Hern R, Prouse A. Shared decision-making using an interactive video disk system for women with early breast cancer. *NT Research* 1998;**3**(6):444–55.  
Maslin AM, Baum M, Walker JS, A'Hern R, Prouse A. Using an interactive video disk in breast cancer patient support. *Nursing Times* 1998;**94**(44):4–10.
- Matlock 2014** *{published data only}*  
Matlock DD, Keech TA, McKenzie MB, Bronsert MR, Nowels CT, Kutner JS. Feasibility and acceptability of a decision aid designed for people facing advanced or terminal illness: a pilot randomized trial. *Health Expectations* 2014;**17**(1):49–59.
- Matloff 2006** *{published data only}*  
Matloff ET, Moyer A, Shannon KM, Niendorf KB, Col NF. Healthy women with a family history of breast cancer: impact of a tailored genetic counseling intervention on risk perception, knowledge, and menopausal therapy decision making. *Journal of Women's Health* 2006;**15**(7):843–56.
- Mazur 1994** *{published data only}*  
Mazur DJ, Hickam DH. The effect of physician's explanations on patients' treatment preferences: five-year survival data. *Medical Decision Making* 1994;**14**(3):255–8.
- McCaffery 2007** *{published data only}*  
McCaffery K, Irwig L, Bossuyt P. Patient decision aids to support clinical decision making: evaluating the decision or the outcomes of the decision. *Medical Decision Making* 2007;**27**(5):619–25.
- McGinley 2002** *{published data only}*  
McGinley AM. *Effect of Web-based Computer-tailoring on Women's Intention to Continue or Begin to Use Hormone Replacement Therapy to Lower their Risk for Osteoporosis [PhD thesis]*. Philadelphia: University of Pennsylvania, 2002.
- McGowan 2008** *{published data only}*  
McGowan J, Hogg W, Campbell C, Rowan M. Just-in-time information improved decision-making in primary care: a randomized controlled trial. *PLOS ONE* 2008;**3**(11):e3785.
- McInerney-Leo 2004** *{published data only}*  
McInerney-Leo A, Biesecker BB, Hadley DW, Kase RG, Giambresini TR, Johnson E, et al. BRCA1/2 testing in hereditary breast and ovarian cancer families: effectiveness of problem-solving training as a counseling intervention. *American Journal of Medical Genetics. Part A* 2004;**130**(3):221–7.
- McLaren 2012** *{published data only}*  
McLaren PJ, Hyde MK, White KM. Exploring the role of gender and risk perceptions in people's decisions to register as a bone marrow donor. *Health Education Research* 2011;**27**(3):513–22.
- Meropol 2013** *{published data only}*  
Meropol NJ, Egleston BL, Buzaglo JS, Balshem A, Benson AB 3rd, Cegala DJ, et al. A web-based communication aid for patients with cancer: the CONNECT study. *Cancer* 2013;**119**(7):1437–45.
- Michie 1997** *{published data only}*  
Michie S, Smith D, McClelland A, Marteau TM. Patient decision making: An evaluation of two different methods of presenting information about a screening test. *British Journal of Health Psychology* 1997;**2**(4):317–26.
- Miller 2014a** *{published data only}*  
Miller MJ, Allison JJ, Coughlin DJ, Ray MN, Saag KG. A group-randomized trial of shared decision making for non-steroidal anti-inflammatory drug risk awareness: primary results and lessons learned. *Journal of Evaluation in Clinical Practice* 2014;**20**:638–48.
- Miller 2014b** *{published data only}*  
Miller SM, Roussi P, Scarpato J, Wen KY, Zhu F, Roy G. Randomized trial of print messaging: the role of the partner and monitoring style in promoting provider discussions about prostate cancer screening among African American men. *Psycho-Oncology* 2014;**23**:404–11.
- Mishel 2009** *{published data only}*  
Mishel MH, Germino BB, Lin L, Pruthi RS, Wallen EM, Crandell J, et al. Managing uncertainty about treatment

- decision making in early stage prostate cancer: a randomized clinical trial. *Patient Education and Counseling* 2009;**77**(3): 349–59.
- Mohammad 2012** *{published data only}*  
 Mohammad-Alizadeh-Charandabi S, Shahnazi M, Jahanbakhsh. Communicating contraceptive effectiveness: a randomized controlled trial. *Journal of Caring Sciences* 2012;**1**(1):1–9.
- Molenaar 2001** *{published data only}*  
 Molenaar S, Sprangers MA, Rutgers EJ, Luiten EJ, Mulder J, Bossuyt PM, et al. Decision support for patients with early-stage breast cancer: effects of an interactive breast cancer CDROM on treatment decision, satisfaction, and quality of life. *Journal of Clinical Oncology* 2001;**19**(6): 1676–87.
- Mulley 2006** *{published data only}*  
 Mulley AG Jr. Developing skills for evidence-based surgery: ensuring that patients make informed decisions. *Surgical Clinics of North America* 2006;**86**(1):181–92.
- Myers 2005a** *{published data only}*  
 Myers RE, Daskalakis C, Cocroft J, Kunkel EJ, Delmoor E, Liberatore M, et al. Preparing African-American men in community primary care practices to decide whether or not to have prostate cancer screening. *Journal of the National Medical Association* 2005;**97**(8):1143–54.
- Myers 2005b** *{published data only}*  
 Myers RE. Decision counseling in cancer prevention and control. *Health Psychology* 2005;**24**(4 Suppl):S71–7.
- Myers 2007** *{published data only}*  
 Myers RE, Sifri R, Hyslop T, Rosenthal M, Vernon SW, Cocroft J, et al. A randomized controlled trial of the impact of targeted and tailored interventions on colorectal cancer screening. *Cancer* 2007;**110**(9):2083–91.
- Myers 2011** *{published data only}*  
 Myers RE, Daskalakis C, Kunkel EJ, Cocroft JR, Riggio JM, Capkin M, et al. Mediated decision support in prostate cancer screening: a randomized controlled trial of decision counseling. *Patient Education and Counseling* 2011;**83**(2): 240–6.
- Myers 2013** *{published data only}*  
 Myers RE, Bittner-Fagan H, Daskalakis C, Sifri R, Vernon SW, Cocroft J, et al. A randomized controlled trial of a tailored navigation and a standard intervention in colorectal cancer screening. *Cancer Epidemiology, Biomarkers & Prevention* 2013;**22**(1):109–17.
- Neubeck 2008** *{published data only}*  
 Neubeck L, Redfern J, Briffa T, Bauman A, Hare D, Freedman SB. The CHOICE (Choice of Health Options In prevention of Cardiovascular Events) replication trial: study protocol. *BMC Cardiovascular Disorders* 2008;**8**:25.
- Newton 2001** *{published data only}*  
 Newton KM, LaCroix AZ, Buist DS, Delaney KM, Anderson LA. Women's responses to a mailed hormone replacement therapy workbook. *Menopause* 2001;**8**(5): 361–7.
- O'Cathain 2002** *{published data only}*  
 O'Cathain A, Walters SJ, Nicholl JP, Thomas KJ, Kirkham M. Use of evidence based leaflets to promote informed choice in maternity care: randomised controlled trial in everyday practice. *BMJ* 2002;**324**(7338):643–6.
- O'Connor 1996** *{published data only}*  
 O'Connor AM, Pennie RA, Dales RE. Framing effects on expectations, decisions, and side effects experienced: the case of influenza immunization. *Journal of Clinical Epidemiology* 1996;**49**(11):1721–6.
- O'Connor 1998a** *{published and unpublished data}*  
 O'Connor AM, Tugwell P, Wells GA, Elmslie T, Jolly E, Hollingworth G, et al. Randomized trial of a portable, self-administered decision aid for postmenopausal women considering long-term preventive hormone therapy. *Medical Decision Making* 1998;**18**:295–303.
- O'Connor 1999a** *{published data only}*  
 O'Connor AM, Wells GA, Tugwell P, Laupacis A, Elmslie T, Drake E. The effects of an 'explicit' values clarification exercise in a women's decision aid regarding postmenopausal hormone therapy. *Health Expectations* 1999;**2**:21–32.
- O'Connor 2009a** *{published data only}*  
 O'Connor PJ, Sperl-Hillen J, Johnson PE, Rush WA, Crain AL. Customized feedback to patients and providers failed to improve safety or quality of diabetes care: a randomized trial. *Diabetes Care* 2009;**32**(7):1158–63.
- O'Connor 2011** *{published data only}*  
 Connor PJ, Sperl-Hillen JM, Rush WA, Johnson PE, Amundson GH, Asche SE, et al. Impact of electronic health record clinical decision support on diabetes care: a randomized trial. *Annals of Family Medicine* 2011;**9**(1): 12–21.
- Owens 2014A** *{published data only}*  
 Owens OL. *A Community-driven Approach to the Development of a Digital Decision Aid to Facilitate Informed Decision Making for Prostate Cancer Screening among African-American men in Communities of Faith [PhD thesis]*. Columbia: University of South Carolina, 2014.
- Patanwala 2011** *{published data only}*  
 Patanwala IM, Brocklebank V, Inglis J, Trewby PN. A randomized questionnaire-based study on the impact of providing numerical information on colorectal cancer screening. *Journal of the Royal Society of Medicine Short Reports* 2011;**2**(6):48.
- Patel 2014** *{published data only}*  
 Patel S, Ngunjiri A, Wan Hee S, Yang Y, Brown S, Friede T, et al. Primum non nocere: shared informed decision making in low back pain - a pilot cluster randomised trial. *BMC Musculoskeletal Disorders* 2014;**15**:282.
- Pearson 2005** *{published data only}*  
 Pearson S, Maddern GJ, Hewett P. Interacting effects of preoperative information and patient choice in adaptation to colonoscopy. *Diseases of the Colon & Rectum* 2005;**48** (11):2047–54.

- Peele 2005** {published data only}  
Peele PB, Siminoff LA, Xu Y, Ravdin PM. Decreased use of adjuvant breast cancer therapy in a randomized controlled trial of a decision aid with individualized risk information. *Medical Decision Making* 2005;**25**(3):301–7.
- Petty 2014** {published data only}  
Petty J. Exploring the effectiveness of an interactive, technology-enabled learning tool to enhance knowledge for neonatal nurses. *Neonatal, Paediatric and Child Health Nursing* 2014;**17**(1):2–10.
- Philip 2010** {published data only}  
Philip E, DuHamel K, Jandorf L. Evaluating the impact of an educational intervention to increase CRC screening rates in the African American community: a preliminary study. *Cancer Causes Control* 2010;**21**(10):1685–91.
- Phillips 1995** {published data only}  
Phillips C, Hill BJ, Cannac C. The influence of video imaging on patients' perceptions and expectations. *Angle Orthodontist* 1995;**65**(4):263–70.
- Pignone 2013** {published data only}  
Pignone MP, Howard K, Brenner AT, Crutchfield TM, Hawley ST, Lewis CL, et al. Comparing 3 techniques for eliciting patient values for decision making about prostate-specific antigen screening: a randomized controlled trial. *JAMA Internal Medicine* 2013;**173**(5):362–8.
- Pinto 2008** {published data only}  
Pinto H, Rumball D, Maskrey V, Holland R. A pilot study for a randomized controlled and patient preference trial of buprenorphine versus methadone maintenance treatment in the management of opiate dependent patients. *Journal of Substance Use* 2008;**13**(2):73–82.
- Powers 2011** {published data only}  
Powers B, Danus S, Grubber J, Olsen M, Oddone E, Bosworth H. The effectiveness of personalized coronary heart disease and stroke risk communication. *American Heart Journal* 2011;**161**(4):673–80.
- Proctor 2006** {published data only}  
Proctor A, Jenkins TR, Loeb T, Elliot M, Ryan A. Patient satisfaction with 3 methods of postpartum contraceptive counseling: a randomized, prospective trial. *Journal of Reproductive Medicine* 2006;**51**(5):377–82.
- Prunty 2008** {published data only}  
Prunty MC, Sharpe L, Butow P, Fulcher G. The motherhood choice: a decision aid for women with multiple sclerosis. *Patient Education and Counseling* 2008;**71**(1):108–15.
- Ranta 2015** {published data only}  
Ranta A, Dovey S, Weatherall M, O'Dea D, Gommans J, Tilyard M. Cluster randomized controlled trial of TIA electronic decision support in primary care. *American Academy of Neurology* 2015;**84**(15):1545–51.
- Rapley 2006** {published data only}  
Rapley T, May C, Heaven B, Murtagh M, Graham R, Kaner EF, et al. Doctor-patient interaction in a randomised controlled trial of decision-support tools. *Social Science & Medicine* 2006;**62**(9):2267–78.
- Raynes-Greenow 2009** {published data only}  
Raynes-Greenow CH, Roberts CL, Nassar N, Trevena L. Do audio-guided decision aids improve outcomes? A randomized controlled trial of an audio-guided decision aid compared with a booklet decision aid for Australian women considering labour analgesia. *Health Expectations* 2009;**12**(4):407–16.
- Raynes-Greenow 2010** {published data only}  
Raynes-Greenow CH, Nassar N, Torvaldsen S, Trevena L, Roberts CL. Assisting informed decision making for labour analgesia: a randomised controlled trial of a decision aid for labour analgesia versus a pamphlet. *BMC Pregnancy and Childbirth* 2010;**10**:15.
- Rimer 2001** {published data only}  
Rimer BK, Halabi S, Sugg Skinner C, Kaplan EB, Crawford Y, Samsa GP, et al. The short-term impact of tailored mammography decision-making interventions. *Patient Education and Counseling* 2001;**43**(3):269–85.
- Rimer 2002** {published data only}  
Rimer BK, Halabi S, Sugg Skinner C, Lipkus IM, Strigo TS, Kaplan EB, et al. Effects of a mammography decision-making intervention at 12 and 24 months. *American Journal of Preventive Medicine* 2002;**22**(4):247–57.
- Robinson 2013** {published data only}  
Robinson JK, Gaber R, Hultgren B, Eilers S, Blatt H, Stapleton J, et al. Skin self-examination education for early detection of melanoma: a randomized controlled trial of internet, workbook and in-person interventions. *Journal of Medical Internet Research* 2013;**16**(1):1–11.
- Ronda 2014** {published and unpublished data}  
Ronda G, Grispen JEJ, Ickenroth M, Dinant GJ, de Vries NK, van der Weijden T. The effects of a web-based decision aid on the intention to diagnostic self-testing for cholesterol and diabetes: a randomized controlled trial. *BMC Public Health* 2014;**14**:921.
- Rostom 2002** {published data only}  
Rostom A, O'Connor A, Tugwell P, Wells G. A randomized trial of a computerized versus an audio-booklet decision aid for women considering post-menopausal hormone replacement therapy. *Patient Education and Counseling* 2002;**46**(1):67–74.
- Roter 2012** {published data only}  
Roter DL, Wexler R, Naragon P, Forrest B, Dees J, Almodovar A, Wood J. The impact of patient and physician computer mediated communication skill training on reported communication and patient satisfaction. *Patient Education and Counseling* 2012;**88**(3):406–13.
- Rotherth 1997** {published and unpublished data}  
Holmes-Rovner M, Kroll J, Rovner DR, Schmitt N, Rotherth M, Padonu G, et al. Patient decision support intervention: increased consistency with decision analytic models. *Medical Care* 1999;**37**(3):270–84.  
\* Rotherth ML, Holmes-Rovner M, Rovner D, Kroll J, Breer L, Talarczyk G, et al. An educational intervention as decision support for menopausal women. *Research in Nursing & Health* 1997;**20**(5):377–87.

- Rovner 2004** {published data only}  
Rovner DR, Wills CE, Bonham V, Williams G, Lillie J, Kelly-Blake K, et al. Decision aids for benign prostatic hyperplasia: applicability across race and education. *Medical Decision Making* 2004;**24**(4):359–66.
- Rubinstein 2011** {published data only}  
Rubinstein W, Acheson L, O'Neill S, Ruffin M, Wang C, Beaumont J, et al. Clinical utility of family history for cancer screening and referral in primary care: a report from the Family Healthcare Impact Trial. *Genetics in Medicine* 2011;**13**(11):956–65.
- Ruddy 2009** {published data only}  
Ruddy KJ, Partridge AH. Breast cancer in young women: clinical decision-making in the face of uncertainty. *Oncology* 2009;**23**(6):474–7.
- Ruehlman 2012** {published data only}  
Ruehlman LS, Karoly P, Enders C. A randomized controlled evaluation of an online chronic pain self management program. *Pain* 2012;**153**(2):319–30.
- Ruland 2013** {published data only}  
Ruland CM, Andersen T, Jensen A, Moore S, Grimsbø GH, Børøsund E, Ellison MC. Effects of an internet support system to assist cancer patients in reducing symptom distress: a randomized controlled trial. *Cancer Nursing* 2013;**36**(1):6–17.
- Ryser 2004** {published data only}  
Ryser FG. Breastfeeding attitudes, intention, and initiation in low-income women: the effect of the best start program. *Journal of Human Lactation* 2004;**20**(3):300–5.
- Sassen 2014** {published data only}  
Sassen B, Kok G, Schepers J, Vanhees L. Supporting health care professionals to improve the processes of shared decision making and self-management in a web-based intervention: randomized controlled trial. *Journal of Medical Internet Research* 2014;**16**(10):e211.
- Saver 2007** {published data only}  
Saver BG, Gustafson D, Taylor TR, Hawkins RP, Woods NF, Dinauer S, et al. A tale of two studies: the importance of setting, subjects and context in two randomized, controlled trials of a web-based decision support for perimenopausal and postmenopausal health decisions. *Patient Education and Counseling* 2007;**66**(2):211–22.
- Sawka 2011** {published data only}  
Sawka AM, Straus S, Gafni A, Brierley JD, Tsang RW, Rotstein L, et al. How can we meet the information needs of patients with early stage papillary thyroid cancer considering radioactive iodine remnant ablation?. *Clinical Endocrinology* 2011;**74**:419–23.
- Scaffidi 2014** {published data only}  
Scaffidi RM, Posmontier B, Bloch JR, Wittmann-Price R. The relationship between personal knowledge and decision self-efficacy in choosing trial of labor after cesarean. *Journal of Midwifery & Women's Health* 2014;**59**(3):246–53.
- Schapira 2000** {published data only}  
Schapira MM, VanRuiswyk J. The effect of an illustrated pamphlet decision-aid on the use of prostate cancer screening tests. *Journal of Family Practice* 2000;**49**(5):418–24.
- Schapira 2007** {published data only}  
Schapira MM, Gilligan MA, McAuliffe T, Garmon G, Carnes M, Nattinger AB. Decision-making at menopause: a randomized controlled trial of a computer-based hormone therapy decision-aid. *Patient Education and Counseling* 2007;**67**(1-2):100–7.
- Schwartz 2009b** {published data only}  
Schwartz LM, Woloshin S, Welch HG. Using a drug facts box to communicate drug benefits and harms: two randomized trials. *Annals of Internal Medicine* 2009;**150**(8):516–27.
- Sears 2007** {published data only}  
Sears SR, Woodward JT, Twillman RK. What do I have to lose? effects of a psycho-educational intervention on cancer patient preference for resuscitation. *Journal of Behavioral Medicine* 2007;**30**(6):533–44.
- Sequist 2011** {published data only}  
Sequist T, Zaslavsky A, Colditz G, Ayanian J. Electronic patient message to promote colorectal cancer screening. *Archives of Internal Medicine* 2011;**171**(7):636–41.
- Shah 2012** {published data only}  
Shah S, Singh K, Ali MK, Mohan V, Kadir MM, Unnikrishnan AG, et al. CARRS Trial Writing Group. Improving diabetes care: multi-component cardiovascular disease risk reduction strategies for people with diabetes in South Asia - the CARRS multi-center translation trial. *Diabetes Research and Clinical Practice* 2012;**98**(2):285–94.
- Sheppard 2012** {published data only}  
Sheppard VB, Wallington SF, Williams KP, Lucas W. A decision-support intervention for black women eligible for adjuvant systematic therapy: Sisters informing sisters about breast cancer treatment - An intervention to reduce treatment disparities. In: Elk R, Landrine H editor(s). *Cancer Disparities: Causes and Evidence-Based Solutions*. American Cancer Society, 2012.
- Sheridan 2004** {published data only}  
Sheridan SL, Felix K, Pignone MP, Lewis CL. Information needs of men regarding prostate cancer screening and the effect of a brief decision aid. *Patient Education and Counseling* 2004;**54**(3):345–51.
- Sheridan 2010** {published data only}  
Sheridan SL, Griffith JM, Behrend L, Gizlice Z, Jianwen C, Pignone MP. Effect of adding a values clarification exercise to a decision aid on heart disease prevention: a randomized trial. *Medical Decision Making* 2010;**30**(4):E28–39.
- Sheridan 2012** {published data only}  
Sheridan SL, Golin C, Bunton A, Lykes JB, Schwartz B, McCormack L, Driscoll D, Bangdiwala SI, Harris RP. Shared decision making for prostate cancer screening: the results of a combined analysis of two practice-based

- randomized controlled trials. *BMC Medical Informatics & Decision Making* 2012;**12**:130.
- Sherman 2014** *{published data only}*  
Sherman KA, Harcourt DM, Lam TC, Shaw LK, Boyages J. BRECONDA: Development and acceptability of an interactive decisional support tool for women considering breast reconstruction. *Psycho-Oncology* 2014;**23**:835–8.
- Shirai 2012** *{published data only}*  
Shirai Y, Fujimori M, Ogawa A, Yamada Y, Nishiwaki Y, Ohtsu A, Uchitomi Y. Patients' perception of the usefulness of a question prompt sheet for advanced cancer patients when deciding the initial treatment: a randomized, controlled trial. *Psycho-Oncology* 2012;**21**(7):706–713.
- Silver 2012** *{published data only}*  
Silver B, Zaman IF, Ashraf K, Majed Y, Norwood EM, Schuh LA, et al. A randomized trial of decision-making in asymptomatic carotid stenosis. *Neurology* 2012;**78**(5):315–21.
- Siminoff 2006** *{published data only}*  
Siminoff LA, Gordon NH, Silverman P, Budd T, Ravdin PM. A decision aid to assist in adjuvant therapy choices for breast cancer. *Psycho-Oncology* 2006;**15**(11):1001–13.  
Vickers AJ, Elkin EB, Peele PB, Dickler M, Siminoff LA. Long-term health outcomes of a decision aid: data from a randomized trial of adjuvant in women with localized breast cancer. *Medical Decision Making* 2009;**29**(4):461–7.
- Simon 2012a** *{published data only}*  
Simon D, Kriston L, von Wolff A, Buchholz A, Vietor C, Hecke T, et al. Effectiveness of a web-based, individually tailored decision aid for depression or acute low back pain: a randomized controlled trial. *Patient Education and Counseling* 2012;**87**(3):360–8.
- Simon 2012b** *{published data only}*  
Simon W, Lambert MJ, Harris MW, Busath G, Vazquez A. Providing patient progress information and clinical support tools to therapists: Effects on patients at risk of treatment failure. *Psychotherapy Research* 2012;**22**(6):638–47.
- Smith 2011a** *{published data only}*  
Smith T, Dow L, Virago E, Khatcheressian J, Matsuyama R, Lyckholm L. A pilot trial of decision aids to give truthful prognostic and treatment information to chemotherapy patients with advanced cancer. *Journal of Supportive Oncology* 2011;**9**(2):79–86.
- Smith 2011b** *{published data only}*  
Smith SW, Nazione S, LaPlante C, Clark-Hitt R, Park HS, Sung R, Leichtman A. Living kidney donor decision making and communication. *Journal of Health Communication: International Perspectives* 2011;**16**(8):870–88.
- Solberg 2010** *{published data only}*  
Solberg LI, Asche SE, Sepucha K, Thygeson NM, Madden JE, Morrissey L, et al. Informed choice assistance for women making uterine fibroid treatment decisions: a practical clinical trial. *Medical Decision Making* 2010;**30**(4):444–52.
- Sorenson 2004** *{published data only}*  
Sorenson JR, Lakon C, Spinney T, Jennings-Grant T. Assessment of a decision aid to assist genetic testing research participants in the informed consent process. *Genetic Testing* 2004;**8**(3):336–46.
- Sparano 2006** *{published data only}*  
Sparano JA. TAILORx: trial assigning individualized options for treatment (Rx). *Clinical Breast Cancer* 2006;**7**(4):347–50.
- Stalmeier 2009** *{published data only}*  
Stalmeier PF, Roosmalen MS. Concise evaluation of decision aids. *Patient Education and Counseling* 2009;**74**(1):104–9.
- Starosta 2015** *{published data only}*  
Starosta AJ, Luta G, Tomko CA, Schwartz MD, Taylor KL. Baseline attitudes about prostate cancer screening moderate the impact of decision aids on screening rates. *Annals of Behavioral Medicine* 2015;**49**:762–768.
- Stein 2013** *{published data only}*  
Stein RA, Sharpe L, Bell ML, Boyle FM, Dunn SM, Clarke SJ. Randomized controlled trial of a structured intervention to facilitate end-of-life decision making in patients with advanced cancer. *Journal of Clinical Oncology* 2013;**31**(27):3403–10.
- Steiner 2003** *{published data only}*  
Steiner MJ, Dalebout S, Condon S, Dominik R, Trussell J. Understanding risk: a randomized controlled trial of communicating contraceptive effectiveness. *Obstetrics & Gynecology* 2003;**102**(4):709–17.
- Stephens 2008** *{published data only}*  
Stephens RL, Xu Y, Volk RJ, Scholl LE, Kamin SL, Holden EW. Influence of a patient decision aid on decisional conflict related to PSA testing: a structural equation model. *Health Psychology* 2008;**27**(6):711–21.
- Stiggelbout 2008** *{published data only}*  
Stiggelbout AM, Molewijk AC, Otten W, van Bockel JH, Bruijninx CM, van der Salm I, et al. The impact of individualized evidence-based decision support on aneurysm patients' decision making, ideals of autonomy, and quality of life. *Medical Decision Making* 2008;**28**(5):751–62.
- Stirling 2012** *{published data only}*  
Stirling C, Leggett S, Lloyd B, Scott J, Blizzard L, Quinn S, Robinson A. Decision aids for respite service choices by carers of people with dementia: development and pilot RCT. *BMC Medical Informatics and Decision Making* 2012;**12**:21.
- Street 1995** *{published data only}*  
Street RLJ, Voigt B, Geyer CJ, Manning T, Swanson GP. Increasing patient involvement in choosing treatment for early breast cancer. *Cancer* 1995;**76**(11):2275–85.
- Street 1998** *{published data only}*  
Street RL Jr, Van Order A, Bramson R, Manning T. Preconsultation education promoting breast cancer screening: does the choice of media make a difference?. *Journal of Cancer Education* 1998;**13**(3):152–61.

- Sundaresan 2011** {published data only}  
Sundaresan P, Turner S, Kneebone A, Pearse M, Butow P. Evaluating the utility of a patient decision aid for potential participants of a prostate cancer trial (RAVES-TROG 08.03). *Radiotherapy and Oncology* 2011;**101**(3):521–4.
- Tabak 1995** {published data only}  
Tabak N. Decision making in consenting to experimental cancer therapy. *Cancer Nursing* 1995;**18**(2):89–96.
- Taylor 2013** {published data only}  
Taylor KL, Williams RM, Davis K, Luta G, Penek S, Barry S, et al. Decision making in prostate cancer screening using decision aids vs usual care: a randomized clinical trial. *JAMA Internal Medicine* 2013;**173**(18):1704–12.
- Ten 2008** {published data only}  
Ten Wolde GB, Dijkstra A, van Empelen P, van den Hout W, Neven AK, Zitman F. Long-term effectiveness of computer-generated tailored patient education on benzodiazepines: a randomized controlled trial. *Addiction* 2008;**103**(4):662–70.
- Thomas 2013** {published data only}  
Thomas KL, Zimmer LO, Dai D, Al-Khatib SM, Allen LaPointe NM, Peterson ED. Educational videos to reduce racial disparities in ICD therapy via innovative designs (VIVID): a randomized clinical trial. *American Heart Journal* 2013;**166**(1):157–63.
- Thomson 2006** {published data only}  
Thomson P, Dowding D, Swanson V, Bland R, Mair C, Morrison A, et al. A computerised guidance tree (decision aid) for hypertension, based on decision analysis: development and preliminary evaluation. *European Journal of Cardiovascular Nursing* 2006;**5**(2):146–9.
- Thornton 1995** {published data only}  
Thornton JG, Hewison J, Lilford RJ, Vail A. A randomised trial of three methods of giving information about prenatal testing. *BMJ* 1995;**311**(7013):1127–30.
- Tiller 2006** {published data only}  
Tiller K, Meiser B, Gaff C, Kirk J, Dudding T, Phillips KA, et al. A randomized controlled trial of a decision aid for women at increased risk of ovarian cancer. *Medical Decision Making* 2006;**26**(4):360–72.
- Tinsel 2013** {published data only}  
Tinsel I, Buchholz A, Vach W, Siegel A, Dürk T, Buchholz A, et al. Shared decision-making in antihypertensive therapy: a cluster randomised controlled trial. *BMC Family Practice* 2013;**14**:135.
- Tomko 2015** {published data only}  
Tomko C, Davis K, Ludin S, Kelly S, Stern A, Luta G, et al. Decisional outcomes following use of an interactive web-based decision aid for prostate cancer screening. *Translational Behavioral Medicine: Practice, Policy, Research* 2015;**5**(2):189–97.
- Ukoli 2013** {published data only}  
Ukoli FA, Patel K, Hargreaves M, Beard K, Moton PJ, Bragg R, et al. A tailored prostate cancer education intervention for low-income African Americans: impact on knowledge and screening. *Journal of Health Care for the Poor and Underserved* 2013;**24**(1):311–331.
- Valdez 2001** {published data only}  
Valdez A, Banerjee K, Fernandez M, Ackerson L. Impact of a multimedia breast cancer education intervention on use of mammography by low-income Latinas. *Journal of Cancer Education* 2001;**16**(4):221–4.
- Van der Krieke 2013** {published data only}  
Van der Krieke L, Emerencia AC, Boonstra N, Wunderink L, de Jonge P, Sytema S. A web-based tool to support shared decision making for people with a psychotic disorder: randomized controlled trial and process evaluation. *Journal of Medical Internet Research* 2013;**15**(10):e216.
- Van Roosmalen 2004** {published and unpublished data}  
Van Roosmalen MS, Stalmeier PF, Verhoef LC, Hoekstra-Weebers JE, Oosterwijk JC, Hoogerbrugge N, et al. Randomised trial of a decision aid and its timing for women being tested for a BRCA1/2 mutation. *British Journal of Cancer* 2004;**90**(2):333–42.  
\* Van Roosmalen MS, Stalmeier PF, Verhoef LC, Hoekstra-Weebers JE, Oosterwijk JC, Hoogerbrugge N, et al. Randomized trial of a shared decision-making intervention consisting of trade-offs and individualized treatment information for BRCA1/2 mutation carriers. *Journal of Clinical Oncology* 2004;**22**(16):3293–301.
- Van Steenkiste 2008** {published data only}  
Van Steenkiste B, van der Weijden TM, Stoffers JHEH, Grol RPTM. Patients' responsiveness to a decision support tool for primary prevention of cardiovascular diseases in primary care. *Patient Education and Counseling* 2008;**72**(1):63–70.
- Van Til 2009** {published data only}  
Van Til JA, Stiggelbout AM, IJzerman MJ. The effect of information on preferences stated in a choice-based conjoint analysis. *Patient Education and Counseling* 2009;**74**(2):264–71.
- Van Tol-Geerdink 2013** {published data only}  
Van Tol-Geerdink JJ, Leer JW, Weijerman PC, van Oort IM, Vergunst H, van Lin EN, et al. Choice between prostatectomy and radiotherapy when men are eligible for both: a randomized controlled trial of usual care vs decision aid. *BJU International* 2013;**111**(4):564–73.
- Veroff 2012** {published data only}  
Veroff D, Sullivan L, Shoptaw EJ, Venator B, Ochoa-Arvelo T, Baxter J, et al. Improving self-care for heart failure for seniors: Impact of video and written education and decision aids. *Population Health Management* 2012;**15**(1):37–45.
- Volandes 2009** {published data only}  
Volandes AE, Paasche-Orlow MK, Barry MJ, Gillick MR, Minaker KL, Chang Y, et al. Video decision support tool for advance care planning in dementia: randomised controlled trial. *BMJ* 2009;**338**:b2159.
- Volandes 2011** {published data only}  
Volandes A, Ferguson L, Davis A, Hull N, Green M, Chang Y, et al. Assessing end-of-life preferences for advanced

- dementia in rural patients using an educational video: a randomised controlled trial. *Journal of Palliative Medicine* 2011;**14**(2):169–77.
- Volandes 2013** *{published data only}*  
Volandes AE, Paasche-Orlow MK, Mitchell SL, El-Jawahri A, Davis AD, Barry MJ, et al. Randomized controlled trial of a video decision support tool for cardiopulmonary resuscitation decision making in advanced care. *Journal of Clinical Oncology* 2013;**31**(3):380–6.
- Volk 2008** *{published data only}*  
Volk RJ, Jibaja-Weiss ML, Hawley ST, Kneuper S, Spann SJ, Miles BJ, et al. Entertainment education for prostate cancer screening: a randomized trial among primary care patients with low health literacy. *Patient Education and Counseling* 2008;**73**(3):482–9.
- Von Wagner 2011** *{published data only}*  
Von Wagner C. A decision aid to support informed choice about bowel cancer screening in people with low educational level improves knowledge but reduces screening uptake. *Evidence-Based Nursing* 2011;**14**(2):36–7.
- Wagner 1995** *{published data only}*  
Wagner EH, Barrett P, Barry MJ, Barlow W, Fowler FJ Jr. The effect of a shared decision making program on rates of surgery for benign prostatic hyperplasia. Pilot results. *Medical Care* 1995;**33**(8):765–70.
- Wakefield 2008a** *{published data only}*  
Wakefield CE, Meiser B, Homewood J, Ward R, O'Donnell S, Kirk J, et al. Randomized trial of a decision aid for individuals considering genetic testing for hereditary nonpolyposis colorectal cancer risk. *Cancer* 2008;**113**(5):956–65.
- Wakefield 2008b** *{published data only}*  
Wakefield CE, Meiser B, Homewood J, Peate M, Taylor A, Lobb E, et al. A randomized controlled trial of a decision aid for women considering genetic testing for breast and ovarian cancer risk. *Breast Cancer Research and Treatment* 2008;**107**(2):289–301.
- Wakefield 2008c** *{published data only}*  
Wakefield CE, Meiser B, Homewood J, Taylor A, Gleeson M, Williams R. A randomized trial of a breast/ovarian cancer genetic testing decision aid used as a communication aid during genetic counseling. *Psycho-Oncology* 2008;**17**(8):844–54.
- Wallston 1991** *{published data only}*  
Wallston KA, Smith RA, King JE, Smith MS, Rye P, Burish TG. Desire for control and choice of antiemetic treatment for cancer chemotherapy. *Western Journal of Nursing Research* 1991;**13**(1):12–23.
- Wang 2004** *{published data only}*  
Wang C, Gonzalez R, Milliron KJ, Strecher VJ, Merajver SD. Genetic counseling for BRCA1/2: a randomized controlled trial of two strategies to facilitate the education and counseling process. *American Journal of Medical Genetics. Part A* 2005;**134**(1):66–73.
- Warner 2015** *{published data only}*  
Warner DO, LeBlanc A, Kadimpati S, Vickers KS, Shi Y, Montori V. Decision aid for cigarette smokers scheduled for elective surgery. *Anesthesiology* 2015;**123**(1):18–28.
- Watts 2014** *{published data only}*  
Watts KJ, Meiser B, Wakefield CE, Barratt AL, Howard K, Cheah BC, et al. Online prostate cancer screening decision aid for at-risk men: a randomized trial. *Health Psychology* 2014;**33**(9):986–97.
- Welschen 2012** *{published data only}*  
Welschen LM, Bot SD, Kostense PJ, Dekker JM, Timmermans DR, van der Weijden T, et al. Effects of cardiovascular disease risk communication for patients with type 2 diabetes on risk perception in a randomized controlled trial: the @RISK study. *Diabetes Care* 2012;**35**:2485–92.
- Wennberg 2010** *{published data only}*  
Wennberg DE, Marr A, Lang L, O'Malley S, Bennett G. A randomized trial of a telephone care-management strategy. *New England Journal of Medicine* 2010;**363**(13):1245–55.
- Westermann 2013** *{published data only}*  
Westermann GM, Verheij F, Winkens B, Verhulst FC, Van Oort FV. Structured shared decision-making using dialogue and visualization: a randomized controlled trial. *Patient Education and Counseling* 2013;**90**(1):74–81.
- Weymann 2015** *{published data only}*  
Weymann N, Dirmaier J, von Wolff A, Kriston L, Härter M. Effectiveness of a web-based tailored interactive health communication application for patients with type 2 diabetes or chronic low back pain: randomized controlled trial. *Journal of Medical Internet Research* 2015;**17**(3):e53 1–21.
- Wilhelm 2009** *{published data only}*  
Wilhelm D, Gillen S, Wirnhier H, Kranzfelder M, Schneider A, Schmidt A, et al. Extended preoperative patient education using a multimedia DVD: impact on patients receiving a laparoscopic cholecystectomy: a randomised controlled trial. *Langenbeck's Archives of Surgery* 2009;**394**(2):227–33.
- Wilkes 2013** *{published data only}*  
Wilkes MS, Day FC, Srinivasan M, Griffin E, Tancredi DJ, Rainwater JA, et al. Pairing physician education with patient activation to improve shared decisions in prostate cancer screening: a cluster randomized controlled trial. *Annals of Family Medicine* 2013;**11**(4):324–34.
- Wilkie 2013** *{published data only}*  
Wilkie DJ, Gallo AM, Yao Y, Molokie RE, Stahl C, Hershberger PE, et al. Reproductive health choices for young adults with sickle cell disease or trait: randomized controlled trial immediate posttest effects. *Nursing Research* 2013;**62**(5):352–61.
- Wilkins 2006** *{published data only}*  
Wilkins EG, Lowery JC, Copeland LA, Goldfarb SL, Wren PA, Janz NK. Impact of an educational video on patient decision making in early breast cancer treatment. *Medical Decision Making* 2006;**26**(6):589–98.

**Willemsen 2006 {published data only}**

Willemsen MC, Wiebing M, van Erst A, Zeeman G. Helping smokers to decide on the use of efficacious smoking cessation methods: a randomized controlled trial of a decision aid. *Addiction* 2006;**101**(3):441–9.

**Williamson 2014 {published data only}**

Williamson LEA, Lawson KL, Downe PJ, Pierson RA. Informed reproductive decision-making: The impact of providing fertility information on fertility knowledge and intentions to delay childbearing. *Journals of Obstetrics and Gynaecology Canada* 2014;**36**(5):400–5.

**Williams-Piehota 2008 {published data only}**

Williams-Piehota PA, McCormack LA, Treiman K, Bann CM. Health information styles among participants in a prostate cancer screening informed decision-making intervention. *Health Education Research* 2008;**23**(3): 440–53.

**Wolmann 2011 {published data only}**

Wolmann EM, Wilkiss SM, Teachout A, McHugo GJ, Drake RE. Trial of an electronic decision support system to facilitate shared decision making in community mental health. *Psychiatric Services* 2011;**62**(1):54–60.

**Wroe 2005 {published data only}**

Wroe AL, Turner N, Owens RG. Evaluation of a decision-making aid for parents regarding childhood immunizations. *Health Psychology* 2005;**24**(6):539–47.

**Yee 2014 {published data only}**

Yee LM, Wolf M, Mullen R, Bergeron AR, Cooper Bailey S, Levine R, Grobman WA. A randomized trial of a prenatal genetic testing interactive computerized information aid. *Prenatal Diagnosis* 2014;**34**(6):552–7.

**Yun 2011 {published data only}**

Yun YH, Lee MK, Park S, Lee JL, Park J, Choi YS, et al. Use of a decision aid to help caregivers discuss terminal disease status with a family member with cancer: a randomized controlled trial. *Journal of Clinical Oncology* 2011;**29**(36): 4811–9.

**Zajac 2012 {published data only}**

Zajac LE. *Making Difficult Health Decisions: A Motivated Decision Processing Model [PhD thesis]*. Pittsburgh: University of Pittsburgh, 2012.

**Zapka 2004 {published data only}**

Zapka JG, Lemon SC, Puleo E, Estabrook B, Luckmann R, Erban S. Patient education for colon cancer screening: a randomized trial of a video mailed before a physical examination. *Annals of Internal Medicine* 2004;**141**(9): 683–92.

**Zikmund-Fisher 2008 {published data only}**

Zikmund-Fisher BJ, Ubel PA, Smith DM, Derry HA, McClure JB, Stark A, et al. Communicating side effect risks in a tamoxifen prophylaxis decision aid: the debiasing influence of pictographs. *Patient Education and Counseling* 2008;**73**(2):209–14.

**Zoffman 2012 {published data only}**

Zoffman V, Kirkevold M. Realizing empowerment in difficult diabetes care: a guided self-determination

intervention. *Qualitative Health Research* 2012;**22**(1): 103–18.

## References to ongoing studies

**ACTRN12615000523505 {published data only}**

ACTRN12615000523505. The Motherhood Choices Decision Aid for Women with Rheumatoid Arthritis Increases Knowledge and Reduces Decisional Conflict: A Randomized Controlled Study. <http://apps.who.int/trialsearch/Trial2.aspx?TrialID=ACTRN12615000523505> (first received May 25, 2015).

**ACTRN12615000843550 {published data only}**

ACTRN12615000843550. Evaluation of decision aids for parents about the benefits and harms of antibiotic use for coughs and colds in children [Pilot randomised controlled trial of decision aids for parents about the benefit and harm of antibiotics for common acute respiratory infections in children to aid informed decision making]. <http://apps.who.int/trialsearch/Trial2.aspx?TrialID=ACTRN12615000843550> (first received August 13, 2015).

**Al-Itejawi 2015 {published data only}**

Al-Itejawi HH, van Uden-Kraan CF, Vis AN, Nieuwenhuijzen JA, Hofstee MJ, van Moorselaar RJ, Verdonck-de Leeuw IM. Development of a patient decision aid for the treatment of localised prostate cancer: a participatory design approach. *J Clin Nurs* 2016;**25**(7-8): 1131–1144.

**Anderson 2014 {published data only}**

Anderson RT, Montori VM, Shah ND, Ting HH, Pencille LJ, Demers M, et al. Effectiveness of the Chest Pain Choice decision aid in emergency department patients with low-risk chest pain: study protocol for a multicenter randomized trial. *Trials* 2014;**15**(166):1–11.

**Aslani 2014 {published data only}**

Aslani A, Tara F, Ghalichi L, Eslami S. The impact of computerized decision aid on mode of delivery - a study protocol. *Studies in Health Technology and Informatics* 2014; **200**:170–2.

**Buhse 2013 {published data only}**

Buhse S, Heller T, Kasper J, Mühlhauser I, Müller UA, Lehmann T, Lenz M. An evidence-based shared decision making programme on the prevention of myocardial infarction in type 2 diabetes: protocol of a randomised-controlled trial. *BMC Family Practice* 2013;**14**(155):1–8.

**Carroll 2012 {published data only}**

Carroll SL, McGillion M, Stacey D, Healey JS, Browne G, Arthur HM, Thabane L. Development and feasibility testing of decision support for patients who are candidates for a prophylactic implantable defibrillator: a study protocol for a pilot randomized controlled trial. *Trials* 2013;**14**(346): 1–9.

**Chambers 2008 {published data only}**

Chambers SK, Ferguson M, Gardiner RA, Nicol D, Gordon L, Occhipinti S, et al. ProsCan for men: randomised controlled trial of a decision support intervention for men with localised prostate cancer. *BMC Cancer* 2008;**8**:207.



- Coylewright 2012** *{published data only}*  
Coylewright M, Shepel K, LeBlanc A, Pencille L, Hess E, Shah N, Montori VM, Ting HH. Shared decision making in patients with stable coronary artery disease: PCI choice. *PLOS One* 2012;7(11):1–8.
- Cuyppers 2015** *{published data only}*  
Cuyppers M, Lamers RE, Kil PJ, van de Poll-Franse LV, de Vries M. Impact of a web-based treatment decision aid for early-stage prostate cancer on shared decision-making and health outcomes: study protocol for a randomized controlled trial. *Trials* 2015;16:231.
- Den Ouden 2015** *{published data only}*  
Den Ouden H, Vos RC, Reidsma C, Rutten GEHM. Shared decision making in type 2 diabetes with a support decision tool that takes into account clinical factors, the intensity of treatment and patient preferences: design of a cluster randomised (OPTIMAL) trial. *BMC Family Practice* 2015;16:27.
- Dirmaier 2013** *{published data only}*  
Dirmaier J, Härter M, Weymann N. A tailored, dialogue-based health communication application for patients with chronic low back pain: study protocol of a randomised controlled trial. *BMC Medical Informatics & Decision Making* 2013;13(66):1–9.
- Geiger 2011** *{published data only}*  
Geiger F, Liethmann K, Hoffmann F, Paschedag J, Kasper J. Investigating a training supporting Shared Decision Making (IT'S SDM 2011): study protocol for a randomized controlled trial. *Trials* 2011;12:232.
- Hersch 2014** *{published data only}*  
Hersch J, Barratt A, Jansen J, Houssami N, Irwig L, Jacklyn G, et al. The effect of information about overdetected breast cancer on women's decision-making about mammography screening: study protocol for a randomised controlled trial. *BMJ Open* 2014;4(5):1–10.
- Hess 2014** *{published data only}*  
Hess EP, Wyatt KD, Kharbanda AB, Louie JP, Dayan PS, Tzimenatos L, et al. Effectiveness of the head CT choice decision aid in parents of children with minor head trauma: study protocol for a multicenter randomized trial. *Trials* 2014;14(253):1–11.
- Jimbo 2012** *{published data only}*  
Jimbo M, Kelly-Blake K, Sen A, Hawley ST, Ruffin MT 4th. Decision Aid to Technologically Enhance Shared decision making (DATES): study protocol for a randomized controlled trial. *Trials* 2013;14(381):1–16.
- Layton 2012** *{published data only}*  
Effects of a web-based decision aid on African American men's prostate screening knowledge and behavior. Ongoing study -.
- LeBlanc 2013** *{published data only}*  
LeBlanc A, Bodde AE, Branda ME, Yost KJ, Herrin J, Williams MD, et al. Translating comparative effectiveness of depression medications into practice by comparing the depression medication choice decision aid to usual care: study protocol for a randomized controlled trial. *Trials* 2013;14(127):1–8.
- Mann 2012** *{published data only}*  
Mann DM, Lin JJ. Increasing efficacy of primary care-based counseling for diabetes prevention: rationale and design of the ADAPT (Avoiding Diabetes Thru Action Plan Targeting) trial. *Implementation Science* 2012;7:6.
- NCT00813033** *{published data only}*  
NCT00813033. Use of a Patient Decision Aid for Gastrologic Endoscopy in a Pediatric Setting [Creation and Pilot Evaluation of a Patient Decision Aid as an Adjunct to the Consenting Process for Gastrointestinal Endoscopy in a Pediatric Setting]. <https://clinicaltrials.gov/show/NCT00813033> (first received December 19, 2008).
- NCT01077037** *{published data only}*  
NCT01077037. Impact of a Decision Aid on Patient Decision Making in Emergency Department Chest Pain Patients [Impact of a Decision Aid on Patient Participation in Decision Making and Resource Use in Low Risk Chest Pain Patients: A Randomized Trial]. [clinicaltrials.gov/show/NCT01077037](https://clinicaltrials.gov/show/NCT01077037) (first received February 24, 2010).
- NCT01152294** *{published data only}*  
NCT01152294. Measuring Quality of Decisions About Treatment of Menopausal Symptoms [Measuring Quality of Decisions About Treatment of Menopausal Symptoms]. [clinicaltrials.gov/show/NCT01152294](https://clinicaltrials.gov/show/NCT01152294) (first received June 22, 2010).
- NCT01152307** *{published data only}*  
NCT01152307. Measuring Quality of Decisions About Treatment of Depression [Measuring Quality of Decisions About Treatment of Depression]. [clinicaltrials.gov/show/NCT01152307](https://clinicaltrials.gov/show/NCT01152307) (first received June 22, 2010).
- NCT01447186** *{published data only}*  
NCT01447186. Adaptation of the American Cancer Society (ACS) Early Detection of Prostate Cancer Patient Decision Aid for Spanish Speaking Men [Adaptation of the American Cancer Society (ACS) Early Detection of Prostate Cancer Patient Decision Aid for Spanish Speaking Men]. [clinicaltrials.gov/show/NCT01447186](https://clinicaltrials.gov/show/NCT01447186) (first received October 3, 2011).
- NCT01618097** *{published data only}*  
NCT01618097. Evaluation of DVD and Internet Decision Aids for Hip and Knee Osteoarthritis: Focus on Health Literacy. [clinicaltrials.gov/show/NCT01618097](https://clinicaltrials.gov/show/NCT01618097) (first received May 29, 2012).
- NCT01713894** *{published data only}*  
NCT01713894. Decision Aid - Extreme Prematurity [Utility of a Clinically Relevant Decision Aid, for Parents Facing Extremely Premature Delivery]. [clinicaltrials.gov/show/NCT01713894](https://clinicaltrials.gov/show/NCT01713894) (first received October 22, 2012).
- NCT01771536** *{published data only}*  
NCT01771536. The PCI Choice Trial: a Pilot Randomized Trial of a Decision Aid for Patients With Stable Coronary Artery Disease. [clinicaltrials.gov/show/NCT01771536](https://clinicaltrials.gov/show/NCT01771536) (first received December 15, 2012).

**NCT01851785 {published data only}**

NCT01851785. African American Preference for Knee Replacement: A Patient-Centered Intervention (ACTION) [Behavioral & Social Science Research on Understanding and Reducing Health Disparities]. [clinicaltrials.gov/show/NCT01851785](http://clinicaltrials.gov/show/NCT01851785) (first received May 8, 2013).

**NCT01941186 {published data only}**

NCT01941186. A Family Centered Intervention to Promote Optimal Child Development [A Family Centered Intervention to Promote Optimal Child Development at the Interface of the Health System and Community]. [clinicaltrials.gov/show/NCT01941186](http://clinicaltrials.gov/show/NCT01941186) (first received September 9, 2013).

**NCT01976325 {published data only}**

NCT01976325. Evaluating the Ottawa Malaria Decision Aid (OMDA) [Incorporation of the 'Ottawa Malaria Decision Aid' Into the Pre-travel Consultation Process: Assessment of Travelers' Knowledge, Decisional Conflict, Preparation for Decision-making and Medication Adherence Compared to Standard Care]. [clinicaltrials.gov/show/NCT01976325](http://clinicaltrials.gov/show/NCT01976325) (first received October 29, 2013).

**NCT02026102 {published data only}**

NCT02026102. A Pilot Trial of Patient Decision Aids for Implantable Cardioverter-Defibrillators (ICDs) [A Pilot Trial of Patient Decision Aids for Implantable Cardioverter-Defibrillators (ICDs)]. [clinicaltrials.gov/show/NCT02026102](http://clinicaltrials.gov/show/NCT02026102) (first received December 12, 2013).

**NCT02084290 {published data only}**

NCT02084290. Evaluating a Shared Decision Making Program for Crohn's Disease [Evaluating a Prediction Tool and Decision Aid for Patients With Crohn's Disease]. [clinicaltrials.gov/show/NCT02084290](http://clinicaltrials.gov/show/NCT02084290) (first received January 29, 2014).

**NCT02110979 {published data only}**

NCT02110979. Validation of a Patient Decision Aid for Type 2 Diabetes [Validation of a Patient Decision Aid for Type 2 Diabetes]. [clinicaltrials.gov/show/NCT02110979](http://clinicaltrials.gov/show/NCT02110979) (first received April 4, 2014).

**NCT02145481 {published data only}**

NCT02145481. Decisional Quality for Patients With Coronary Artery Disease (DeQCAD). [clinicaltrials.gov/show/NCT02145481](http://clinicaltrials.gov/show/NCT02145481) (first received May 15, 2014).

**NCT02198690 {published data only}**

NCT02198690. Trial of a Mammography Decision Aid for Women Aged 75 and Older [Randomized Trial of a Mammography Decision Aid for Women Aged 75 and Older]. [clinicaltrials.gov/show/NCT02198690](http://clinicaltrials.gov/show/NCT02198690) (first received July 16, 2014).

**NCT02235571 {published data only}**

NCT02235571. iChoose Decision Kidney Aid for End-Stage Renal Disease Patients [iChoose Kidney Decision Aid for Treatment Options Among End-Stage Renal Disease (ESRD) Patients]. [clinicaltrials.gov/show/NCT02235571](http://clinicaltrials.gov/show/NCT02235571) (first received September 6, 2014).

**NCT02248974 {published data only}**

NCT02248974. Development & Testing of a Decision Aid for LVAD Placement (VADDA) [Development and User Testing of a Decision Aid for Left Ventricular Assist Device (LVAD) Placement]. [clinicaltrials.gov/show/NCT02248974](http://clinicaltrials.gov/show/NCT02248974) (first received September 17, 2014).

**NCT02259699 {published data only}**

NCT02259699. Ovarian Cancer Patient-Centered Decision Aid (PCOA) [Ovarian Cancer Patient-Centered Decision Aid]. [clinicaltrials.gov/show/NCT02259699](http://clinicaltrials.gov/show/NCT02259699) (first received May 15, 2014).

**NCT02308592 {published data only}**

NCT02308592. Patient Decision Aid for Antidepressant Use in Pregnancy [Patient Decision Aid (PDA) for Antidepressant Use In Pregnancy]. [clinicaltrials.gov/show/NCT02308592](http://clinicaltrials.gov/show/NCT02308592) (first received December 2, 2014).

**NCT02319525 {published data only}**

NCT02319525. Individualized Patient Decision Making for Treatment Choices Among Minorities With Lupus [Individualized Patient Decision Making for Treatment Choices Among Minorities With Lupus]. [clinicaltrials.gov/show/NCT02319525](http://clinicaltrials.gov/show/NCT02319525) (first received November 5, 2014).

**NCT02326597 {published data only}**

NCT02326597. Decision Aid for Therapeutic Options In Sickle Cell Disease [Comparative Effectiveness of a Decision Aid for Therapeutic Options in Sickle Cell Disease]. [clinicaltrials.gov/show/NCT02326597](http://clinicaltrials.gov/show/NCT02326597) (first received December 18, 2014).

**NCT02344576 {published data only}**

NCT02344576. Trial of a Decision Support Intervention for Patients and Caregivers Offered Destination Therapy Heart Assist Device (DECIDE-LVAD) [A Multicenter Trial of a Shared Decision Support Intervention for Patients and Their Caregivers Offered Destination Therapy for End-Stage Heart Failure]. [clinicaltrials.gov/show/NCT02344576](http://clinicaltrials.gov/show/NCT02344576) (first received January 16, 2015).

**NCT02488317 {published data only}**

NCT02488317. Empowering Patients On Choices for Renal Replacement Therapy (Aim 3) (EPOCH-RRT) (EPOCH-RRT) [Empowering Patients On Choices for Renal Replacement Therapy (Aim 3)]. [clinicaltrials.gov/show/NCT02488317](http://clinicaltrials.gov/show/NCT02488317) (first received June 15, 2015).

**NCT02488603 {published data only}**

NCT02488603. Decision Aids for Tamoxifen Treatment in Breast Cancer Patients [Utilization of Decision Aids for Tamoxifen Treatment in Breast Cancer Patients: A Randomized Controlled Trial.]. [clinicaltrials.gov/show/NCT02488603](http://clinicaltrials.gov/show/NCT02488603) (first received June 25, 2015).

**NCT02492009 {published data only}**

NCT02492009. Patient Decision Aid for Antidepressant Use in Pregnancy [Patient Decision Aid (PDA) for Antidepressant Use In Pregnancy: a Pilot RCT]. [clinicaltrials.gov/show/NCT02492009](http://clinicaltrials.gov/show/NCT02492009) (first received June 22, 2015).

**NCT02503553 {published data only}**

NCT02503553. Decision Aids in Cerebral Aneurysm Treatment. [clinicaltrials.gov/show/NCT02503553](http://clinicaltrials.gov/show/NCT02503553) (first received July 17, 2015).

**NCT02516449 {published data only}**

NCT02516449. Assessment of Shared Decision Making Aids in Asthma [Utility of Two Patients Decision Aids About Asthma Inhaled Controller Medication Use in Adult Patients With Asthma]. [clinicaltrials.gov/show/NCT02516449](http://clinicaltrials.gov/show/NCT02516449) (first received July 2, 2015).

**NCT02540044 {published data only}**

NCT02540044. Supporting Patient Care With Electronic Resource (SuPER) (SuPER) [Supporting Patient Care With Electronic Resource (SuPER): Efficacy of an Online Decision Aid for Patients Considering Biologic Therapy for Rheumatoid Arthritis]. [clinicaltrials.gov/show/NCT02540044](http://clinicaltrials.gov/show/NCT02540044) (first received September 1, 2015).

**NCT02611050 {published data only}**

NCT02611050. Treatment Decisions for Multi-vessel CAD [Treatment Decisions for Multi-vessel Coronary Artery Disease Patients]. [clinicaltrials.gov/show/NCT02611050](http://clinicaltrials.gov/show/NCT02611050) (first received November 9, 2015).

**Oostendorp 2011 {published data only}**

Oostendorp L, Ottevanger P, van der Graaf W, Stalmeier P. Assessing the information desire of patients with advanced cancer by providing information with a decision aid, which is evaluated in a randomized trial: a study protocol. *BMC Medical Informatics & Decision Making* 2011;**11**(9):1–9.

**Yu 2015 {published data only}**

Yu CH, Ivers NM, Stacey D, Rezmovitz J, Telner D, Thorpe K, et al. Impact of an interprofessional shared decision-making and goal-setting decision aid for patients with diabetes on decisional conflict: study protocol for a randomized controlled trial. *Trials* 2015;**16**(1):286.

**Additional references****Andrews 2013**

Andrews JC, Schünemann HJ, Oxman AD, Pottie K, Meerpohl JJ, Coello PA, et al. GRADE guidelines 15: Going from evidence to recommendation - determinants of a recommendation's direction and strength. *Journal of Clinical Epidemiology* 2013;**66**(7):726–35. [DOI: 10.1016/j.jclinepi.2013.02.003]

**Barry 2008**

Barry MJ, Wescott PH, Reifler EJ, Chang Y, Moulton BW. Reactions of potential jurors to a hypothetical malpractice suit. Alleging failing to perform a prostate-specific-antigen test. *Journal of Law, Medicine & Ethics* 2008;**Summer**: 396–402.

**Bekker 2003**

Bekker HL, Legare F, Stacey D, O'Connor A, Lemyre L. Is anxiety a suitable measure of decision aid effectiveness: a systematic review. *Patient Education and Counseling* 2003;**50**(3):255–62.

**Bennett 2010**

Bennett C, Graham ID, Kristjansson E, Kearing SA, Clay KF, O'Connor AM. Validation of a preparation for decision making scale. *Patient Education and Counseling* 2010;**78**(1): 130–33.

**Brehaut 2003**

Brehaut JC, O'Connor AM, Wood TJ, Hack TF, Siminoff L, Gordon E, et al. Validation of a decision regret scale. *Medical Decision Making* 2003;**23**(4):281–92.

**Brouwers 2010**

Brouwers M, Stacey D, O'Connor A. Knowledge creation: synthesis, tools and products. *CMAJ* 2010;**182**(2):E68–72.

**Brown 2015**

Brown JG, Joyce KE, Stacey D, Thomson MD. Patients or volunteers? The impact of motivation for trial participation on the efficacy of patient decision aids: a secondary analysis of a Cochrane Systematic Review. *Medical Decision Making* 2015;**35**(4):419–35.

**Charles 1997**

Charles C, Gafni A, Whelan T. Shared decision-making in the medical encounter: what does it mean?. *Social Science and Medicine* 1997;**44**(5):681–92.

**Charles 2010**

Charles C, Gafni A, Freeman E. Implementing shared treatment decision making and treatment decision aids: a cautionary tale. *Psychooncologia* 2010;**7**(2-3):243–55.

**Clinical Evidence 2013**

Clinical Evidence. How much do we know?. Available from: [clinicalevidence.bmj.com/x/set/static/cms/efficacy-categorisations.html](http://clinicalevidence.bmj.com/x/set/static/cms/efficacy-categorisations.html) (accessed 29 October 2013).

**Coyne 2013**

Coyne I, O'Mathuna DP, Gibson F, Shields L, Sheaf G. Interventions for promoting participation in shared decision-making for children with cancer. *Cochrane Database of Systematic Reviews* 2013, Issue 6. [DOI: 10.1002/14651858.CD008970.pub2]

**Degner 1992**

Degner LF, Sloan JA. Decision making during serious illness: what role do patients really want to play. *Journal of Clinical Epidemiology* 1992;**45**(9):941–50.

**Duncan 2010**

Duncan E, Best C, Hagen S. Shared decision making interventions for people with mental health conditions. *Cochrane Database of Systematic Reviews* 2010, Issue 1. [DOI: 10.1002/14651858.CD007297.pub2]

**Durand 2008**

Durand MA, Stiel M, Boivin J, Elwyn G. Where is the theory? Evaluating the theoretical frameworks described in decision support technologies. *Patient Education and Counseling* 2008;**71**(1):125–35.

**Durand 2014**

Durand MA, Carpenter L, Dolan H, Bravo P, Mann M, Bunn F, et al. Do interventions designed to support shared decision-making reduce health inequalities? A systematic review and meta-analysis. *PLOS One* 2014;**9**(4):1–14.

**Elwyn 2005**

Elwyn G, Hutchings H, Edwards A, Rapport F, Wensing M, Cheung WY, et al. The OPTION scale: measuring the extent that clinicians involve patients in decision-making tasks. *Health Expectations* 2005;**8**(1):34–42.

**Elwyn 2006**

Elwyn G, O'Connor A, Stacey D, Volk R, Edwards A, Coulter A, et al. Developing a quality criteria framework for patient decision aids: online international Delphi consensus process. *BMJ* 2006;**333**(7565):417.

**Elwyn 2013**

Elwyn G, I Scholl, Tietbohl C, Mann M, Edwards AGK, Clay C, et al. "Many miles to go...": A systematic review of the implementation of patient decision support interventions into routine clinical practice. *BMC: Medical Informatics and Decision Making* 2013;**13**(Suppl 2):S14.

**Gentles 2013**

Gentles SJ, Stacey D, Bennett C, Alshurafa M, Walter SD. Factors explaining the heterogeneity of effects of patient decision aids on knowledge of outcome probabilities: a systematic review sub-analysis. *Systematic Reviews* 2013;**2**:95.

**GRADEpro GDT [Computer program]**

GRADE Working Group, McMaster University. GRADEpro GDT. Version accessed prior to 21 March 2017. Hamilton (ON): GRADE Working Group, McMaster University, 2014.

**Gravel 2006**

Gravel K, Legare F, Graham ID. Barriers and facilitators to implementing shared decision-making in clinical practice: a systematic review of health professionals' perceptions. *Implementation Science* 2006;**1**:16.

**Hays 1993**

Hays RD, Sherbourne CD, Mazel RM. The RAND 36-item Health Survey 1.0. *Health Economics* 1993;**2**(3):217–27.

**Hibbard 1997**

Hibbard JH, Slovic P, Jewett JJ. Informing consumer decisions in health care: Implications from decision-making research. *Milbank Quarterly* 1997;**75**(3):395–414.

**Hibbard 2013**

Hibbard JH, Greene J. What the evidence shows about patient activation: better health outcomes and care experiences; fewer data on costs. *Health Affairs* 2013;**32**(2):207–14.

**Higgins 2011**

Higgins JPT, Green S (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from [www.cochrane-handbook.org](http://www.cochrane-handbook.org).

**Hoffman 2015**

Hoffmann TC, Del Mar C. Patients' expectations of the benefits and harms of treatments, screening, and tests: a systematic review. *Journal of the American Medical Association* 2015;**175**(2):274–86.

**Hollinghurst 2010**

Hollinghurst S, Emmett C, Peters TJ, Watson H, Fahey T, Murphy DJ, et al. Economic evaluation of the DIAMOND randomized trial: cost and outcomes of 2 decision aids for mode of delivery among women with previous caesarian section. *BMJ* 2010;**30**:453–63.

**IPDAS 2005a**

International Patient Decision Aid Standards Collaboration. Background Document. 2005. [ipdas.ohri.ca/IPDAS\\_Background.pdf](http://ipdas.ohri.ca/IPDAS_Background.pdf) (accessed 29 Oct 2013).

**IPDAS 2005b**

International Patient Decision Aid Standards Collaboration. IPDAS Voting Document - 2nd Round. 2005. [ipdas.ohri.ca/IPDAS\\_Second\\_Round.pdf](http://ipdas.ohri.ca/IPDAS_Second_Round.pdf) (accessed 29 Oct 2013).

**IPDAS 2013**

Volk RJ, Llewellyn-Thomas H, Stacey D, Elwyn G. Ten years of the International Patient Decision Aid Standards collaboration: evolution of the core dimensions for assessing the quality of patient decision aids. *BMC: Medical Informatics and Decision Making* 2013;**13**(Suppl 2):S1.

**Joseph-Williams 2013**

Joseph-Williams N, Newcombe R, Politi M, Durand MA, Sivell S, Stacey D, et al. Toward minimum standards for certifying patient decision aids: a modified Delphi consensus process. *Medical Decision Making* 2014;**34**(6):699–710. [DOI: 10.1177/0272989X13501721]

**Kiesler 2006**

Kiesler DJ, Auerbach SM. Optimal matches of patient preferences for information, decision-making and interpersonal behavior: evidence, models and interventions. *Patient Education and Counseling* 2006;**61**(3):319–41.

**LeBlanc 2010**

LeBlanc A, Kenny DA, O'Connor AM, Legare F. Decisional conflict in patients and their physicians: a dyadic approach to shared decision making. *Medical Decision Making* 2010;**29**(1):61–8.

**Legare 2008b**

Legare F, Ratte S, Gravel K, Graham ID. Barriers and facilitators to implementing shared decision-making in clinical practice: update of a systematic review of health professionals' perceptions. *Patient Education and Counseling* 2008;**73**(3):526–35.

**Legare 2010**

Legare F, Ratte S, Stacey D, Kryworuchko J, Gravel K, Graham ID, et al. Interventions for improving the adoption of shared decision making by healthcare professionals. *Cochrane Database of Systematic Reviews* 2010, Issue 5. [DOI: 10.1002/14651858.CD006732.pub2]

**Legare 2014**

Legare F, Stacey D, Turcotte S, Cossi MJ, Kryworuchko J, Graham ID, et al. Interventions for improving the adoption of shared decision making by healthcare professionals. *Cochrane Database of Systematic Reviews* 2014, Issue 9. [DOI: 10.1002/14651858.CD006732.pub3]

**Makoul 2006**

Makoul G, Clayman ML. An integrative model of shared decision making in medical encounters. *Patient Education and Counseling* 2006;**60**(3):301–12.

**Michie 2002**

Michie S, Dormandy E, Marteau TM. The multi-dimensional measure of informed choice: a validation study. *Patient Education and Counseling* 2002;**48**(1):87–91.

**Mulley 1995**

Mulley A. Outcomes research: implications for policy and practice. In: Smith R, Delamother T editor(s). *Outcomes in Clinical Practice*. London: BMJ Publishing Group, 1995.

**Munro 2016**

Munro S, Stacey D, Lewis KB, Bansback N. Choosing treatment and screening options congruent with values: do decision aids help? Sub-analysis of a systematic review. *Patient Education & Counseling* 2016;**99**(4):491–500.

**NCGC/NICE 2012**

National Clinical Guideline Centre. Patient experience in adult NHS services: improving the experience of care for people using adult NHS services. 2012. [www.nice.org.uk/nicemedia/live/13668/58283/58283.pdf](http://www.nice.org.uk/nicemedia/live/13668/58283/58283.pdf). London, UK: The Author, (accessed prior to 27 March 2017).

**O'Connor 1995**

O'Connor AM. Validation of a decisional conflict scale. *Medical Decision Making* 1995;**15**(1):25–30.

**O'Connor 1998b**

O'Connor AM, Tugwell P, Wells GA, Elmslie T, Jolly E, Hollingworth G, et al. A decision aid for women considering hormone therapy after menopause: decision support framework and evaluation. *Patient Education and Counseling* 1998;**33**(3):267–79.

**O'Connor 2007**

O'Connor AM, Wennberg JE, Legare F, Llewellyn-Thomas HA, Moulton BW, Sepucha KR, et al. Toward the 'tipping point': decision aids and informed patient choice. *Health Affairs* 2007;**26**(3):716–25.

**RevMan 2014 [Computer program]**

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

**RNAO 2009**

Registered Nurses' Association of Ontario. Decision support for adults living with chronic kidney disease. 2009. [rnao.ca/bpg/guidelines/decision-support-adults-living-chronic-kidney-disease](http://rnao.ca/bpg/guidelines/decision-support-adults-living-chronic-kidney-disease). Toronto, Ontario: The Author, (accessed prior to 27 March 2017).

**Rothert 1987**

Rothert M, Talarczyk GJ. Patient compliance and the decision making process of clinicians and patients. *Journal of Compliance in Health Care* 1987;**2**(1):55–71.

**Schünemann 2011**

Schünemann HJ, Oxman AD, Higgins JPT, Vist GE, Glasziou P, Guyatt GH. Chapter 11: Presenting results

and 'Summary of findings' tables. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from [www.handbook.cochrane.org](http://www.handbook.cochrane.org).

**Sepucha 2013**

Sepucha KR, Borkhoff CM, Lally J, Levin CA, Matlock DD, Ng CJ, et al. Establishing the effectiveness of patient decision aids: key constructs and measurement instruments. *BMC: Medical Informatics and Decision Making* 2013;**13** (Suppl 2):S12.

**Spielberger 1970**

Spielberger CD, Gorsuch RL, Lushene RE. *Manual for the State-Trait Anxiety Inventory (Self-evaluations questionnaire)*. Palo Alto, CA: Consulting Psychologists Press, 1970.

**Stewart 1992**

Stewart AL, Ware JE Jr (editors). *Measuring Functioning and Well-being: The Medical Outcomes Study Approach*. Durham NC: Duke University Press, 1992.

**Trenaman 2014**

Trenaman L, Stirling B, Bansback N. The cost-effectiveness of patient decision aids: a systematic review. *Healthcare* 2014;**2**(4):2510257.

**Trenaman 2016**

Trenaman L, Selva A, Desroches S, Singh K, Bissonnette J, Bansback N, et al. A measurement framework for adherence in patient decision aid trials applied in a systematic review subanalysis. *Journal of Clinical Epidemiology* 2016;**77**: 15–23.

**Trikalinos 2014**

Trikalinos TA, Wieland LS, Adam GP, Zgodic A, Ntzani EE. *Decision Aids for Cancer Screening and Treatment*. Rockville, MD: Agency for Healthcare Research and Quality, 2014.

**Washington State 2016**

Washington State Health Authority. Patient decision aid certification criteria. 2016. [www.hca.wa.gov/hw/Documents/sdm'cert'criteria.pdf](http://www.hca.wa.gov/hw/Documents/sdm'cert'criteria.pdf) (accessed prior to 27 March 2017).

**Weston 2001**

Weston WW. Informed and shared decision-making: the crux of patient-centered care. *CMAJ* 2001;**165**(4):438–9.

**References to other published versions of this review****O'Connor 1999b**

O'Connor AM, Rostom A, Fiset V, Tetroe J, Entwistle V, Llewellyn-Thomas H, et al. Decision aids for patients facing health treatment or screening decisions: systematic review. *BMJ* 1999;**319**(7212):731–4.

**O'Connor 2001**

O'Connor AM, Stacey D, Rovner D, Holmes-Rovner M, Tetroe J, Llewellyn-Thomas H, et al. Decision aids for people facing health treatment or screening decisions. *Cochrane Database of Systematic Reviews* 2001, Issue 3. [DOI: 10.1002/14651858.CD001431]

**O'Connor 2003**

O'Connor AM, Stacey D, Rovner D, Holmes-Rovner M, Tetroe J, Llewellyn-Thomas H, et al. Decision aids for people facing health treatment or screening decisions. *Cochrane Database of Systematic Reviews* 2003, Issue 1. [DOI: 10.1002/14651858.CD001431]

**O'Connor 2009b**

O'Connor AM, Bennett C, Stacey D, Barry M, Col NF, Eden KB, et al. Decision aids for people facing health treatment or screening decisions. *Cochrane Database of Systematic Reviews* 2009, Issue 3. [DOI: 10.1002/14651858.CD001431.pub2]

**Stacey 2011**

Stacey D, Bennett CL, Barry MJ, Col NF, Eden KB, Holmes-Rovner M, et al. Decision aids for people facing health treatment or screening decisions. *Cochrane Database of Systematic Reviews* 2011, Issue 10. [DOI: 10.1002/14651858.CD001431.pub3]

**Stacey 2014b**

Stacey D, Legare F, Col NF, Bennett CL, Barry MJ, Eden KB, et al. Decision aids for people facing health treatment or screening decisions. *Cochrane Database of Systematic Reviews* 2014, Issue 1. [DOI: 10.1002/14651858.CD001431.pub4]

\* Indicates the major publication for the study

## CHARACTERISTICS OF STUDIES

### Characteristics of included studies [ordered by study ID]

#### Allen 2010

Methods	Cluster-randomized to decision aid vs usual care
Participants	398 + 414 men considering prostate cancer screening in the USA
Interventions	DA: computer tailored programme on clinical problem, outcome probabilities, explicit values clarification, others' opinion and guidance (step-by-step process for making the decision; interactive computer programme: inherently guided the patient through the decision aid and decision making process), tailored printout given to patients to promote discussion with others (practitioner, significant others) Comparator: no intervention
Outcomes	Primary outcomes: decisional status, knowledge, decision self-efficacy, decisional consistency Secondary outcomes: desire for involvement in decision making, decisional conflict, preferred options Outcomes assessed pre- and postintervention
Notes	-

#### *Risk of bias*

#### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Sites were blocked on size and percent of male employees and randomly assigned by computer-generated random numbers to condition within blocks" (p 2173, Setting)
Allocation concealment (selection bias)	Unclear risk	The study does not address this criterion.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	The study does not address this criterion.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes measured were not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data and low rate of attrition that was consistent between groups
Selective reporting (reporting bias)	Unclear risk	No mention of protocol

Other bias	Low risk	Intervention delivery: mention of money incentive to complete paperwork, but was judged to have no effect on outcomes measured (p 2175)
------------	----------	-----------------------------------------------------------------------------------------------------------------------------------------

## Arterburn 2011

Methods	Randomized to decision aid vs usual care
Participants	75 + 77 participants considering bariatric surgery in the USA
Interventions	DA: booklet + video on options' outcomes, clinical problem, outcome probabilities, others' opinion, guidance (list of questions to discuss with clinician) Comparator: usual care (general information pamphlets on clinical problem)
Outcomes	Primary outcomes: knowledge, values, values concordance Secondary outcomes: treatment preference, decisional conflict, decisional self-efficacy, proportion undecided Primary outcomes assessed at baseline, postintervention and 3 months follow-up; secondary outcomes assessed at baseline and postintervention
Notes	-

**Risk of bias****Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"[U]sed computer-assisted, block randomisation process to ensure balanced allocation of participants" (p 1670, Participants and randomization)
Allocation concealment (selection bias)	Unclear risk	No mention of allocation concealment and no mention of impact on study
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"[S]tudy was not blinded" (p 1670, Participants and randomization); no mention of impact on study
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subject to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Measures: mentioned 4 choices for treatment preference (surgery, drug therapy, diet and/or exercise programme and unsure) but only reported on surgery and unsure options (p 1671); minimal attrition that was consis-



**Arterburn 2011** (Continued)

		tent between groups
Selective reporting (reporting bias)	Unclear risk	No mention of study protocol or trial registration; all pre-specified outcomes included
Other bias	Low risk	The study appears to be free of other sources of bias

**Auvinen 2004**

Methods	Randomized to decision aid vs usual care
Participants	103 + 100 men newly diagnosed with prostate cancer in Finland
Interventions	DA: pamphlet patient decision aid created for study on options' outcomes, outcome probability, guidance Comparator: usual care by clinical guideline
Outcomes	Primary outcome: uptake of options Secondary outcome: participation in decision making Other outcomes (from Huang 2014): death (5 years), disease-free survival (10-years), biochemical failure (serum PSA elevation) (5 years), biochemical failure-free survival (5 years), disease progression (5 years), disease progression-free survival (5 years) (data from 104 + 106 men)
Notes	-

***Risk of bias***

***Risk of bias***

<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Auvinen 2001, p 2: "randomized centrally, using software based on a random number generator"; no blocking used Auvinen 2004, (primary study), p 1: "randomized using a computer algorithm based on random numbers"
Allocation concealment (selection bias)	Unclear risk	Auvinen 2001, p 2, Patients and Methods: randomized centrally at the Finnish Cancer Registry Auvinen 2004, (primary study), p 1: randomized centrally Comment: central allocation confers low risk
Blinding of participants and personnel (performance bias) All outcomes	High risk	Auvinen 2001, p 3: "recognized carry-over effect because same physician in charge for intervention and control groups, diminish

**Auvinen 2004** (Continued)

		contrast between groups, as these physicians were more motivated to inform patients than those physicians not participating” Auvinen 2004 (primary study): no blinding but primary outcome is choice of treatment for prostate, objectively recorded. But unsure how physicians may have influenced decisions
Blinding of outcome assessment (detection bias) All outcomes	Low risk	No blinding but primary outcome is choice of treatment for prostate, objectively recorded
Incomplete outcome data (attrition bias) All outcomes	Low risk	Auvinen 2001, p 3: flow-chart “Imbalance in the numbers of patients between the arms within two hospitals. Not expected to affect the results in any way”; “some participants refused to give informed consent, health deterioration, not seen by urologist” (p 4) Auvinen 2004 (primary study), p 2: flow diagram and results; low attrition and consistent between groups
Selective reporting (reporting bias)	Unclear risk	No indication that trial registered in central trials registry Auvinen 2001, p 2: “The study protocol was approved by an ethical committee in each participating hospital” Auvinen 2004 (primary study), p 1: “The study protocol was approved by the institutional review board at each participating hospital”
Other bias	Low risk	Appears to be free of other potential biases

**Barry 1997**

Methods	Randomized to decision aid vs usual care
Participants	104 + 123 patients considering benign prostatic hyperplasia treatment in the USA
Interventions	DA: Health Dialog interactive videodisc on options’ outcomes, clinical problem, outcome probability, others’ opinion Comparator: usual care using general information on the clinical problem
Outcomes	Primary outcome: knowledge Secondary outcomes: uptake of option, satisfaction with DM process, satisfaction with decision, interest in DM, general health outcomes, condition specific health outcomes

**Barry 1997** (Continued)

Notes	-	
<b><i>Risk of bias</i></b>		<b><i>Risk of bias</i></b>
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	"Stratified by study site in concealed blocks of 10" (p 2)
Allocation concealment (selection bias)	Low risk	Study coordinator opening serially numbered, opaque, sealed envelopes (p 2)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	No blinding but phase 1 eliminated risk of contamination
Blinding of outcome assessment (detection bias) All outcomes	Low risk	No blinding but phase 1 eliminated risk of outcome assessor interfering with decision
Incomplete outcome data (attrition bias) All outcomes	Low risk	Patient accrual and follow-up reported; post-randomization withdrawals could have biased the results (more in intervention group) - however they reported no evidence of a differential effect of the study group (p 3)
Selective reporting (reporting bias)	Unclear risk	No indication that trial registered in central trials registry
Other bias	Low risk	Appears to be free of other potential biases

**Bekker 2004**

Methods	Randomized to detailed vs routine consultation
Participants	59 + 58 pregnant women who have received a maternal serum screening positive test result for Down syndrome in the UK
Interventions	DA (in consult): decision analysis plus routine consultation on options' outcomes, clinical problem, outcome probability, values clarification, guidance/coaching Comparator: routine consultation on options' outcomes, outcome probability
Outcomes	Primary outcome: anxiety Secondary outcomes: uptake of option, knowledge, decisional conflict, informed decision making, satisfaction with consultation, consultation length

**Bekker 2004** (Continued)

Notes	-	
<i>Risk of bias</i>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Bekker 2003, p 2 - section 2.3 Sample and Procedure: "randomly allocated... using previously numbered... envelopes" Bekker 2004 (primary study), p 3: "Participants were randomly allocated by previously numbered envelopes"; does not mention how sequence was generated
Allocation concealment (selection bias)	Low risk	Bekker 2003, p 2 - section 2.3 Sample and Procedure: "Using previously numbered, sealed, opaque envelopes" Bekker 2004 (primary study), p 3: previously numbered, sealed, opaque envelopes
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Participants blinded, personnel not blinded. Same personnel did control & intervention. Tape recorded sessions to ensure no bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Bekker 2003 flow diagram indicates post-randomization attrition with more attrition in decision aid group; no discussion on implications of attrition Bekker 2004 (primary study), p 4: results/flow diagram; baseline characteristics not included
Selective reporting (reporting bias)	Unclear risk	Bekker 2003: the coding frame was developed from literature. Does not mention protocol Bekker 2004 (primary study): no information provided about central trials registry
Other bias	Unclear risk	Bekker 2003: does not directly address baseline characteristics of participants Bekker 2004 (primary study): appears to be free of other potential biases

## Bernstein 1998

Methods	Randomized to decision aid vs usual care
Participants	65 + 53 patients with coronary artery disease considering revascularization surgery in the USA
Interventions	DA: Health Dialog video on options' outcomes, clinical problem, outcome probability, others' opinion Comparator: usual care (no information provided)
Outcomes	Primary outcome: satisfaction with decision and decision making process Secondary outcomes: uptake of option, knowledge, satisfaction with care, general health outcomes, condition specific health outcomes
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Randomization was stratified by study site in blocks of 10" (p 3)
Allocation concealment (selection bias)	Low risk	"[R]andomization performed by a study coordinator opening opaque, sealed envelopes at study headquarters" (p 3)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Neither subjects nor study staff were blinded to treatment assignment - could lead to different satisfaction ratings based on knowing the treatment received
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured
Incomplete outcome data (attrition bias) All outcomes	Low risk	Flow diagram (p 3); low attrition of eligible participants randomized and consistent between group
Selective reporting (reporting bias)	Unclear risk	No information provided indicating trial was included in central trials registry
Other bias	Low risk	Appears to be free of other potential biases

## Berry 2013

Methods	Randomized to decision aid vs usual care
Participants	266 + 228 men considering prostate cancer treatment in the USA
Interventions	DA: interactive web based video on options' outcomes, clinical problem, outcome probabilities, others' opinion, guidance (list of questions to ask doctor and automated summary) Comparator: usual care
Outcomes	Primary outcome: decisional conflict Secondary outcome: preferred/actual treatment choice (pre- and post-DA), proportion undecided Other outcomes (Bosco 2012): choice concordance (6 months post-DA). (Data from 239 + 209 men)
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Methods section- second paragraph, p 3: "Participants were randomized automatically by the P3P application to study groups (1:1 using a simple randomization scheme with no blocking)"
Allocation concealment (selection bias)	Low risk	Methods section, p 3: "Participants were randomized automatically by the P3P application to study groups"
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Participants were not blinded and study does not address the effect on the results
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear whether outcome assessors are blinded, but outcomes are not subject to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used intention-to-treat analysis and low dropout (p 4)
Selective reporting (reporting bias)	Low risk	Protocol made available
Other bias	Unclear risk	Was a multicentre trial which could have lead to contamination, protocol violation and biased questionnaire completion

## Bjorklund 2012

Methods	Randomized to decision aid vs usual care
Participants	236 + 247 women less than 11 weeks pregnant considering Down syndrome screening in Sweden
Interventions	DA: linear video on options' outcomes, clinical problem, outcome probabilities, others' opinion, and guidance (step-by-step process for making the decision) Comparator: usual care using pamphlet
Outcomes	Primary outcomes: knowledge (post-DA), attitude (post-DA), uptake of combined ultrasound and biochemical screening (post-DA) Secondary outcomes: values congruent with chosen option (post-DA)
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"The midwife allocated the participants randomly by sealed envelopes" (p 391) but does not state the actual sequence generation method
Allocation concealment (selection bias)	Low risk	Used sealed envelopes, "prepared, sequentially coded and distributed to the maternity units by the research group" (p 391)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"It was not possible to blind neither [sic] the midwives nor the participants due to the characteristics of the intervention" (p 395). The study does not address the effects of this on the results
Blinding of outcome assessment (detection bias) All outcomes	Low risk	No blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No mention of why some participants' data were excluded in Tables 2, 3 and 4
Selective reporting (reporting bias)	Unclear risk	No mention of study protocol
Other bias	Low risk	Appears to be free of other sources of bias

## Bozic 2013

Methods	Randomized to decision aid vs usual care
Participants	95 + 103 participants with hip and/or knee osteoarthritis considering hip/knee surgery
Interventions	DA: DVD and booklet on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, others' opinions, and guidance/coaching with health coach Comparator: usual care using pamphlet
Outcomes	Primary outcomes: informed decision/knowledge (pre, immediately post, and 6 weeks follow-up) Secondary outcomes: preferred treatment choice (pre and immediately post), patient and provider satisfaction (immediately post), length of consultation time
Notes	Trial registration: NCT01492257

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The randomization was blocked with use of random permuted blocks in groups of four, six, or eight to help ensure that the groups were balanced" (p 1634)
Allocation concealment (selection bias)	Low risk	"Patients were randomized to either the intervention group or the control group with use of the sealed envelop method" (p 1634)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"[S]urgeons were not blinded to the intervention" (p 1635). Knowing the allocation of participants, surgeons' favourable scoring could be due to greater investment in decision-making. Insufficient information to make a judgment
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcomes are objectively measured and not subject to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	62% (123/198) retention rate therefore high attrition rate - however the attrition was balanced between groups
Selective reporting (reporting bias)	Low risk	Protocol available
Other bias	Low risk	Appears to be free of other sources of bias



# Brazell 2014

Methods	Randomized to DA + standard counselling vs usual care + standard counselling
Participants	53 + 51 women presenting for the management and treatment of pelvic organ prolapse
Interventions	DA: paper-based or web-based DA on clinical problem, options' outcomes, outcome probabilities, patient stories and standard counselling Comparator: standard counselling alone
Outcomes	Primary outcomes: decisional conflict (immediately postconsultation) Secondary outcomes: choice (3 months after making decision), decisional regret (3 months after making decision)
Notes	-

## *Risk of bias*

## *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Patients were randomized 1:1 using a random numbers table in blocks of 6" (p 231)
Allocation concealment (selection bias)	Unclear risk	Insufficient information provided to make judgment
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information provided to make judgment
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information provided to make judgment
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	High attrition but balanced between groups: "39 randomized subjects were either missed by the research assistant at their new patient visit and thus did not receive a DCS questionnaire to complete or they canceled their appointments and did not reschedule a new one" (p 233). There was a 48% (50/104) attrition rate for Decisional Regret measures
Selective reporting (reporting bias)	Low risk	Trial registered
Other bias	High risk	Risk of contamination due to same physicians in both groups. Also, outcomes measured after the PtDA and physician consult

## Chabrera 2015

Methods	Randomized to DA vs usual care
Participants	73 + 74 men recently diagnosed with prostate cancer considering treatment options
Interventions	DA: 2-part decision support booklet with clinical problem, options' outcomes, outcome probabilities, patient stories, explicit values clarification, and guidance Comparator: usual care
Outcomes	Primary outcomes: knowledge, decisional conflict, satisfaction with decision-making process Secondary outcome: coping Outcomes assessed at 3 months postintervention
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"[S]tudy participants were randomized into 1 of 2 arms using a computer-generated random list with unequal blocks" (p E44)
Allocation concealment (selection bias)	Unclear risk	Insufficient information provided to make judgement
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information provided to make judgement
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information provided to make judgement
Incomplete outcome data (attrition bias) All outcomes	Low risk	Balanced attrition in both groups
Selective reporting (reporting bias)	Unclear risk	No protocol provided; trial not registered
Other bias	Unclear risk	Prostate cancer in Catalonia is common; however, only 147 were recruited for this trial (p E44)

## Chambers 2012

Methods	Randomized to DA vs usual care
Participants	74 + 77 healthcare workers who did not receive the influenza vaccine considering receiving the vaccine in Canada
Interventions	DA: web-based DA on options' outcomes, clinical problem, outcome probabilities, explicit values clarification and guidance Comparator: usual care using pamphlet
Outcomes	Primary outcomes: confidence in decision (post-DA) Secondary outcomes: impact on immunization intent (post-DA), proportion undecided
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The randomization list was generated using the randomization function in Excel 2002 (version 10.6856.6856 SP3)" (p 199)
Allocation concealment (selection bias)	Low risk	"The list was imported from Excel into a Microsoft SQL Server database. The online application would sequentially assign a random identification number and their decision aid status (seeing the decision aid or not) from the randomization list when users logged into the survey." (p 199)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not reported whether or not they were blinded during the course of the intervention
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Questionnaire scores are objective and not subject to interpretation
Incomplete outcome data (attrition bias) All outcomes	High risk	65% completion rate in intervention arm and 77% completion rate in control arm: attrition could be different where the respondents and non-respondents are different
Selective reporting (reporting bias)	Low risk	Protocol available
Other bias	Unclear risk	Figure 1 numbers for exclusion are not logical

## Clancy 1988

Methods	Randomized to decision aid vs usual care
Participants	753 + 263 health physicians considering Hep B vaccine in the USA
Interventions	DA: pamphlet on options' outcomes, clinical problem, outcome probability, explicit values clarification (personal decision analysis), guidance/coaching Comparator: usual care (no information provided)
Outcomes	Uptake of option
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random numbers table; all incoming residents were assigned to Group 2 (non-randomized residents identified as subgroup) (p 2)
Allocation concealment (selection bias)	Unclear risk	No information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No blinding of participants or personnel. Did not report on how this may affect their findings
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but decisions for screening were retrieved from health records (objective data)
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Flow chart not included. Insufficient information to make a judgment
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	High risk	Potential selection bias - non-randomized residents were added to group 2 and therefore potential unbalanced distribution (p 287) Low response rate among those offered decision analysis

## Davison 1997

Methods	Randomized to decision aid + audio-taped consultation vs usual care
Participants	30 + 30 men with prostate cancer considering treatment in Canada
Interventions	DA: written + audiotape consultation of options' outcomes, clinical problem, outcome probability, others' opinion Comparator: usual care (general information pamphlets on clinical problem)
Outcomes	Primary outcomes: role in decision making Secondary outcomes: anxiety, depression
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The group to which subjects were assigned was predetermined by a block randomization procedure. This ensured there were an equal number of subjects in both groups for each physician." (p 5, Data collection)
Allocation concealment (selection bias)	Unclear risk	Not mentioned; group assignment predetermined by block randomization procedure (p 5)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No blinding; study does not report on how the results could be influenced by lack of blinding
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Unclear blinding and whether outcomes could be affected by unblinded assessor
Incomplete outcome data (attrition bias) All outcomes	Low risk	No flow diagram; p 12 explains why certain men did not listen to audiotape. All men approached by study investigator agreed to participate; only 1 man refused to complete the second set of questionnaires
Selective reporting (reporting bias)	Unclear risk	Protocol not mentioned
Other bias	Low risk	Appears to be free of other sources of bias; similar baseline characteristics

## De Achaval 2012

Methods	Randomized to detailed vs simple vs usual care
Participants	70 + 70 + 71 patients diagnosed with knee osteoarthritis considering treatment in the USA
Interventions	Complex DA: video booklet + interactive joint analysis on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, others' opinion and guidance (list of questions) Comparator DA: video booklet on options' outcomes, clinical problem, outcome probabilities, others' opinion and guidance (list of questions) Comparator: usual care receiving generic booklet
Outcomes	Decisional conflict (baseline and postintervention)
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer generated list with uneven blocks (p 231)
Allocation concealment (selection bias)	Low risk	Numbered, sealed and opaque envelopes (p 231)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Likely not blinded, but low threat of bias in study (p 231)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Participants were not blinded but outcome was objectively measured (p 231)
Incomplete outcome data (attrition bias) All outcomes	Low risk	3 dropouts; missing data effect size unlikely to have significant impact on study outcome
Selective reporting (reporting bias)	Unclear risk	Protocol not available
Other bias	Low risk	Appears to be free of other sources of bias

## Dolan 2002

Methods	Randomized to decision aid vs usual care
Participants	50 + 47 average risk for colorectal cancer considering screening in the USA
Interventions	DA: computer with analytic hierarchy process on options' outcomes, clinical problem, outcome probability, explicit values clarification, guidance/coaching Comparator: usual care with information on options, clinical problem
Outcomes	Primary outcomes: uptake of option, decisional conflict Secondary outcomes: role in decision making
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"[R]andomization schedules were created using a computer random number generator" (p 2, Study interventions)
Allocation concealment (selection bias)	Low risk	Computer-based (p 2, Study interventions)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding of participants. All patient interviews in both the experimental and control groups were done by the same investigator, unclear on how this could contribute to risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	See flow diagram - low attrition
Selective reporting (reporting bias)	Unclear risk	Nothing specifically mentioned re study protocol
Other bias	Low risk	Appears to be free of other sources of bias

## Evans 2010

Methods	Randomized to online decision aid vs paper decision aid vs questionnaire vs usual care
Participants	129 + 126 + 127 + 132 men considering PSA screening in Wales

Interventions	DA: online programme on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, others' opinion, guidance (interactive computer programme; summary) Comparator: paper version of online DA on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, others' opinion, guidance (interactive computer programme; summary) Comparator: received a questionnaire Comparator: received nothing
Outcomes	Primary outcomes: knowledge (post-DA) Secondary outcomes: attitude (post-DA), intention to undergo PSA testing (post-DA), anxiety (post-DA), uptake of PSA test (post-DA), total decisional conflict
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"[A] random sample of 100 men was selected from the list." "The process ensured individual level randomization" (p 4, Recruitment process)
Allocation concealment (selection bias)	Low risk	"[A]ffirmative consent forms from each practice were transferred to the research officer who allocated each participant with a number provided remotely by the trial statistician to ensure concealment" (p 4, Recruitment process)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	The study does not address this outcome
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	See flow diagram indicating high attrition consistently across groups
Selective reporting (reporting bias)	Low risk	Registered as a trial
Other bias	Low risk	The study appears free of other sources of bias



## Fagerlin 2011

Methods	Decision aid vs delayed intervention vs control
Participants	382 + 159 + 100 women with an elevated 5-year risk of breast cancer considering breast cancer prevention medication in the USA
Interventions	DA: tailored DA on options' outcomes, clinical problem, outcome probabilities, and explicit values clarification Comparator 1: given DA after 3-month follow-up Comparator 2: given DA after all outcome measures were taken
Outcomes	Decisional conflict (post-DA), behavioural intent (post-DA), actual behaviour (post-DA), proportion undecided, perception of benefits (post-DA), perception of risk (post-DA) Other outcomes: <ul style="list-style-type: none"> <li>Banegas 2013: decisional conflict (post-DA) (data from 690 + 160 + 162 women), proportion undecided (3 months)</li> <li>Korfage 2013: knowledge (immediately post and 3 months post-DA), attitudes (immediately post and 3 months post-DA), behavioural intent (post-DA), actual behaviour (3 months post-DA), informed decision defined as "participants with sufficient knowledge about chemoprevention behavior, whose attitudes were concordant with their intentions or decisions to engage in chemoprevention behavior" (data from 383 + 102 + 100 women).</li> </ul>
Notes	Primary outcome was not specified

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random sequence generation was provided by the author
Allocation concealment (selection bias)	Low risk	Central and web-based allocation
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding - using an online decision aid would have avoided control participants accessing the decision aid
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Does not report exclusions; inadequate reporting on participant flow through the study to determine risk for attrition bias or incomplete outcome data
Selective reporting (reporting bias)	Unclear risk	No mention of study protocol

**Fagerlin 2011** (Continued)

Other bias	Low risk	Appears to be free of other sources of bias
------------	----------	---------------------------------------------

**Fraenkel 2007**

Methods	Randomized to decision aid vs usual care
Participants	47 + 40 patients with knee pain considering treatment options in the USA
Interventions	DA: interactive computer tool options' outcomes, outcome probability, explicit values clarification Comparator: usual care using the Arthritis Foundation information pamphlet
Outcomes	Decisional self-efficacy, preparation for decision making
Notes	Primary outcome was not specified

<i><b>Risk of bias</b></i>		<i><b>Risk of bias</b></i>
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Computer-generated randomization sequence (p 2)
Allocation concealment (selection bias)	Unclear risk	No information provided; computer generated
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No blinding but study does not report if it had an impact on the outcomes measured
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Low risk of attrition bias - outcome data for all 40 controls and 44 of 47 intervention (p 3, Results)
Selective reporting (reporting bias)	Unclear risk	No information provided; no indication of trial was registered centrally
Other bias	Low risk	Appears to be free of other potential biases

## Fraenkel 2012

Methods	Cluster-randomized control trial of clinics to decision aid versus usual care
Participants	69 + 66 patients with nonvalvular atrial fibrillation considering anticoagulation with aspirin or warfarin
Interventions	DA: computer-based tool on options' outcomes, clinical problem, options' probabilities, guidance, explicit values clarification Comparator: control arm (no further information provided)
Outcomes	Primary outcomes: feeling informed and having clear values (baseline, immediately post) Secondary outcomes: knowledge (baseline, immediately post), accuracy of risk (baseline, immediately post), anxiety (baseline, immediately post), worry (baseline, immediately post), rationale for preferred treatment (during the encounter - DA group only), discussion of related outcomes (during the encounter as captured on audiotape), change in treatment plan (post intervention), anxiety, accurate risk expectations (stroke, bleeding)
Notes	Trial registration NCT00829478

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Inadequate information on random sequence generation
Allocation concealment (selection bias)	Unclear risk	inadequate information on allocation concealment
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"To avoid contamination, participants were randomized at the level of the firm so that all participants in one firm received the intervention, and all participants in the second firm were included in the control arm" (p 1435)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"An interviewer blinded to the participant's group assignment reassessed the primary and secondary outcomes after participant's primary care visit" (p 1436)
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Does not appear to be incomplete outcome data; flow diagram does not report participation beyond randomization
Selective reporting (reporting bias)	Low risk	Protocol available
Other bias	Low risk	Does not appear to be any other potential sources of bias

## Frosch 2008a

Methods	Randomized to decision aid vs. decision aid + chronic disease trajectory vs chronic disease trajectory vs usual care (Internet information)
Participants	155 + 152 + 153 + 151 men considering prostate cancer screening
Interventions	DA: information on options' outcomes, clinical problem, outcome probabilities, others' opinions Comparator 1: information on options' outcomes, clinical problem, outcome probabilities, others' opinions, explicit values clarification (utilities for outcomes associated with prostate cancer) Comparator 2: explicit values clarification (utilities for outcomes associated with prostate cancer) Comparator 3: usual care using public information on prostate cancer screening on American Cancer Society and Centers for Disease Control and Prevention websites 2005-2006
Outcomes	Primary outcomes: knowledge, actual option, decisional conflict Secondary outcomes: concern about prostate cancer, treatment preference if prostate cancer diagnosed
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer algorithm randomly assigned participants to the 4 study groups
Allocation concealment (selection bias)	Low risk	Revealed after signed consent and completed baseline measures
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Accessed a secure Internet site that hosted all study materials; participants had unlimited access to assigned intervention, unclear blinding of personnel
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were measured via questionnaires and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used intention-to-treat analysis; imputed missing data for participants who did not complete follow-up assessments; minimal attrition
Selective reporting (reporting bias)	Unclear risk	No indication of published protocol

**Frosch 2008a** (Continued)

Other bias	Low risk	Appears to be free of other potential biases
------------	----------	----------------------------------------------

**Gattellari 2003**

Methods	Randomized to decision aid vs usual care
Participants	126 + 122 men considering PSA testing in Australia
Interventions	DA: pamphlet on options' outcomes, clinical problem, outcome probability, explicit values clarification Comparator: usual care using brief information on screening test and chances of false-positive results
Outcomes	Preferred option, knowledge, decisional conflict, accurate risk perceptions, perceived ability to make an informed choice
Notes	Primary outcome was not specified

<i><b>Risk of bias</b></i>		<i><b>Risk of bias</b></i>
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Pre-randomized code - no further information (p 1)
Allocation concealment (selection bias)	Low risk	Pre-randomized code unobtrusively marked on envelopes (p 1)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Consenting men were blinded to allocation, but unclear if personnel were blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Pre-test characteristics included. Flow chart not included and reasons for attrition not mentioned; some attrition but balanced between groups
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Low risk	Appears to be free of other potential biases

## Gattellari 2005

Methods	Randomized to decision aid booklet vs decision aid video vs usual care
Participants	140 + 141 + 140 men considering PSA testing in Australia
Interventions	DA: pamphlet on options' outcomes, clinical problem, outcome probability, explicit values clarification Comparator 1: video on clinical problem, outcome probability, others' opinion Comparator 2: usual care using brief information on screening test and chances of false-positive results
Outcomes	Preferred option, knowledge, decisional conflict, perceived ability to make an informed choice
Notes	Primary outcome was not specified

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Unique identification codes assigned to participants according to date and time enrolled into the interventional component of the study. Block randomization of identification codes then performed via computer software (p 2 - 2.3.1)
Allocation concealment (selection bias)	Low risk	"Allocation concealment was ensured as the interviewers, responsible for enrolling participants onto the trial, were blinded to the randomized study design while one of the authors (MG) was responsible for randomisation. Hence, it was not possible for either participants or interviewers to be aware of the randomisation sequence." (p 2 - 2.3.1)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Participants and interviewers were blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	At post-test, it was not possible to blind the interviewers but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Minimal attrition that is consistent across groups (figure 1)
Selective reporting (reporting bias)	Unclear risk	"[S]uccess of study protocol" limitation to protocol: men not confronted with actual

**Gattellari 2005** (Continued)

		decision to undergo PSA screening; no indication that trial registered in central trials registry (p 13, paragraph 5)
Other bias	Low risk	“[H]igh follow-up rate and allocation concealment; study not subjected to selection bias” (p 13, paragraph 5). Appears to be free of other sources of bias

**Green 2001**

Methods	Randomized to decision aid + counselling vs counselling alone vs usual care
Participants	29 + 14 women with a first degree relative with breast cancer interested in learning about genetic testing in the USA
Interventions	DA: CD-ROM plus counselling on options’ outcomes, clinical problem, others’ opinions, guidance/coaching Comparator: counselling Comparator: usual care
Outcomes	Primary outcome: preferred options Secondary outcome: knowledge
Notes	-

<i><b>Risk of bias</b></i>		<i><b>Risk of bias</b></i>
<b>Bias</b>	<b>Authors’ judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	“[B]lock randomization schedule to one of three groups in a 2:2:1 ratio” (p 2)
Allocation concealment (selection bias)	Unclear risk	No information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	“[G]enetic counsellor blinded to randomization until just prior to the session” (p 2), unclear if participants were blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	“Values do not always add up to the number of participants due to missing data”; reasons not mentioned (p 4). “Participants’ baseline knowledge was reflected in the control group’s answers”; participants bal-

**Green 2001** (Continued)

		anced in study groups
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Low risk	Appears to be free of other sources of bias

**Hamann 2006**

Methods	Cluster-randomized trial of decision aid vs usual care
Participants	54 + 59 patients with schizophrenia considering treatment options (cluster-RCT with 12 wards paired and randomized) in Germany
Interventions	DA: 16-page booklet on options' outcomes, outcome probabilities, explicit values clarification, coaching/guidance Comparator: usual care
Outcomes	Knowledge, participation in decision making (COMRADE - doctor gave me a chance to decided which treatment I thought was best for me), uptake of psycho-education, rehospitalization, adherence, satisfaction with care, severity of illness (baseline only), attitudes about drug use, decision making preference
Notes	Primary outcome was not specified

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"[O]ne member of each pair being randomly assigned to the control or to the interventional condition" (p 266). Sequence generation method was not stated
Allocation concealment (selection bias)	Unclear risk	No mention of allocation concealment
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Reasons for attrition mentioned
Selective reporting (reporting bias)	Unclear risk	No information provided



**Hamann 2006** (Continued)

Other bias	High risk	Clustering was not accounted for in the analysis
------------	-----------	--------------------------------------------------

**Hanson 2011**

Methods	Randomized to decision aid vs usual care
Participants	127 + 129 patients diagnosed with advanced dementia and eating problems considering long-term feeding tube placement in the USA
Interventions	DA: booklet or audio recording on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, others' opinion, guidance (steps in decision making, worksheet, summary) Comparator: usual care
Outcomes	Primary outcomes: decisional conflict (3 months post-DA) Secondary outcomes: surrogate knowledge, risk perceptions, frequency of communication with providers (3 months post-DA), feeding treatment use (3, 6 and 9 months post-DA), participation in decision making, satisfaction with the decision, decisional regret
Notes	-

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computerized random number generation (p 2010, Randomization)
Allocation concealment (selection bias)	Unclear risk	No description of method used to conceal allocation (p 2010, Randomization)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"Cluster randomization prevented double blinding and may have introduced bias due to site effects" (p 2014, Discussion); study authors unsure of effect on study
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"[B]ecause of cluster randomization, data collectors were not blinded to group assignment" (p 2010, Randomization); authors believe has little impact on study
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Intervention group missing data for 1 participant, reason for omission not reported (table 1) No explanation for number of participants in each group (n = 127) given numbers vary

**Hanson 2011** (Continued)

		from those in 'recruitment and retention' figure (table 4)
Selective reporting (reporting bias)	Low risk	Registered with clinicaltrials.gov, protocol on website
Other bias	Low risk	Appears to be free of other potential biases

**Heller 2008**

Methods	Randomized to decision aid vs usual care
Participants	66 + 67 breast cancer patients eligible for breast reconstruction in the USA
Interventions	DA: interactive software programme on options' outcomes, others' opinions Comparator: standard patient education
Outcomes	Knowledge, anxiety, satisfaction with treatment choice, satisfaction with decision-making ability
Notes	Primary outcome was not specified

<i>Risk of bias</i>		<i>Risk of bias</i>
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	"upon study entry, the participants were randomized (computer generated) to one of two groups" (p 2)
Allocation concealment (selection bias)	Unclear risk	Not enough information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No information provided
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No information provided
Incomplete outcome data (attrition bias) All outcomes	Low risk	Baseline anxiety and knowledge included in graphs. Participant numbers between study groups balanced (p 3). Reasons for incomplete questionnaires and study withdrawals mentioned
Selective reporting (reporting bias)	Unclear risk	No information provided re protocol

**Heller 2008** (Continued)

Other bias	Low risk	Appears to be free of other potential biases
------------	----------	----------------------------------------------

**Hess 2012**

Methods	Randomized to decision aid vs usual care
Participants	103 + 105 patients in the the emergency department with primary symptoms of nontraumatic chest pain and were being considered of admission to the emergency department observation unit for monitoring and cardiac stress testing within 24 hours
Interventions	DA (in consultation): 1-page printout on options' outcomes, clinical problem, and outcome probabilities Comparator: usual care
Outcomes	Primary outcomes: knowledge Secondary outcomes: risk perceptions, decisional conflict, actual choice, satisfaction with decision making process, patient-practitioner communication
Notes	-

<i><b>Risk of bias</b></i>		<i><b>Risk of bias</b></i>
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	"Patients were randomized to either usual care or shared decision making through a Web-based, computer-generated allocation sequence in a 1:1 concealed fashion" (p 253)
Allocation concealment (selection bias)	Low risk	"Patients were randomized to either usual care or shared decision making through a Web-based, computer-generated allocation sequence in a 1:1 concealed fashion" (p 253)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Personnel were blinded, but unclear if patients were blinded (p 253, Outcome measures). However, the primary outcome is unlikely to be biased
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Investigators assessing outcomes were blinded (p 253, Outcome measures).
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Some of the numbers of patients reported in the results did not match the flow chart
Selective reporting (reporting bias)	Low risk	Protocol is available

Other bias	Low risk	Appears to be free of other biases
------------	----------	------------------------------------

**Jibaja-Weiss 2011**

Methods	Randomized to decision aid vs usual care
Participants	51 + 49 women diagnosed with breast cancer considering surgical treatment in the USA
Interventions	DA: computer programme on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, others' opinion and guidance (step-by-step process for making the decision) Comparator: usual care + breast cancer treatment educational materials normally provided to patients
Outcomes	Surgical treatment preference (post-DA), breast cancer knowledge (pre, post-DA, post-DA and consult), satisfaction with surgical decision (post-DA), satisfaction with decision-making process (post-DA), decisional conflict (pre, post-DA, post-DA and consult), proportion undecided
Notes	Primary outcome was not specified

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Patients at each hospital were randomized using permuted blocks" (p 42, Methods section)
Allocation concealment (selection bias)	Unclear risk	Not addressed in the study
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not addressed in the study
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	There is no way to know if the plots include all of the participants' data since they do not specify what was the number of patients used to obtain these mean scores
Selective reporting (reporting bias)	Unclear risk	No mention of protocol
Other bias	Low risk	Appears to be free of other potential biases

## Johnson 2006

Methods	Randomized to decision aid vs usual care
Participants	32 + 35 patients considering endodontic treatment options in the USA
Interventions	DA (in consultation): decision board on options' outcomes, clinical problem, outcome probability, guidance Comparator: usual care
Outcomes	Primary outcomes: knowledge, satisfaction with decision making process, anxiety
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"[F]our computerized random generation lists to assign to one of two groups" (p 3)
Allocation concealment (selection bias)	Unclear risk	Not for residents: computer-generated randomization lists (1 for each resident) were prepared by the PI (p 3-4); therefore residents would have had pre-generated lists; Unclear for patients: "allocation was concealed from patients" (p 3) but does not explain how
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Blinding not mentioned. Allocation was concealed from patients only (p 3)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Flow diagram (p 6); all 40 patients agreed to participate in the study, but only 32 questionnaires were useable several residents did not understand need for entering data on the envelope and placing matched questionnaire in it (p 5)
Selective reporting (reporting bias)	Unclear risk	No indication that the trial was registered in a central trials registry
Other bias	Unclear risk	"[B]aseline data obtained because possible that clinicians training in the EndoDB would alter usual care discussions" (p 5) . Mentions taking baseline characteristics,

**Johnson 2006** (Continued)

	but not included in article
--	-----------------------------

**Kasper 2008**

Methods	Randomized to decision aid vs usual care
Participants	150 + 147 multiple sclerosis patients considering immunotherapy in Germany
Interventions	DA: booklet and worksheet on options' outcomes, clinical problem, outcome probabilities, explicit values clarification (based on IPDAS) Comparator: information material on immunotherapy (80 pages)
Outcomes	Primary outcomes: role in decision making Secondary outcomes: choice, feeling undecided, helpfulness with making a decision, attitudes toward immunotherapy, expectations of side effects realized at 6 months
Notes	-

<i>Risk of bias</i>			<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement	
Random sequence generation (selection bias)	Low risk	"[A]llocation using computer generated random numbers" (p 5)	
Allocation concealment (selection bias)	Unclear risk	Randomization was carried out by concealed allocation, but method of concealment was not described (p 2, Assignment)	
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Participants were not told whether the information they received was standard information or the newly developed DA (p 3, Masking)	
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Assessors were not told whether the information they received was standard information or the newly developed DA (p 3, Masking)	
Incomplete outcome data (attrition bias) All outcomes	Low risk	Flow of participants (p 2, Fig 1); baseline data/characteristics included	
Selective reporting (reporting bias)	Low risk	"The protocol of this study has been published with the trial registration at <a href="http://controlled-trials.com/ISRCTN25267500">http://controlled-trials.com/ISRCTN25267500</a> " (p 2)	

**Kasper 2008** (Continued)

Other bias	Unclear risk	Difference in preferred interaction style between groups at baseline (P value 0.04) (p 5)
------------	--------------	-------------------------------------------------------------------------------------------

**Kennedy 2002**

Methods	Randomized to decision aid + coaching vs decision aid only vs usual care
Participants	215 + 206 + 204 women considering treatment for menorrhagia in the UK
Interventions	DA: video + booklet on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, others' opinions, guidance/coaching Coaching: ~ 20 minute coaching with explicit values clarification by a registered nurse prior to seeing physician Comparator: usual care
Outcomes	Primary outcomes: general quality of life Secondary outcomes: uptake of option, satisfaction, menorrhagia severity, cost-effectiveness
Notes	-

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Allocation sequence was generated by computer and stratified by consultant and the age at which the woman left full-time education (p 3)
Allocation concealment (selection bias)	Low risk	"Secure randomization ensured by using a central telephone randomization system" (p 3)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Possibility of contamination bias; clinicians could have applied the experience gained from consultations with the interventions groups in their consultations with the control group (p 6)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear if blinding used but most outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Table 1 and Figure 1 flow diagram (p 4-5)

**Kennedy 2002** (Continued)

Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Low risk	Appears to be free from other risks of bias

**Knops 2014**

Methods	Randomized to decision aid vs usual care
Participants	91 + 87 patients with asymptomatic abdominal aortic aneurysm considering elective surgery vs watchful waiting
Interventions	DA: interactive CD-ROM on options' outcomes, clinical problem, outcome probabilities, explicit values clarification Comparator: usual care with regular information
Outcomes	Primary outcomes: decisional conflict (baseline, 1, 4, and 10 months) Secondary outcomes: patient knowledge (baseline and 1 month), anxiety (baseline, 1, 4, and 10 months), satisfaction with conversation with the surgeon (baseline and 1 month), final treatment choice (10 months), aneurysm rupture (10 months), possible date of surgery (10 months), postoperative morbidity and mortality (10 months), physical quality of life (baseline, 1, 4, and 10 months)
Notes	Trial registration: NTR1524

**Risk of bias**

**Risk of bias**

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Computer-generated randomisation ALEA v.2.2, NKI-AVL, the Netherlands) was performed by the investigators" (p 2)
Allocation concealment (selection bias)	Low risk	"Computer-generated randomisation ALEA v.2.2, NKI-AVL, the Netherlands) was performed by the investigators" (p 2)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Patients and investigators could not be blinded after group assignment, a factor which is inherent to the decision aid and the design of the study. Surgeons and nurses involved in the outpatient care of the participants were blinded to the patient's allocation group, although patients were not prohibited from sharing their allocation with them." (p 3)



**Knops 2014** (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcome measurement is not likely to be influenced by lack of blinding as all outcomes were measured objectively using validated scales and data retrieved from medial records
Incomplete outcome data (attrition bias) All outcomes	Low risk	Appears to have similar attrition between groups. The proportion of values missing varied from 2% to 9% per outcome measure. Missing values were completed by multiple imputation analysis. If one of the outcome measures had more than 25% missing values, that outcome measure for that patient was excluded from analysis. Therefore, missing data have been handled appropriately (p 3)
Selective reporting (reporting bias)	Unclear risk	Insufficient information to make judgment
Other bias	High risk	<p>“Considerable number of patients could not be included, were not asked to participate, or declined to participate. Selection bias may have occurred in patients that were not included” (p 6)</p> <p>“Both patients and surgeons were aware of the aim and subject of the study and could not be blinded to the allocation. It is possible that surgeons in the contributing centres offered more than average information to their patients” (p 6). Performance bias may have been introduced in terms of altered communication style</p>

**Krist 2007**

Methods	Randomized to decision aid booklet vs decision aid web-based vs usual care
Participants	196 + 226 + 75 patients considering prostate cancer screening in the USA
Interventions	DA: 4 page pamphlet with options' outcomes, clinical problem, outcome probability Comparator: web-site with same information as paper based DA Comparator: usual care
Outcomes	Primary outcomes: role in decision making Secondary outcomes: knowledge, decisional conflict, time spent discussing screening, choice (PSA test ordered)
Notes	-

**Krist 2007** (Continued)

<i>Risk of bias</i>			<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement	
Random sequence generation (selection bias)	Low risk	"[C]oordinator referred to pre-generated randomisation tables to inform the participant to which arm he was randomised" (p 2)	
Allocation concealment (selection bias)	Low risk	At the time of enrolment, the allocation was concealed from the coordinator (p 2)	
Blinding of participants and personnel (performance bias) All outcomes	High risk	Physicians were not blinded - could affect decision making process and uptake of screening	
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation	
Incomplete outcome data (attrition bias) All outcomes	Low risk	p 3, Results; p 4, Flow diagram	
Selective reporting (reporting bias)	Unclear risk	No information provided	
Other bias	Unclear risk	Uneven groups but done intentionally, ration of 1:3:3 but appears to be free of other potential biases	

**Kupke 2013**

Methods	Cluster-randomized trial of 2 groups of dental students to decision board group and non-decision board group. Patients randomized to students in either group
Participants	57 + 36 patients with defect in posterior tooth (Class II defect) considering 6 treatment options, including no therapy
Interventions	DA (in consultation): options' outcomes, outcome probabilities Comparator: usual care with discussion of the treatment options
Outcomes	Knowledge (costs/self-payment, survival rate, characteristics and treatment time) (postintervention); overall satisfaction with consultation (postintervention)
Notes	Primary outcome not specified
<i>Risk of bias</i>	

**Kupke 2013** (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomly assigned by a dice (selection of students and patient allocation) (p 20)
Allocation concealment (selection bias)	High risk	"The patients were assigned to the students according to common standards of the university independently and without knowing which group the student belonged to." (p 20)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Patients were assigned to the students independently and without knowing which group the students belonged to" (p 20)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to judge if blinding of outcome assessment occurred
Incomplete outcome data (attrition bias) All outcomes	Low risk	Similar attribution in both groups; "missing answers were treated as incorrect answers, while illegible answers were treated as missing values" (p 22)
Selective reporting (reporting bias)	Unclear risk	No mention of study protocol or trial registration. No way to ensure the outcomes they intended to measure are fully reported
Other bias	High risk	Did not adjust for clustering in analysis

**Kuppermann 2014**

Methods	Randomized to decision aid vs usual care
Participants	375 + 369 11-week pregnant women who had not yet undergone prenatal screening or diagnostic testing
Interventions	DA: describes clinical condition, options, outcome probabilities, values clarification Comparator: usual care
Outcomes	Primary outcomes: invasive prenatal diagnostic testing (3 to 6 months) Secondary outcomes: testing strategy undergone (3 to 6 months), knowledge (3 to 6 months), accurate risk perception (procedure related miscarriage, DS affected fetus) (3 to 6 months), decisional conflict (3 to 6 months), decisional regret (3 to 6 months)
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"A computer generated random allocation sequence assigned participants to experimental groups within permuted blocks of random size, with a 1:1 allocation ratio, stratified by age, clinical site, parity, and interviewer" (p 1211)
Allocation concealment (selection bias)	Low risk	"The randomization code was not available to any study-related personnel until data analysis was complete" (p 1211)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Different research associates facilitated baseline and follow-up interviews and medical record review to ensure blinding to the randomization assignment" (p 1211)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Different research associates facilitated baseline and follow-up interviews and medical record review to ensure blinding to the randomization assignment" (p 1211)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Similar attrition in both groups. "[A]ll reported analyses were based on a modified intention-to-treat sample" (p 1211)
Selective reporting (reporting bias)	Low risk	Trial registered
Other bias	Low risk	Appears to be free of other sources of bias

**Lam 2013**

Methods	Randomized to decision aid or standard information booklet after initial consultation
Participants	138 + 138 women considering breast cancer surgery for early-stage breast cancer
Interventions	DA: take-home booklet on clinical problem, options' outcomes, outcome probabilities, guidance, explicit values clarification Comparator: standard information booklet
Outcomes	Primary outcomes: treatment decision making difficulties and decisional conflict scale at 1 week post consultation, knowledge at 1-week postconsultation, decision regret at 1 month after surgery Secondary outcomes: postoperative psychological distress (anxiety and depression) at

	1, 4, and 10 months after surgery, decision regret at 4 and 10 months after surgery, treatment decision
--	---------------------------------------------------------------------------------------------------------

Notes	-
-------	---

<b><i>Risk of bias</i></b>	<b><i>Risk of bias</i></b>
----------------------------	----------------------------

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Patient assignment to treatment and control arms was performed using a prior computer-generated random-number sequence" (p 2880)
Allocation concealment (selection bias)	Low risk	"A serially labeled, opaque, sealed-envelope method was used for block randomization" (p 2880)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Two research staff members - one responsible for preintervention assessment and block allocation and the other for postintervention assessments - ensured that the researcher performing follow-up assessments was blinded regarding women's allocation status." "Blinding surgeons to allocation status proved impractical." (p 2880)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	1 research staff member was responsible for postintervention assessments to ensure that the researcher performing follow-up assessments was blinded regarding women's allocation status (p 2880)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Does not appear to be missing any outcome data; similar attrition in both groups
Selective reporting (reporting bias)	Low risk	Study protocol available online with published study
Other bias	Low risk	Does not appear to be subject to other sources of bias

## Langston 2010

Methods	Randomized to decision aid + coaching vs usual care
Participants	114 + 108 women pregnant women in their first trimester considering use of contraceptives in the USA
Interventions	DA: double-sided flip chart on clinical problem, outcome probabilities, guidance (administered by a research assistant), coaching (structured, standardized, non-directive contraceptive counselling) + usual care Comparator: usual care
Outcomes	Primary outcomes: proportion of participants choosing very effective contraceptive method (post-DA and consult) Secondary outcomes: actual choice on day of procedure (post-DA and consult), adherence of very effective and/or effective methods at 3 months and at 6 months (post-DA and consult)
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Using a random-number table, we determined the sequence for 1:1 allocation constrained by blocks of 10" (p 363, Methods-study procedures)
Allocation concealment (selection bias)	Low risk	"Randomization assignments were sealed inside numbered, opaque envelopes" (p 363, Methods-study procedures)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"No blinding of participants or coordinators was feasible due to the nature of the intervention. Physician-providers did not know the participant's allocation group, did not discuss the study with patients, and were asked not to change their counselling" (p 363, Methods-study procedures)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	For "method initiation on the day of the procedure" it is only said that the "[p]articipants in the intervention group were not more likely to initiate the requested method immediately compared to those in the usual care group"; possible that

**Langston 2010** (Continued)

		the results contradicted the hypothesis and were excluded for this reason
Selective reporting (reporting bias)	Unclear risk	No mention of study protocol; not enough information to permit judgement
Other bias	Low risk	Appears to be free of other potential biases

**Laupacis 2006**

Methods	Randomized to decision aid vs usual care
Participants	60 + 60 patients undergoing elective open heart surgery considering pre-operative autologous blood donation in Canada
Interventions	DA: audiotape booklet on options' outcomes, clinical problem, outcome probability, explicit values clarification, guidance (Ottawa Decision Support Framework) Comparator: usual care
Outcomes	Primary outcomes: knowledge, decisional conflict Secondary outcomes: uptake of option, satisfaction with decision making process, satisfaction with decision, accurate risk perceptions
Notes	-

<i><b>Risk of bias</b></i>		<i><b>Risk of bias</b></i>
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	"Randomization envelopes were prepared centrally by a statistician" (p 2)
Allocation concealment (selection bias)	Low risk	"The envelopes were labeled with identification numbers and contained a card specifying the patient's group assignment. The envelopes were opened by the interviewer after completion of the baseline interview." (p 2)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation

**Laupacis 2006** (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Results, p 4; fig 1, flow diagram
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Low risk	Appears to be free of other potential biases

**LeBlanc 2015**

Methods	Randomized to decision aid vs individualized score only vs usual care
Participants	32 + 33 + 14 women over 50 years diagnosed with osteopenia or osteoporosis not taking biphosphonates or other prescription medication
Interventions	DA (in consultation): clinical problem, individualized risk of condition, options' outcomes, guidance Comparator 1: individualized risk Comparator 2: usual care
Outcomes	Primary outcomes: knowledge (immediately post), decisional conflict (immediately post), participation in decision-making process (immediately post), decision to start (immediately post), adherence (6 months), acceptability (timing not specified), satisfaction with the decision-making process (not specified), quality of life (not specified), time (review of video consultation) Secondary outcome: decision quality (not reported)
Notes	-

***Risk of bias***

***Risk of bias***

<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	"Patients were allocated using a computer-generated sequence that randomized them 1:1:1 in a concealed fashion" (p 5)
Allocation concealment (selection bias)	Low risk	"Patients were allocated using a computer-generated sequence that randomized them 1:1:1 in a concealed fashion" (p 5)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Patients and clinicians were aware of the overall objective, presented as improvement in communication between patients and clinicians during the clinical encounter, but remained blinded to the specific aims" (p 5)



**LeBlanc 2015** (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	"After randomization, only data analysts remained blind to allocation" (p 5)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Used intention-to-treat analysis; similar attrition in both groups
Selective reporting (reporting bias)	Unclear risk	Trial registered; Checklists available for CONSORT and protocol. Sample size originally calculated based on adherence but re-calculated for decisional conflict given inability to reach original target
Other bias	High risk	"Possible contamination at the clinician level (i.e. clinician who, having used the decision aid with a prior patient, recreates elements of the decision aid with a subsequent patient allocated to receive FRAX alone or usual care) was monitored by a detailed review of the available video recorded encounters" (p 5)

**Legare 2008a**

Methods	Randomized to decision aid vs usual care
Participants	45 + 45 women considering use of natural health products for managing menopausal symptoms
Interventions	DA: booklet with worksheet on options' outcomes, clinical problem, explicit values clarification, guidance/coaching (Ottawa Decision Support Framework) Comparator: general information brochure on the clinical problem (did not address risks and benefits)
Outcomes	Primary outcomes: decisional conflict Secondary outcomes: knowledge of natural health products in general (not specific option outcomes), preferred choice, values-choice agreement, proportion undecided
Notes	-

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The randomization scheme was carried out by a biostatistician using computer-generated unequal blocks

**Legare 2008a** (Continued)

Allocation concealment (selection bias)	Low risk	Sealed opaque envelopes containing 1 or the other documents (a PDA in the intervention group and a general information brochure in the control group) were prepared by another individual, external to the study
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	The investigators were blinded but no mention of blinding of participants
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	See Figure 1 for flow diagram, reason for loss to follow-up was described
Selective reporting (reporting bias)	Low risk	Trial registration identifier is NCT00325923
Other bias	Low risk	No statistically significant difference in women's characteristics between groups (Table 1)

**Legare 2011**

Methods	Cluster-randomized to decision aid vs usual care
Participants	245 + 214 patients with non-emergent acute respiratory infections considering using antibiotics in Canada
Interventions	DA (in consultation): pamphlet on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, guidance and coaching Comparator: delayed intervention
Outcomes	Primary outcomes: <ul style="list-style-type: none"> <li>• Patient outcomes: actual choice (pre and post-DA), perceived decision quality (pre and post-DA), decisional conflict (pre and post-DA), decision regret (pre and post-DA), general health outcomes</li> <li>• Practitioner outcomes: decision, perceived decision quality, decisional conflict</li> </ul> Secondary outcomes: <ul style="list-style-type: none"> <li>• Patient outcomes: intention to engage in future SDM (pre and post-DA), participation in decision making</li> <li>• Practitioner outcomes: intention to engage in future SDM and comply with clinical practice guidelines</li> </ul>
Notes	-

**Legare 2011** (Continued)

<i>Risk of bias</i>		<i>Risk of bias</i>
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	"A biostatistician simultaneously randomised all FMGs and allocated them to groups using Internet-based software" (p 99)
Allocation concealment (selection bias)	Low risk	"Using Internet-based software" (p 99)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding of participants and personnel: only biostatistician was blinded (p 99)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Biostatistician who assesses the outcomes is blinded, outcomes were objectively measured (p 99)
Incomplete outcome data (attrition bias) All outcomes	Low risk	There appear to be no missing data
Selective reporting (reporting bias)	Low risk	No missing pre-specified outcomes
Other bias	Low risk	Appears to be free of other sources of bias

**Legare 2012**

Methods	Cluster-randomized controlled trial to decision aid vs usual care
Participants	239+210 adults and children with with a diagnosis of acute respiratory infection (e.g., bronchitis, otitis media, pharyngitis, rhinosinusitis)
Interventions	DA (in consultation): pamphlet on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, guidance and coaching (participating physicians also received training in the form of a 2-hour online tutorial and a 2-hour on-site interactive workshop) Comparator: usual care
Outcomes	Primary outcome: use of antibiotics (immediately post consultation) Secondary outcomes: decisional conflict (immediately post), control preference scale (immediately post), quality of decision (immediately post), adherence to the decision (2 weeks post), repeat consultation (2 weeks post), decisional regret (2 weeks post), quality of life (2 weeks post) and intention to engage in SDM in future consultations regarding antibiotics for acute respiratory infections (2 weeks post)
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"A biostatistician used internet-based software to simultaneously randomize all 12 family practice teaching units to either the intervention group or control group. The teaching units were stratified according to rural or urban location" (p E728)
Allocation concealment (selection bias)	Low risk	"A biostatistician used internet-based software to simultaneously randomize all 12 family practice teaching units to either the intervention group or control group. The teaching units were stratified according to rural or urban location" (p E728)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Patients with symptoms suggestive of an acute respiratory infection were initially recruited by a RA in the waiting room before consultation with a physician" (p E728)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"The biostatistician was unaware of group allocation, the researchers and research assistants who recruited patients and collected data were not" and "Statistical analysis was performed by a statistician who was unaware of the teaching unit allocations" (p E729)
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	Protocol registered and published
Other bias	Low risk	"To avoid contamination bias, access to the online tutorial was denied to providers in the control group during the trial" (p E728)

## Leighl 2011

Methods	Randomized to DA + usual care vs usual care
Participants	107 + 100 patients diagnosed with metastatic CRC considering advanced chemotherapy in Australia and Canada

**Leighl 2011** (Continued)

Interventions	DA: booklet and audiotape on option' outcomes, clinical problem, outcome probabilities, explicit values clarification and guidance (steps in decision making + worksheet) Comparator: usual care
Outcomes	Primary outcomes: knowledge (post-DA), satisfaction with decision (post-DA) Secondary outcomes: anxiety (pre and post-DA), satisfaction with consultation (post-DA), choice leaning (post-DA), decisional conflict (post-DA). achievement of their information preference (post-DA), participation in decision making (post-DA), acceptability (post-DA), quality of life (post-DA)
Notes	-

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer generated randomized lists (p 2078, Study design)
Allocation concealment (selection bias)	Low risk	Code concealed in sealed envelopes until time of random assignment (p 2078, Study design)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Patients not blinded and subjective outcomes may be affected by them knowing their assignment
Blinding of outcome assessment (detection bias) All outcomes	Low risk	All outcomes are not subjected to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	31% dropout rate, but similar losses across all groups
Selective reporting (reporting bias)	Unclear risk	Protocol not available
Other bias	Low risk	Appears to be free of other sources of bias

**Lepore 2012**

Methods	Randomized to decision support intervention (decision coaching by telephone + educational pamphlet) vs control
Participants	244 + 246 African American men aged 45-70 in the USA

Interventions	DA: condition-specific educational pamphlet on prostate cancer screening and tailored telephone education on options' outcomes, explicit values clarification, others' opinions, and guidance (decision coaching) Comparator: attention control (education on fruit and vegetable consumption)
Outcomes	Primary outcomes: knowledge (pretest and post-test at 8 months postrandomization), decisional conflict (posttest), physician visit to discuss testing (post-test), adherence as congruence between testing intentions and behaviors (post-test) Secondary outcomes: testing intention (post-test), benefit-to-risk ratio of testing (post-test), PSA screening (post-test), anxiety (pretest and post-test)
Notes	Trial registration NCT01415375

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The principal investigator used a computer-generated randomization schedule to randomize the participant." (p 322)
Allocation concealment (selection bias)	Unclear risk	"The principal investigator used a computer-generated randomization schedule to randomize the participant and emailed the randomization assignment to the interventionist." (p 322)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Interventionists were not blind to condition. We can assume that patients were blinded as the study design was a telephone call for both intervention and control groups (p 322)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Data collectors were blind to condition but the interventionists were not" (p 322)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Does not appear to be missing any outcome data
Selective reporting (reporting bias)	Low risk	Appears to have reported on all pre-specified outcomes (protocol)
Other bias	Low risk	Appears to be free of other potential sources of bias

## Lerman 1997

Methods	Randomized to decision aid vs waiting list control
Participants	122 + 114 + 164 women considering BRCA1 gene testing in the USA
Interventions	DA: education and counselling on options' outcomes, clinical problem, outcome probability, explicit values clarification, others' opinions, guidance/coaching Comparator: no intervention
Outcomes	Primary outcome: preferred option Secondary outcomes: knowledge, accurate risk perceptions, perceived personal risk/benefits/limitations, agreement between values and choice
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No information provided
Allocation concealment (selection bias)	Unclear risk	No information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Of 440 women, 400 completed 1-month follow-up interviews; no reasons provided; baseline data/characteristics included (p 2)
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Low risk	Appears to be free of other potential biases

## Lewis 2010

Methods	Cluster-randomized to decision aid vs usual care
Participants	211 + 232 patients considering colorectal cancer screening in the USA
Interventions	DA: web-based, DVD and VHS videotape formats + stage targeted brochures (and booster kit if patients had not been screened) on options' outcomes, clinical problem, outcome probabilities, others' opinion, guidance (encouraged patients to communicate

	with their practitioners by asking questions and sharing preferences; summary) Comparator: usual care using Aetna annual reminders to obtain CRC screening
Outcomes	Knowledge of the age at which screening should begin (post-DA), completion of colorectal cancer screening (pre, post-DA), intrusive thoughts (pre, post-DA), interest in CRC screening (pre, post-DA), intent to ask provider about screening (pre, post-DA), readiness to be screened (pre, post-DA), perceived risk of colon cancer (pre, post-DA), general beliefs about colon cancer (pre, post-DA), fears about colorectal cancer screening (pre, post-DA), perceptions about whether participants had enough information (post-DA), whether participants had enough information about specific screening tests (post-DA), willingness to pay for screening tests (post), desire to participate in medical decision (post) <b>Practice level measures:</b> assess CRC screening practices (pre, post-DA), referrals (pre, post-DA), quality improvement initiatives
Notes	Primary outcome was not specified

***Risk of bias******Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Randomisation was done using matched pairs and a blocking procedure." (p 2, Practice recruitment and randomization section)
Allocation concealment (selection bias)	Unclear risk	"Thus, purposive assignment to treatment group was used, resulting in a hybrid randomisation" (p 3, Practice recruitment and randomization section). There is no mention of the effect of this purposive assignment on the study
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	As mentioned above, staff used purposive assignment and were therefore not blinded, but there is no mention of the effect on the study
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The study did not address this outcome, but outcomes were objectively measured
Incomplete outcome data (attrition bias) All outcomes	Low risk	There appear to be no missing outcome data
Selective reporting (reporting bias)	Unclear risk	No mention of study protocol
Other bias	High risk	Unadjusted cluster analysis



## Loh 2007

Methods	Cluster-randomized to decision aid vs usual care
Participants	263 + 142 patients with physician diagnosed depression (cluster RCT with 30 general practitioners randomized) in Germany
Interventions	DA (in consultation): options' outcomes, clinical problem, explicit values clarification, guidance/coaching Comparator: usual care
Outcomes	Participation in decision making, adherence, satisfaction with clinical care, depression severity, consultation length
Notes	Primary outcome was not specified

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"[T]wo-thirds of the general practitioners were randomly assigned to the intervention group by drawing blinded lots under the supervision of the principal investigator and two researchers" (p 3)
Allocation concealment (selection bias)	Low risk	Drawing blinded lots (p 3 - 2.1)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Unclear blinding, not enough information provided to assess whether this contributes to bias on outcomes not measured by using a scale (e.g. consultation time was documented in minutes by the physicians following each consultation)
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	"Further results resting on the baseline phase of this trial were already presented elsewhere" (p 5, fig); "unequal distribution of physicians was due to possibility of higher dropout rate in intervention group because of additional time and effort" (p 3)
Selective reporting (reporting bias)	Unclear risk	No indication that the trial was registered in a central trials registry

**Loh 2007** (Continued)

Other bias	Low risk	Appears to be free of other potential biases (p 5-6, details pt and physician baseline characteristics). Statistically significant differences were controlled for in outcome analyses
------------	----------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

**Man-Son-Hing 1999**

Methods	Randomized to decision aid vs usual care
Participants	139 + 148 patients on atrial fibrillation trial considering continuing on aspirin vs change to Warfarin in Canada
Interventions	DA: audiotape booklet on options' outcomes, clinical problem, outcome probability, explicit values clarification, others' opinions, guidance (Ottawa Decision Support Framework) Comparator: usual care
Outcomes	Primary outcomes: uptake of options, adherence Secondary outcomes: help with making a decision, knowledge, accurate risk perceptions, decisional conflict, satisfaction with decision making process, role in decision making
Notes	-

<i><b>Risk of bias</b></i>		<i><b>Risk of bias</b></i>
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Computer-generated scheme (p 2)
Allocation concealment (selection bias)	Low risk	Administered from a central location (p 2)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Unclear blinding however, "contamination, physicians may have provided DA information to patients receiving usual care" (p 7)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	P 4, fig 2 flow chart. Reasons for attrition not mentioned. Baseline data not included
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Low risk	No other potential risks of bias

## Mann D 2010

Methods	Randomized to decision aid vs usual care
Participants	80 + 70 participants diagnosed with diabetes considering the use of statins to reduce coronary risk
Interventions	DA (in consultation): healthcare provider led discussion using developed tool (Statin Choice) on options' outcomes, outcome probabilities, guidance (step-by-step process for making the decision; administered by the physician in the consultation) Comparator: usual primary care visit + pamphlet
Outcomes	Knowledge (postconsult and post-DA), decisional conflict (postconsult and post-DA), risk estimation (postconsult and post-DA), beliefs (postconsult and post-DA), adherence (3 and 6 months postconsult and post-DA)
Notes	Primary outcome was not specified

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Participants were randomized but there is no mention of method used (p 138, Methods section)
Allocation concealment (selection bias)	Unclear risk	Not reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not reported
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Baseline data was provided
Selective reporting (reporting bias)	Unclear risk	Only reports on improvement (i.e. decisional conflict scale); does not present outcome data to fullest (no numerical data on knowledge results between groups, only describes in words)
Other bias	Unclear risk	"We did not adjust the clustering of effects given that few participants received care by the same clinicians" (p 139, Analysis section). No mention of magnitude in change of data due to this choice

## Mann E 2010

Methods	Randomized to decision aid vs usual care
Participants	278 + 139 participants considering diabetes screening in the UK
Interventions	DA: screening invitation on clinical problem, outcome probabilities and explicit values clarification Comparator: usual care using screening invitation on clinical problem
Outcomes	Primary outcomes: preferred option (post-DA) Secondary outcomes: whether invitation type impacts on intention (post-DA), impact on knowledge (post-DA), impact on attitude (post-DA), risk perception
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"Invitation taken from the top of a randomly ordered pile (either standard or one of two versions of an informed decision choice invitation). The materials were ordered in a way that the invitation type was hidden until the recruitment process was completed" (p 2-3, Methods, Participants section). Unclear how invitation type was hidden
Allocation concealment (selection bias)	Low risk	"Invitation taken from the top of a randomly ordered pile; materials were ordered in a way that the invitation type was hidden until the recruitment process was completed" (p 2-3, Methods, Participants section)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Interviewers were not aware of the direction of anticipated effect of materials, and materials were dummy-coded so that no sense of intervention or control would have been communicated to interviewers or participants (p 3, Methods, Participants section)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Study did not address this outcome, but outcomes were objectively measured and not subject to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data

**Mann E 2010** (Continued)

Selective reporting (reporting bias)	Unclear risk	No mention of protocol; insufficient information to permit judgment
Other bias	Unclear risk	“Present sample was ... not necessarily representative of the highest risk individuals in this age group”; “£5 incentive might have also added a selection bias”; “Lack of anonymity with verbally delivered questionnaire might encourage socially desirable responding” (p 6, Discussion section)

**Marteau 2010**

Methods	Randomized to decision aid vs usual care
Participants	633 + 639 patients considering diabetes screening in England
Interventions	DA: screening invitation on clinical problem, outcome probabilities and explicit values clarification Comparator: usual care using screening invitation on clinical problem
Outcomes	Primary outcome: attendance for screening (post-DA and consult) Secondary outcomes: intention to make changes to lifestyle (post-DA and consult), satisfaction with decisions made among attenders (post-DA and consult)
Notes	-

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	“[G]enerated simultaneously in a batch by random numbers using Excel spreadsheet software, stratifying by number of participants in household” (p 2, Randomization section)
Allocation concealment (selection bias)	Low risk	“Randomisation ... was undertaken by the study statistician from a central site” (p 2, Randomization section)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Personnel were blinded and appears that patients were unaware which arm they were in (members of the same household received the same intervention) (p 2, Randomization section)

**Marteau 2010** (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Clinical and trial staff taking measurements and entering data were unaware of the study arm to which participants had been assigned (p 2, Randomization section)
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	Published protocol (p 2, Methods)
Other bias	Low risk	Appears free of other potential biases

**Mathers 2012**

Methods	Cluster-randomized controlled trial of 49 general practices in the UK to decision aid, healthcare professional training workshop and use of PDA in consultation, or usual care	
Participants	95 + 80 participants with type 2 diabetes considering adding or changing to insulin therapy	
Interventions	DA: booklet about clinical problem, treatment options, options' outcomes, outcome probabilities, explicit values clarification, structured guidance Comparator: usual care	
Outcomes	Primary outcomes: decisional conflict (immediately postintervention), glycaemic control (glycosolated haemoglobin, HbA1c) at 6 months Secondary outcomes: knowledge (immediately post), realistic expectations (immediately post), preference option (immediately post), proportion undecided (immediately post), participation in decision-making (immediately post), regret (6 months), adherence with chosen option (6 months)	
Notes	Trial registration: ISRCTN14842077	

<i><b>Risk of bias</b></i>		<i><b>Risk of bias</b></i>
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	"All eligible and willing practices were randomly allocated by a computer" (p 3)
Allocation concealment (selection bias)	Low risk	"A statistician generated the random allocation sequence while a secretary who was not involved in the research study assigned participants to either the intervention or control groups" (p 3)

**Mathers 2012** (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	“Blinding of the intervention and assessment of the process measures were not feasible in view of the nature of the intervention studied” (p 3)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	“Blinding of the intervention and assessment of the process measures were not feasible in view of the nature of the intervention studied” (p 3)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Does not appear to be missing any outcome data
Selective reporting (reporting bias)	Low risk	Trial registered
Other bias	Unclear risk	Cannot make a judgment with information provided regarding cessation of recruitment at 175 (yet 320 required to allow detection of 0.5% difference in HbA1c)

**Mathieu 2007**

Methods	Randomized to decision aid versus usual care
Participants	367 + 367 women aged 70 to 71 years and considering a subsequent screening mammography in Australia
Interventions	DA: booklet on options' outcomes, clinical problem, outcome probability, explicit values clarification, others' opinions, guidance with worksheet (Ottawa Decision Support Framework) Comparator: BreastScreen NSW brochure - includes information for women 70 + but no numeric information about the outcomes of screening
Outcomes	Primary outcomes: actual decision, informed choice Secondary outcomes: knowledge (includes 5 questions about risk perceptions), anxiety, decisional conflict, breast cancer worry, preference/intension, attitudes about screening, relationship between objective and perceived risk of breast cancer
Notes	-

***Risk of bias***

***Risk of bias***

<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Computer programme, which assigned allocations in accordance with a simple randomization schedule (p 2, Methods)

**Mathieu 2007** (Continued)

Allocation concealment (selection bias)	Low risk	Randomized by interview staff who accessed a previously concealed computer programme (p 2, Methods)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Interviewers [at follow-up] were blinded, outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Fig 1 flow diagram (p 2)
Selective reporting (reporting bias)	Low risk	"The trial was registered with the Australian Clinical Trials Registry and the Clinical Trials Registration System" (p 5)
Other bias	Low risk	Appears to be free of other potential biases

**Mathieu 2010**

Methods	Randomized to decision aid vs usual care
Participants	189 + 223 women considering mammography screening
Interventions	DA: Internet programme + worksheet on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, others' opinions, guidance (worksheet with questions relevant to decision making process; one or more questions that asked patients to clarify their preferences; summary) Comparator: delayed intervention
Outcomes	Primary outcomes: knowledge (post-DA), risk perception Secondary outcomes: intention (post-DA), values (post-DA), informed choice (post-DA), proportion undecided
Notes	-

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"[C]omputer generated simple randomization schedule" (p 66, Randomization and baseline questions section)



**Mathieu 2010** (Continued)

Allocation concealment (selection bias)	Unclear risk	“[R]andomization was conducted in a concealed manner” (p 66). Method of allocation concealment not stated
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not reported
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	All outcomes mentioned in Outcome measures section were reported in the results section (p 68, Table 2; information for intention as well as anxiety and acceptability can be found in text format in the secondary outcomes section on pg.67-68)
Selective reporting (reporting bias)	Unclear risk	No mention of protocol
Other bias	Low risk	Appears to be free of other potential sources of bias

**McAlister 2005**

Methods	Cluster-randomized to decision aid vs usual care
Participants	219 + 215 patients considering antithrombotic therapy for nonvalvular atrial fibrillation (cluster-RCT with 102 primary care practices randomized) in Canada
Interventions	DA: audiotape booklet on options’ outcomes, clinical problem, outcome probabilities, explicit values clarification, others’ opinions, guidance (Ottawa Decision Support Framework) Comparator: usual care
Outcomes	Primary outcomes: uptake of (appropriate) option Secondary outcomes: knowledge, decisional conflict, accurate risk perceptions
Notes	-

***Risk of bias***

***Risk of bias***

Bias	Authors’ judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	“[C]luster randomization at level of primary care practice to minimize contamination; randomization was done centrally

**McAlister 2005** (Continued)

		to preserve allocation concealment using a computer generated sequence” (p 2)
Allocation concealment (selection bias)	Low risk	Randomization was done centrally to preserve allocation concealment (p 2, Methods)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not blinded, but not sure whether the lack of blinding would affect the outcomes
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcome assessors blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	Results and Fig 1 - flow diagram (p 3)
Selective reporting (reporting bias)	Low risk	DAAFI trial protocol, including copies of the various questionnaires we employed, has been published (p 1, Methods)
Other bias	Low risk	Appears to be free of other potential biases

**McBride 2002**

Methods	Randomized to decision aid vs usual care
Participants	289 + 292 perimenopausal women considering hormone replacement therapy in the USA
Interventions	DA: options’ outcomes, clinical problem, outcome probability, values clarification, others’ opinions, guidance/coaching Comparator: delayed intervention
Outcomes	Primary outcome: accurate risk perceptions Secondary outcomes: satisfaction with decision, confidence with knowledge and making/discussing decision
Notes	-

***Risk of bias***

***Risk of bias***

<b>Bias</b>	<b>Authors’ judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	No information provided; Bastian 2002, no information provided - Study design is described elsewhere (p 4)

**McBride 2002** (Continued)

Allocation concealment (selection bias)	Unclear risk	No information provided; Bastian 2002, no information provided - Study design is described elsewhere (p 4)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Complete data are available for 520 (90%) of the women (p 2). Reasons why not mentioned (Bastian 2002, p 5, Results; p 6, Baseline characteristics/data included)
Selective reporting (reporting bias)	Unclear risk	No indication that the trial was registered in a central trials registry
Other bias	Low risk	Appears to be free of other potential biases; Bastian 2002, p 8 - Eligible participants were willing to consider HRT and this may have favoured recruitment of women with higher SES and those who had prior experience with HRT

**McCaffery 2010**

Methods	Randomized to decision aid + informed choice vs HPV testing vs repeat smear
Participants	104 + 104 + 106 women screened as HPV indeterminate considering HPV testing in Australia
Interventions	DA: pamphlet on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, others' opinion and guidance (worksheet) Comparator 1: no decision support, received immediate HPV testing Comparator 2: no decision support, received a repeat cervical smear at 6 months
Outcomes	Primary outcomes: quality of life (post-DA) Secondary outcomes: waiting time anxiety (post-DA), , perceived risk (post-DA), perceived seriousness of cancer (post-DA), worriedness (post-DA), intrusive thoughts (post-DA), satisfaction with care (post-DA), anxiety (post-DA), distress and concerns (post-DA), self-esteem (post-DA), effect on sexual behaviour (post-DA), help seeking behaviour (post-DA), knowledge (post-DA)
Notes	-

<i>Risk of bias</i>			<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement	
Random sequence generation (selection bias)	Low risk	"Participants were randomised centrally by the research team within each clinic in blocks of three" (p 2, Design)	
Allocation concealment (selection bias)	Low risk	"Participants were randomised centrally by the research team within each clinic in blocks of three" (p 2, Design)	
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Patients and staff were unblinded, but objective outcomes were used	
Blinding of outcome assessment (detection bias) All outcomes	Low risk	All outcomes are on questionnaires; not subject to interpretation	
Incomplete outcome data (attrition bias) All outcomes	Low risk	Figure 3: sensitivity analysis was done to include most of the patients	
Selective reporting (reporting bias)	Low risk	Protocol available	
Other bias	Low risk	Appears to be free of other sources of bias	

## Miller 2005

Methods	Randomized to decision aid vs usual care
Participants	279 women considering BRCA1-BRCA2 gene testing in the USA
Interventions	DA: educational intervention on options' outcomes, personal family cancer history; clinical problem, outcome probability, explicit values clarification, others' opinions, guidance/coaching Comparator: provision of general information about cancer risk
Outcomes	Preferred option, knowledge, perceived risk, satisfaction
Notes	Primary outcome was not specified

<i>Risk of bias</i>			<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement	

**Miller 2005** (Continued)

Random sequence generation (selection bias)	Low risk	“[R]andomized by the CATI system” (p 4) after self-initiated telephone contact
Allocation concealment (selection bias)	Low risk	“[C]omputerized assisted telephone interview system (CATI)” (p 4)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Blinding was not addressed
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Reasons stated for initial drop-out of study participants (p 8). Patients contacted offered reasons for dropping out. Study protocol allowed patients to be reached up to 13 times at follow-up; but still not able to be reached
Selective reporting (reporting bias)	Unclear risk	No indication that the trial was registered in a central trials registry
Other bias	Low risk	Appears to be free of other sources of bias

**Miller 2011**

Methods	Decision aid vs attention placebo
Participants	132 + 132 participants considering colon cancer screening in the USA
Interventions	DA: computer-based web programme on options’ outcomes, clinical problem, outcome probabilities, others’ opinion, guidance (encourages patient-practitioner communication, summary) Comparator: computer-based web programme on prescription drug refills and safety
Outcomes	Primary outcomes: receipt of CRC screening (post-DA) Secondary outcomes: ability to state a preference, change in readiness to receive screening (pre and post-DA), CRC test ordering (post-DA), proportion undecided
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors’ judgement	Support for judgement

**Miller 2011** (Continued)

Random sequence generation (selection bias)	Low risk	Block-randomized, stratified by literacy level (p 609, Methods)
Allocation concealment (selection bias)	Unclear risk	Study does not address this domain
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Health care providers were not notified of patients' enrolment in the study at any time (p 609, Methods) RAs that administered post-DA questionnaire were not blinded but believed to be a low risk of bias (p 613, Discussion)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"[C]linical outcome assessors were [blinded]" (p 613, Discussion)
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	Protocol on ClinicalTrials.gov
Other bias	Unclear risk	USD 10 gift card for participation could affect participant pool

**Montgomery 2003**

Methods	Randomized to decision aid + decision analysis vs decision analysis vs decision aid vs usual care
Participants	51 + 52 + 55 + 59 newly diagnosed hypertensive patients considering drug therapy for blood pressure in the UK
Interventions	DA: decision analysis plus information video and leaflet on options' outcomes, clinical problem, outcome probability, explicit values clarification Comparator: decision analysis on options' outcomes, outcome probability, explicit values clarification Comparator: video and leaflet on options' outcomes, clinical problem Comparator: usual care
Outcomes	Primary outcomes: decisional conflict Secondary outcomes: uptake of option, knowledge, anxiety
Notes	-

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Montgomery 2003 (Continued)

Random sequence generation (selection bias)	Low risk	Allocation schedule was computer-generated by an individual not involved in the study (p 2)
Allocation concealment (selection bias)	Low risk	"[A]llocation was concealed to the author in advance by the nature of the minimization procedure" (p 2)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not blinded - unclear if this would introduce bias to outcome assessed
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Flow diagram (p 5)
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Low risk	Appears to be free of other potential biases

Montgomery 2007

Methods	Randomized to decision aid with values clarification vs decision aid without values clarification vs usual care
Participants	245 + 250 + 247 women with previous caesarean section in the UK
Interventions	DA: options' outcomes, clinical problem, outcome probability, explicit values clarification Comparator: options' outcomes, clinical problem, outcome probability Comparator: usual care
Outcomes	Primary outcomes: decisional conflict Secondary outcomes: choice, anxiety, knowledge, satisfaction with decision
Notes	-

*Risk of bias*

*Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Blocked by using randomly permuted and selected blocks of sizes 6, 9, 12, and 15 generated by computer (p 2 Methods, Ran-

Montgomery 2007 (Continued)

		domization)
Allocation concealment (selection bias)	Low risk	1 member of the study team generated the randomization sequence by computer, and another member of staff with no other involvement in the trial performed the allocation (p 2 Methods, Randomization)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	See flow of women through the study
Selective reporting (reporting bias)	Low risk	Trials registry ISRCTN84367722
Other bias	Low risk	Recruited more than planned to account for lost data (p 4, Sample size); baseline characteristics were balanced

Montori 2011

Methods	Randomized to decision aid vs usual care + booklet
Participants	52 + 48 women with low bone mass or osteoporosis considering taking bisphosphonates in the USA
Interventions	DA (in consultation): worksheet on options' outcomes, clinical problem, outcome probabilities, guidance (administered by physician) Comparator: usual care + general information booklet on osteoporosis
Outcomes	Patient knowledge (post-DA), satisfaction with knowledge transfer (post-DA), decisional conflict (post-DA), patient-clinician communication (OPTION), trust with physician (during intervention), clinician's perception of decision quality (post-DA), clinician's satisfaction with knowledge transfer (post-DA), uptake (post-DA), adherence (post-DA), fidelity (post-DA), contamination (post-DA), risk perception
Notes	Primary outcome was not specified

*Risk of bias*

*Risk of bias*

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------



**Montori 2011** (Continued)

Random sequence generation (selection bias)	Low risk	“computer generated allocation” (p 551, Randomization)
Allocation concealment (selection bias)	Low risk	Patients randomized “in a concealed fashion (using a secure study website)” (p 551, Randomization)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No mention of participants being blinded to their allocation; only mention of data collectors and analysts blinding (p 551, Randomization)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	“After randomization, data collectors and data analysts were blind to allocation” (p 551, Randomization); Outcomes were not subject to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	“The protocol for this trial has been reported in full” (p 550, Design)
Other bias	Unclear risk	Appears to be free of other potential biases

**Morgan 2000**

Methods	Randomized to decision aid vs usual care
Participants	120 + 120 patients with ischaemic heart disease considering revascularization surgery in Canada
Interventions	DA: Health Dialog interactive videodisc on options’ outcomes, clinical problem, outcome probability, others’ opinions Comparator: usual care
Outcomes	Primary outcome: satisfaction with the decision making process Secondary outcomes: uptake of option, knowledge
Notes	-

***Risk of bias***

***Risk of bias***

Bias	Authors’ judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Morgan 1997, p 29: all randomization enrolment was performed by telephone at which time the participant was assigned

**Morgan 2000** (Continued)

		Morgan 2000 (primary study), p 2, Methods, Patient Population: "Only the statistician was privy to the two randomisation schedules and blocking factor used"
Allocation concealment (selection bias)	Low risk	Morgan 1997, p 29: only the statistician was privy to the two randomization schedules and blocking factor; Morgan 2000, (primary study), p 2, Methods, Patient Population: "only the statistician was privy to the two randomisation schedules and blocking factor used. All randomization enrolment was performed by telephone"
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"[D]ue to nature of trial, neither patients or investigators were blinded to the study" - may introduce bias to subjective outcomes such as satisfaction
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Morgan 1997, p 39, Patient accrual and follow-up: baseline characteristics included Morgan 2000 (primary study): 78% completed follow-up (90 of 120 in the intervention; 97 of 120 in the control). reasons for attrition were provided
Selective reporting (reporting bias)	Unclear risk	No indication that the trial was registered in a central trials registry
Other bias	Unclear risk	Morgan 1997, p 56: significant number of patients were lost to follow-up (25%); Morgan 2000 (primary study): baseline data imbalance (high school grad, income, no. of diseased arteries). Dropout group reported lower incomes, may have affected results. (discussion par. 6) "Selection bias was minimized by enrolling available consecutive patients"

**Mott 2014**

Methods	Randomized to shared decision-making process with DA versus usual care
Participants	13 +14 military veterans in USA diagnosed with PTSD and had served in Iraq or Afghanistan
Interventions	DA: booklet on clinical problem, options' outcomes, structured guidance Comparator: usual care
Outcomes	Satisfaction with SDM qualitatively (postintervention), perceived advantages and disadvantages of SDM qualitative (postintervention), treatment preferences (4 months), adherence using treatment engagement (4 months)
Notes	Not reported as registered in trials database; no primary outcome reported

<i><b>Risk of bias</b></i>		<i><b>Risk of bias</b></i>
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	"Participants were randomized to SDM or UC using a computer-generated randomization sequence" (p 146)
Allocation concealment (selection bias)	Low risk	"[R]andomization envelopes were prepared by the study statistician to ensure that study staff remained masked to randomization sequence" (p 146)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information provided to make judgment
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Study staff not blinded but because outcomes were taken from medical records. "At 4-month follow-up, study staff reviewed participants' medical records to extract information on treatment preferences and engagement. Medical-record reviews were conducted by a single rater trained in use of the dataextraction form. A second rater, masked to initial ratings, reextracted data from 20% of patients" (p 146)
Incomplete outcome data (attrition bias) All outcomes	High risk	27 participants were consented and enrolled , yet only 20 (UC = 11; SMD = 9) completed the study (p 146-147). Only 5 participants in the SDM arm completed the exit interview. No mention of missing data

**Mott 2014** (Continued)

Selective reporting (reporting bias)	Low risk	No protocol available but all expected outcomes reported on
Other bias	Low risk	Does not appear to be any other sources of bias

**Mullan 2009**

Methods	Cluster-randomized to decision aid vs usual care
Participants	48 + 37 patients with type 2 diabetes considering treatment options (cluster RCT with 40 clinicians randomized) in the USA
Interventions	DA (in consultation): decision cards with information on options, outcomes, outcome probability, explicit values clarification Compare: 12-page pamphlet on oral antihyperglycaemic medications
Outcomes	Knowledge, decisional conflict, participation in decision making, acceptability of the information, change in medication, adherence, HbA1C levels, trust in physician, OPTION to analyse audio-taped encounters
Notes	Primary outcome was not specified

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer generated
Allocation concealment (selection bias)	Low risk	Central allocation
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Patients were blinded, the clinicians were not, but each session was recorded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Reasons for attrition not included
Selective reporting (reporting bias)	Low risk	Trial registration no. at clinicaltrials.gov reported
Other bias	Low risk	Appears to be free of other sources of bias

**Murray 2001a**

Methods	Randomized to decision aid vs usual care
Participants	57 + 55 men considering treatment for benign prostatic hypertrophy in the UK
Interventions	DA: Health Dialog interactive videodisc on options, outcomes, clinical problem, outcome probability, others' opinions Comparator: usual care
Outcomes	Primary outcomes: uptake of option, prostate symptoms, costs, anxiety Secondary outcomes: decisional conflict, role in decision making, general health status, utility
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"[R]andomisation schedule, stratified according to recruitment centre, was generated by computer" (p 4)
Allocation concealment (selection bias)	Low risk	"Allocation were sealed in opaque numbered envelopes, opened by the study nurse" (p 4)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not blinded but not sure how this would introduce bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Flow diagram (p 5); baseline data/characteristics included and balanced
Selective reporting (reporting bias)	Unclear risk	No indication that the trial was registered in a central trials registry
Other bias	Low risk	Appears to be free of other sources of bias

**Murray 2001b**

Methods	Randomized to decision aid vs usual care
Participants	102 + 102 women considering hormone replacement therapy in the UK
Interventions	DA: Health Dialog interactive videodisc on options outcomes, clinical problem, outcome probability, other's opinion Comparator: usual care
Outcomes	Primary outcomes: preferred option Secondary outcomes: help with making a decision, decisional conflict, role in decision making anxiety, menopausal symptoms, costs, utility, general health status
Notes	-

<i><b>Risk of bias</b></i>		<i><b>Risk of bias</b></i>
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	"[R]andomisation schedule, stratified according to recruitment centre, was generated by computer" (p 3 Methods, Randomization)
Allocation concealment (selection bias)	Low risk	"Allocations were sealed in opaque numbered envelopes, opened by the study nurse after collection of the baseline data" (p 3 Methods, Randomization)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	See page 3 figure for Progress of patients through trial
Selective reporting (reporting bias)	Unclear risk	Protocol is not mentioned
Other bias	Low risk	Similar baseline characteristics, appears to be free of other potential biases. Educational achievement was higher in control group. Quote "Subsequent analysis showed that educational level not related to use of HRT nor was there an interaction between educational attainment and the intervention"

## Nagle 2008

Methods	Cluster-randomized to decision aid vs usual care
Participants	167 + 172 women in early pregnancy considering genetic testing (26 + 29 general physicians) (cluster RCT with 60 general practitioners randomized) in Australia
Interventions	DA: 24-page booklet and worksheet on options, benefits and risks, test limitations, outcomes; clinical problem, outcome probability, explicit values clarification, opinions of others', guidance (Ottawa Decision Support Framework) Comparator: standard pamphlet on prenatal testing
Outcomes	Primary outcomes: informed choice, decisional conflict Secondary outcomes: anxiety, depression, attitudes toward pregnancy, acceptability of the intervention, choice
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated random numbers (p 3)
Allocation concealment (selection bias)	Low risk	Computer-generated random numbers by an independent statistician; allocation concealment was achieved (p 3)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"Due to the nature of the intervention, it was not possible to blind women, GP's or researchers" (p 3); unclear if this would introduce bias to outcome assessed
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Researchers were not blinded but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Results, p 4; Fig 1 - flow diagram, p 5
Selective reporting (reporting bias)	Low risk	Trial Registration - The ADEPT trial was registered in the UK with Current Controlled Trials [ISRCTN22532458] and with the Australian Clinical Trials Registry (No: 012606000234516) (p 4)
Other bias	Low risk	Appears to be free of other potential biases (p 8); selection bias but was adjusted for in analysis

# Nassar 2007

Methods	Randomized to decision aid vs usual care
Participants	102 + 98 women diagnosed with a breech presentation from 34 weeks gestation considering external cephalic version in Australia
Interventions	DA: 24-page booklet, 30-minute audio-CD and worksheet; clinical problem, outcome probability, explicit values clarification, opinions of others', guidance (Ottawa Decision Support Framework) Comparator: usual care counselling and information on the management of breech presentation
Outcomes	Primary outcomes: knowledge, decisional conflict, anxiety, satisfaction with the decision, Secondary outcomes: preferred role in decision making, preferred choice
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"[R]andomly generated using computer and stratified by parity and center using random variable block sizes" (p 2)
Allocation concealment (selection bias)	Low risk	"[P]articipants were randomized by telephoning a remote, central location" (p 2)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Womens were not blinded - unclear if this would introduce bias to outcome assessed
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Loss to follow-up because of onset of labour or incomplete data forms (p 3). Baseline characteristics are included and equal. Minimum of 84 participants in each study group achieved; p 4 - flow diagram
Selective reporting (reporting bias)	Low risk	ISRCTN14570598
Other bias	Low risk	"Maternal characteristics and baseline measures of cognitive and affective outcomes were comparable between groups" (p 3 Results, Table 1) "Blinding clinicians and employment of a research midwife to interact with women"



## Oakley 2006

Methods	Randomized to decision aid vs usual care
Participants	16 + 17 postmenopausal women with osteoporosis considering treatment options to prevent further bone loss in the UK
Interventions	DA: audiotape booklet on options' outcomes, clinical problem, outcome probability, explicit values clarification, others' opinions, guidance (Ottawa Decision Support Framework) Comparator: usual care
Outcomes	Satisfaction with information, decisional conflict (intervention group only), improvement in adherence
Notes	Primary outcome was not specified

***Risk of bias******Risk of bias***

<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	No information provided
Allocation concealment (selection bias)	Low risk	Group allocation was done by a third party, unconnected to the study and blinded to the identity of the patients (p 1)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Unclear blinding, some outcomes were assessed by open-ended questions, do not know whether this contributes to risk of bias
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Sample characteristics not included; baseline satisfaction score included. "No evaluation was carried out to determine the reasons for non-participation" (p 2)
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Unclear risk	No baseline characteristics (p 2). Only 16 patients in intervention group and 17 in control group; small sample size

Methods	Randomized to decision aid + standard counselling vs usual care (standard counselling)
Participants	15 + 15 women considering breast cancer prevention in the USA
Interventions	DA (in consultation): interactive computer decision aid on options outcomes, outcome probability Comparator: standard counselling
Outcomes	Primary outcomes: consultation length Secondary outcomes: knowledge, decisional conflict, satisfaction with the decision, acceptability of the decision aid, physician satisfaction with the consultation
Notes	-

***Risk of bias******Risk of bias***

<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	Patients were randomized evenly between groups; no information provided about generation (p 149)
Allocation concealment (selection bias)	Unclear risk	No information provided (p 149)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Demographic data included; reasons for attrition mentioned
Selective reporting (reporting bias)	Unclear risk	No reference to study protocol
Other bias	Unclear risk	Small sample size, does not say how many physicians participated in study, mentions that there were observed changes in physician behaviour (based on doing both intervention and control)

## Partin 2004

Methods	Randomized to decision aid with others' opinions vs decision aid without others' opinions vs usual care
Participants	384 + 384 + 384 men considering PSA testing in the USA
Interventions	DA: Health Dialog video on options' outcomes, clinical problem, outcome probability, others' opinions Comparator 1: pamphlet on options' outcomes, clinical problem, outcome probability Comparator 2: usual care
Outcomes	Primary outcomes: knowledge Secondary outcomes: preferred option, help with making a decision, decisional conflict
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Using a computer-generated algorithm (p 2)
Allocation concealment (selection bias)	Unclear risk	No information provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"[P]roviders were blinded to the fact that their patients were participating in a trial" "coordinator did not have direct contact with subjects" (p 5)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"[F]ollow-up interviewers blinded, statisticians were not". Outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Flow diagram (p 2); reasons for attrition mentioned and participants balanced across study groups. Sample characteristics included
Selective reporting (reporting bias)	Unclear risk	No indication that the trial was registered in a central trials registry
Other bias	Low risk	Appears to be free of other potential biases

## Pignone 2000

Methods	Randomized to decision aid vs usual care
Participants	125 + 124 adults considering colon cancer screening in the USA
Interventions	DA: video of options' outcomes, clinical problem, others' opinion Comparator: video on car safety
Outcomes	Primary outcome: uptake of options
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"[C]omputerized random number generator" (p 2, Methods, Group assignment)
Allocation concealment (selection bias)	Low risk	"[R]andomization was performed centrally and was not balanced among centers. Assignments were placed in sealed, opaque, sequentially numbered envelopes and were distributed to the three sites" (p 2, Methods, Group assignment)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"The providers and staff were not blinded to intervention status" "3 to 6 months after, different RA blinded to participant intervention examined clinic records" (p 2) Does not mention whether patients were blinded; unclear if lack of blinding contributed to potential risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	A different research assistant who was blinded to participants' intervention status examined participants' clinic records in a standardized and validated manner to determine whether colon cancer screening tests were actually completed within 3 months of the index visit
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Because of an administrative error, 18 controls did not complete the second and third questionnaires (p 4)
Selective reporting (reporting bias)	Unclear risk	Protocol was not mentioned

**Pignone 2000** (Continued)

Other bias	Low risk	Baseline characteristics similar, appear to be no other potential sources of biases. Minimized bias from repeated measurements by administering the same questionnaires to the intervention and control participants
------------	----------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

**Protheroe 2007**

Methods	Randomized to decision aid vs usual care	
Participants	60 + 56 women considering treatment options for menorrhagia in the UK	
Interventions	DA: interactive computerized DA on options' outcomes, clinical problem, outcome probability, explicit values clarification, guidance Comparator: information leaflet	
Outcomes	Primary outcomes: decisional conflict Secondary outcomes: knowledge, anxiety, condition specific health outcomes, treatment preference, undecided	
Notes	-	

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer generated randomization, stratified by practice and minimized according to age (p 2, Methods)
Allocation concealment (selection bias)	Unclear risk	Random allocation was concealed from the individual who was making judgments of eligibility, but the method of concealment was not stated (p 2, Methods)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Fig 6 flow diagram (p 5); baseline data/characteristics included and balanced (p 4)
Selective reporting (reporting bias)	Low risk	ISRCTN72253427

Other bias	Low risk	Appears to be free of other potential biases
------------	----------	----------------------------------------------

**Rubel 2010**

Methods	Randomized to pretest + decision aid + post-test vs decision aid + post-test vs pretest + posttest vs posttest
Participants	50 + 50 + 50 + 50 men considering prostate cancer screening in the USA
Interventions	DA: booklet on options' outcomes, clinical problem, outcome probabilities, others' opinions + pretest and post-test Comparator : booklet on options' outcomes, clinical problem, outcome probabilities, others' opinions + post-test Comparator: pretest + post-test Comparator: post-test
Outcomes	Knowledge (pre, post-DA), decisional anxiety (post-DA), decisional conflict (post-DA), participation in decision making (pre, post-DA), schema for PSA testing (pre, post-DA), perception of quality and interpretation of recommendation (post-DA)
Notes	Primary outcome was not specified

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Electronically generated random number sequence (p 309, Study design section)
Allocation concealment (selection bias)	Low risk	They were given sealed, sequentially numbered packets (p 309, Study design section)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No mention of blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding, but the outcomes were objectively measured and not subject to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	Protocol followed CONSORT checklist (p 310, Study design section)
Other bias	Low risk	Appears to be free of other potential biases

## Ruffin 2007

Methods	Randomized to decision aid vs usual care
Participants	87 + 87 community dwelling adults not previously screened for CRC in the USA
Interventions	DA: interactive website with information on options' outcomes, clinical problem, outcome probability, explicit values clarification, others' opinion, guidance Comparator: non-interactive website with information on clinical problem
Outcomes	Primary outcome: uptake of option
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"A block randomisation process programmed by the study computer support staff and verified by a statistician was used including two strata, race and gender" (p 3)
Allocation concealment (selection bias)	Unclear risk	No information provided
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Both blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The investigators, data collectors, data entry, and data analyst were all blinded to study arm assignment
Incomplete outcome data (attrition bias) All outcomes	Low risk	Flow diagram (p 3)
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Low risk	Appears to be free of other potential biases

## Sawka 2012

Methods	Randomized to decision aid vs usual care
Participants	37 + 37 individuals with early-stage papillary thyroid cancer
Interventions	DA: web-based decision aid with clinical problem, options' outcomes, outcome probabilities, guidance, printout summary Comparator: usual care (consultation with a specialized head and neck surgeon, and with 1 or more medical specialist)

Outcomes	Primary outcomes: knowledge (baseline and immediately post intervention) Secondary outcomes: decisional conflict, undecided, treatment decision (baseline, immediately post intervention, 6 to 12 months), individual primarily responsible for the treatment decision (6 to 12 months)
Notes	Trial registration: NCT01083550

<i>Risk of bias</i>			<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement	
Random sequence generation (selection bias)	Low risk	"Central computerized randomization in a 1:1 ratio was performed at a patient level by using variable block sizes of 2 and 4 (allocation designed by a study statistician)" (p 2908)	
Allocation concealment (selection bias)	Low risk	"Before the random assignment/testing visit, neither the participant, study staff, investigators, nor treating physicians were aware of the allocation, because it had not yet been assigned" (p 2908)	
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"There was no blinding of participants, study staff, or treating physicians after random assignment was completed" (p 2908), yet it is unlikely that the outcomes are affected by the lack of blinding	
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	"There was no blinding of participants, study staff, or treating physicians after random assignment was completed. However, the statistician was blinded to the allocation of groups at the time of data analysis." (p 2908)	
Incomplete outcome data (attrition bias) All outcomes	Low risk	There does not appear to be any missing outcome data	
Selective reporting (reporting bias)	Unclear risk	Authors state the trial is registered, but no link to trial number	
Other bias	Low risk	Appears to be free of other potential sources of bias	



## Schroy 2011

Methods	Randomized to detailed vs simple decision aid vs control
Participants	223 + 212 + 231 average-risk patients considering CRC screening in the USA
Interventions	Detailed DA: CRC risk assessment + web-based interactive audio-visual DA on options' outcomes, clinical problem, outcome probabilities, others' opinion and guidance Comparator 1: web-based decision aid only Comparator 2: usual care using pamphlet
Outcomes	Knowledge (pre and post-DA), satisfaction with decision making process (pre and post-DA), preferred choice (pre and post-DA)
Notes	Primary outcome was not specified

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No mention of randomization process
Allocation concealment (selection bias)	Unclear risk	No mention of allocation concealment
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Providers were not blinded, subjective outcomes such as satisfaction with decision-making process could have been affected, unclear if participants were blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Assessors not blinded but outcome measures not believed to be influenced by it
Incomplete outcome data (attrition bias) All outcomes	Low risk	No data appears to be missing
Selective reporting (reporting bias)	Unclear risk	No mention of examination of selective outcome reporting or study protocol
Other bias	Low risk	Appears to be free of other sources of bias

## Schwalm 2012

Methods	Randomized to decision aid vs usual care
Participants	76 + 74 patients undergoing coronary angiography

**Schwalm 2012** (Continued)

Interventions	DA: booklet on options' outcomes, clinical problem, outcome probabilities, explicit values clarification and guidance Comparator: usual care
Outcomes	Primary outcomes: decisional conflict Secondary outcomes: knowledge, risk perception, value congruent with chosen option
Notes	-

<b>Risk of bias</b>		<b>Risk of bias</b>
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Computerized random number generator (p 261, Study design)
Allocation concealment (selection bias)	Low risk	Sealed envelopes (p 261, Study design)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Patients and physicians were not blinded to the allocation (p 261, Study design)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear if DCS score assessed by unblinded individuals, but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Did not seem to have incomplete data
Selective reporting (reporting bias)	Low risk	Protocol is available
Other bias	Low risk	Appeared to be free of other biases

**Schwartz 2001**

Methods	Randomized to decision aid vs usual care
Participants	181 + 190 Ashkenazi Jewish women considering genetic testing in the USA
Interventions	DA: 16-page booklet on genetic testing with options' outcomes, clinical problem Comparator: general information on breast cancer, <i>Understanding Breast Changes: A Health Guide for all Women</i> , published by the National Cancer Institute
Outcomes	Primary outcome: preferred option Secondary outcomes: knowledge, accurate risk perceptions
Notes	-

**Schwartz 2001** (Continued)

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated (p 3)
Allocation concealment (selection bias)	Unclear risk	No information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	High retention rate, baseline data and reasons for lost to follow-up were provided (p 2, Participants section)
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Low risk	Appears to be free of other potential biases

**Schwartz 2009a**

Methods	Randomized to decision aid + genetic counselling vs genetic counselling alone
Participants	100 + 114 women considering prophylactic mastectomy for being BRCA1/2 mutation carriers in the USA
Interventions	DA: CD-Rom on options' outcomes, clinical problem, risk communication with individually tailored risk graphs, explicit values clarification, others' opinion; guidance/counselling - genetic counselling as usual care (Ottawa Decision Support Framework) Comparator: genetic counselling on benefits and risks of testing, clinical problem (risk assessment, cancer risks associated with mutations, process of testing and interpretation of results) plus written letter outlining all guidelines and recommendations
Outcomes	Primary outcomes: decisional conflict, satisfaction with decision, actual choice (risk reduction mastectomy) Secondary outcomes: remaining undecided
Notes	-

<i>Risk of bias</i>	<i>Risk of bias</i>
---------------------	---------------------

**Schwartz 2009a** (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomized via computer-generated random number in a 1:1 ratio (p 3, Procedure)
Allocation concealment (selection bias)	Unclear risk	No information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Fig. 1 - flow diagram (p 3)
Selective reporting (reporting bias)	Unclear risk	Protocol not mentioned
Other bias	Low risk	Appears to be free of other sources of bias (p 8) "when variable for not watching DA cd was considered in multivariate models, the results did not change substantively (data not shown)"

**Sheridan 2006**

Methods	Randomized to decision aid vs usual care (list of risk factors)
Participants	49 + 38 adults with no history of cardiovascular disease in the USA
Interventions	DA: computerized decision aid on options' outcomes, outcome probabilities Comparator: list of CHD risk factors to present to doctor
Outcomes	Patient-practitioner communication (e.g. discussion with doctor, specific plan to reduce risk discussed with doctor)
Notes	Primary outcome was not specified

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"[C]omputerized random number generator" (p 2)

**Sheridan 2006** (Continued)

Allocation concealment (selection bias)	Low risk	"[S]ealed in security envelopes" (p 2)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Participants were blinded but the doctors who saw both groups were not
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcome was patient reported
Incomplete outcome data (attrition bias) All outcomes	Low risk	Results (p 5); Flow diagram (p 10); Baseline characteristics/data included
Selective reporting (reporting bias)	Low risk	ClinicalTrials.gov NCT00315978
Other bias	Low risk	Appears to have no other potential risk of bias

**Sheridan 2011**

Methods	Randomized to decision aid + tailored messages vs usual care
Participants	81 + 79 patients with moderate or high risk for CHD considering CHD prevention strategies in the USA
Interventions	DA: web-based decision aid on options' outcomes, clinical problem, outcome probabilities, explicit values clarification and guidance Comparator: usual care using computer programme
Outcomes	Preferred choice (post-DA), adherence Other outcomes (Sheridan 2014): patient-provider communication (post-DA), patient participation (post-DA), patients perceptions of discussions and the health care visit (post-DA), preferred choice (baseline and post-DA) (data from 81 +79 patients)
Notes	Primary outcome was not specified

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"Patients were randomised by study staff who accessed an online randomised schedule" (p 2). Sequence generation method not stated
Allocation concealment (selection bias)	Low risk	"Patients were randomised by study staff who accessed an online randomised schedule" (p 2)

**Sheridan 2011** (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	Patients blinded and physicians unblinded but objective outcomes are not likely affected by lack of blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcomes deemed objective therefore lack of blinding did not influence assessment
Incomplete outcome data (attrition bias) All outcomes	Low risk	There appears to be no missing data
Selective reporting (reporting bias)	Low risk	Protocol made available
Other bias	Low risk	Appears to be free of other sources of bias

**Shorten 2005**

Methods	Randomized to decision aid vs usual care
Participants	85 + 84 pregnant women who have experienced previous cesarean section considering birthing options in Australia
Interventions	DA: decision aid booklet on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, guidance (Ottawa Decision Support Framework) Comparator: usual care
Outcomes	Primary outcomes: knowledge, decisional conflict Secondary outcomes: preferred option, help with making a decision
Notes	-

***Risk of bias***

***Risk of bias***

<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Computer-based randomized generation (p 3, Procedure)
Allocation concealment (selection bias)	Low risk	"[O]paque envelopes containing a random allocation for each participant code number" (p 3)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Participants/midwives/ doctors were blinded to patients' allocation. However, women who used the decision aid as specified and in a process of consultation with their midwife or doctor would have negated the blinding of their clinicians, and

## Shorten 2005 (Continued)

		perhaps of the women themselves. For the intervention group, this may have affected the level and type of information exchanged between them and their caregivers
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	16 women were lost to follow-up from the intervention group and 18 from the control group (no reasons listed) (p 4, Results)
Selective reporting (reporting bias)	Low risk	Reference to published protocol
Other bias	Low risk	Appears to be free of other potential biases

## Shourie 2013

Methods	Cluster-randomized controlled trial of GP practices to web-based MMR DA + usual care, MMR leaflet + usual care, versus usual care
Participants	50 + 93 + 77 parents' of children facing their first dose MMR vaccination
Interventions	Web-based DA: clinical problem, options' outcomes, explicit values clarification, guidance MMR leaflet: Health Scotland leaflet, 'MMR: your questions answered' Comparator: usual care
Outcomes	Primary outcomes: decisional conflict (baseline and 2 weeks postintervention) Secondary outcomes: choice uptake of first dose MMR (when child was 15 months), knowledge (baseline and 2 weeks; results not provided), MMR immunization cognitions (baseline and 2 weeks post; results not provided), immunization trade-off beliefs (baseline and 2 weeks post; results not provided), anxiety (baseline and 2 weeks post; results not provided), use of the intervention (baseline and 2 weeks post)
Notes	Trial registration: UK Clinical Research Network - UKCRN ID 4811

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Simple randomisation using a computer-generated random list allocated GP practices on a 1:1:1 basis" (p 3)

**Shourie 2013** (Continued)

Allocation concealment (selection bias)	Low risk	“An independent researcher who had no contact with participants generated the allocation sequence and assigned the GP practices to their allocated arm” (p 3)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	“On receipt of the completed baseline questionnaire and consent form, the appropriate intervention was delivered. At this point the researchers and participants were no longer blind to allocation” (p 3). We don't know if receiving the intervention had an effect on the ultimate decision that was made
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcome data assessment does not depend on the assessor. It is an objective questionnaire
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing primary outcome data.
Selective reporting (reporting bias)	Unclear risk	Protocol registered. Primary outcome reported as stated. Secondary outcomes are not reported (p 3)
Other bias	Unclear risk	Difference in allocation to groups (50 + 93 + 77). Unclear what effect this difference had on the results

**Smith 2010**

Methods	Randomized to detailed vs simple decision aid vs usual care
Participants	196 + 188 + 188 socioeconomically disadvantaged participants diagnosed with average or slightly above average risk of bowel cancer considering bowel cancer screening in Australia
Interventions	DA: booklet + DVD + worksheet + question prompt list on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, guidance (step-by-step process for making the decision; worksheet; encourages patients to communicate with practitioners by asking questions; summary) Comparator: booklet + DVD + worksheet on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, guidance (step-by-step process for making the decision; worksheet; encourages patients to communicate with practitioners by asking questions; summary) Comparator: usual care using standard information booklet



Outcomes	<p>Primary outcomes: values congruent with chosen option (post-DA), participation in decision making (pre, post-DA)</p> <p>Secondary outcomes: knowledge (pre, post-DA), attitude, actual choice (post-DA), decisional conflict (post-DA), decision satisfaction (post-DA), confidence in decision making (post-DA), general anxiety (post-DA), worry about developing bowel cancer (pre, post-DA), risk perception</p> <p>Other outcomes (Smith 2014): screening participation (357 + 173 participants)</p>
----------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Notes	-
-------	---

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Participants who verbally consented to take part were then randomised to one of the three groups using random permuted blocks of size 6 and 9 for each sex stratum" (p 3, Participants and recruitment section)
Allocation concealment (selection bias)	Low risk	Central allocation; "interviewers responsible for recruiting participants were not aware of the randomization sequence or allocation and therefore did not know which intervention respondents would receive" (p 3, Participants and recruitment section)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"It was not possible for the reviewers to be blinded to the group allocation. However, all questions used standardised wording with pre-coded responses and were asked within a supervised environment, where interviewer performances were regularly monitored to ensure scripts were read as written" (p 3, Outcome measures section)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"[A]nalyses were by intention to treat and carried out blinded to intervention" (p 5, Statistical analysis section); outcomes measured were not subject to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Explanation for the missing data reported at base of tables
Selective reporting (reporting bias)	Low risk	Study protocol available (ClinicalTrials.gov NCT00765869 and Australian New Zealand Clinical Trials Registry

**Smith 2010** (Continued)

		12608000011381)
Other bias	Low risk	Appears to be free of other potential sources of bias

**Stacey 2014a**

Methods	Randomized to decision aid vs usual care
Participants	71 + 71 adults diagnosed with knee osteoarthritis considering joint replacement in Canada
Interventions	DA: DVD + booklet + worksheet on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, others' opinion, guidance (1 page summary for the surgeon) Comparator: usual care
Outcomes	Primary outcomes: feasibility (including recruitment, data collection), preliminary effectiveness Secondary outcomes: knowledge (post-DA, pre-surgeon consult), informed values-congruent with chosen option (post-DA, pre-surgeon consult), uptake of chosen option at 1 year; decisional conflict (SURE test), preparation for decision making (4 items), wait times
Notes	Trial registration: NCT00743951

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The allocation schedule was computer-generated centrally by a statistician using a permuted block design with randomly varying block lengths of 4, 6, or 8." (p 3)
Allocation concealment (selection bias)	Low risk	"Allocations were concealed in numbered opaque sealed envelopes" (p 3)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Patients were not informed of the intervention characteristics" (p 3)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Although the research assistant was not blinded to group allocation, study outcomes for effectiveness were objective and obtained from clinic data (e.g. date of surgery or wait list status)" (p 3)

Stacey 2014a (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data
Selective reporting (reporting bias)	Low risk	Protocol registered on ClinicalTrials.gov
Other bias	Low risk	Appears to be free of other potential sources of bias

Steckelberg 2011

Methods	Randomized to decision aid vs usual care
Participants	785 + 792 patients with no CRC history considering CRC screening in Germany
Interventions	DA: brochure on options' outcomes, clinical problem, and outcome probabilities Comparator: usual care using pamphlet
Outcomes	Primary outcomes: values congruent with chosen option (post-DA) Secondary outcomes: knowledge (post-DA), combination of actual and planned uptake (post-DA), risk perception
Notes	-

*Risk of bias*

*Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer generated sequence (p 2, Randomization and blinding)
Allocation concealment (selection bias)	Low risk	Allocation was concealed. Identity numbers were independent of allocation, and study members did not have access to the data. (p 2, Randomization and blinding)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Trial staff who sent out questionnaires and reminders were not aware of study arm, unclear if participants were blinded (p 2, Randomization and blinding)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Trial staff and statistician who entered data were blinded (p 2, Randomization and blinding)
Incomplete outcome data (attrition bias) All outcomes	Low risk	12% missing one or both questionnaires in intervention group vs 9.2% in control; judged to have low impact on study out-

		come (p 2)
Selective reporting (reporting bias)	Low risk	Protocol available
Other bias	Unclear risk	Participants who completed the trial do not add up

## Taylor 2006

Methods	Randomized to print DA versus video DA versus wait list control
Participants	98 + 95 + 92 African American men with no history of prostate cancer to consider prostate cancer screening
Interventions	Print DA: clinical problem; outcome probabilities; guidance (list of questions to ask at next appointment); others' opinions Video DA: clinical problem; others' opinions Wait list comparator: no information provided until 1 month postrandomization (baseline assessment for this group coincided with 1-month assessment of print and video arms)
Outcomes	Prostate cancer screening intention (baseline and 1 month; not reported), prostate screening uptake (1 year; not included because wait list received intervention before 1 year) process variables including use and perception of the intervention materials (1 month), prostate cancer knowledge (baseline and 1 month post), decisional conflict (baseline and 1 month post), satisfaction with screening decision (baseline and 1 month post)
Notes	No primary outcome reported; not found in trials registry

***Risk of bias******Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information related to random sequence generation
Allocation concealment (selection bias)	Unclear risk	Insufficient information to judge allocation concealment
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to judge blinding; however, participants were requested to not share intervention materials with others to prevent contamination between groups (p 2180)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to judge blinding of outcome assessment

**Taylor 2006** (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Does not appear to be missing any outcome data
Selective reporting (reporting bias)	Unclear risk	No protocol registered or published
Other bias	Unclear risk	<p>“All participants were mailed \$25 for their participation following completion of the 1-month interview” (p 2181)</p> <p>“Men who reported that they had not yet had a chance to read/watch the materials were given an additional week to do so and called again to complete the follow-up assessment” (p 2181)</p>

**Thomson 2007**

Methods	Randomized to decision aid vs usual care by clinical guidelines
Participants	69 + 67 patients with atrial fibrillation considering treatment options in the UK
Interventions	DA (in consultation): computerized decision on options’ outcomes, clinical problem, outcome probabilities, explicit values clarification, guidance/coaching by physician Comparator: guidelines applied as direct advice
Outcomes	Primary outcome: decisional conflict Secondary outcomes: anxiety, knowledge, resource use, choice, health outcomes (stroke, transient ischaemic attack, bleeding events)
Notes	-

***Risk of bias***

***Risk of bias***

Bias	Authors’ judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	“[E]lectronically-generated random permuted blocks via a web-based randomisation service” (p 2, Recruitment and randomization)
Allocation concealment (selection bias)	Low risk	“[E]lectronically-generated random permuted blocks via a web-based randomisation service” (p 2, Recruitment and randomization)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Physicians were blinded. Unclear if patients are blinded and how that may affect the outcome

**Thomson 2007** (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	See flow diagram
Selective reporting (reporting bias)	Low risk	ISRCTN24808514
Other bias	Low risk	Baseline characteristics similar, sample size similar, not stopped early

**Trevena 2008**

Methods	Randomized to decision aid vs usual care by consumer guidelines
Participants	157 + 157 patients not previously screened for colorectal cancer in Australia
Interventions	DA: age-gender-family history specific DA booklet with information on options, outcome probabilities, explicit values clarification, guidance (personal worksheet with steps in decision making) (Theory of planned behaviour) Comparator: consumer guidelines recommending faecal occult blood testing
Outcomes	Primary outcome: informed choice Secondary outcomes: knowledge, values, screening intention (choice); test uptake, anxiety, acceptability of the intervention, satisfaction with the decision
Notes	-

***Risk of bias***

***Risk of bias***

<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	"Sequential ID numbers were randomly assigned by computer program to DA or Guidelines (G) in blocks of four" (p 3)
Allocation concealment (selection bias)	Low risk	"Allocation was concealed via the password-protected program" (p 3)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Participants were blinded to the intervention type - not sure about GPs
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Researchers were blinded to allocation for all telephone interviews, outcomes were objectively measured

**Trevena 2008** (Continued)

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Baseline characteristics included (p 3). Fig 2 flow chart (p 5). Reasons for loss to follow-up not mentioned
Selective reporting (reporting bias)	Low risk	ClinicalTrials.gov - NCT00148226
Other bias	Low risk	Appears to be free of other potential biases

**Van Peperstraten 2010**

Methods	Randomized to decision aid vs usual care
Participants	152 + 156 infertile women on wait list for in vitro fertilization in the Netherlands
Interventions	DA: self-administered booklet on options' outcomes, clinical problem, outcome probabilities, explicit values clarification, guidance (step-by-step process for making decision, worksheet with questions relevant to decision-making process; 1 or more questions that asked patients to clarify their preferences; summary to be shared with practitioner) , coaching (by trained in vitro fertilization nurse) + standard in vitro fertilization care Comparator: standard in vitro fertilization care, including a session in which the number of embryos transferred was discussed
Outcomes	Primary outcomes: actual choice (postintervention and consult) Secondary outcomes: knowledge (pre, post-DA and consult), empowerment (pre, post-DA and consult), participation in decision making, decisional conflict (post-DA and consult), levels of anxiety (pre, post-DA and consult), depression (pre, post-DA and consult), cost evaluation of empowerment strategy (post-DA and consult), condition-specific health outcomes (pregnancies) (post-DA and consult)
Notes	-

***Risk of bias***

***Risk of bias***

<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Computer generated list (p 2, Methods section)
Allocation concealment (selection bias)	Low risk	Central allocation (p 2, Methods section)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Because of the nature of the intervention it was not possible to blind the participants or in vitro fertilisation doctors to the allocation. Participation in our trial did not change the normal in vitro routine." (p 2, Methods section)

**Van Peperstraten 2010** (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes assessed were not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	There are categories in each column of table 1 (p 3) where the denominators do not match the number of people in the group and no reason was given to explain why this would be or if this affects the study
Selective reporting (reporting bias)	Low risk	Outcomes same as those registered with ClinicalTrials.gov
Other bias	Low risk	The study appear to be free of other sources of bias

**Vandemheen 2009**

Methods	Randomized to decision aid vs usual care
Participants	70 + 79 patients with cystic fibrosis considering referral for lung transplantation in Canada
Interventions	DA: self-administered booklet with clinical problem, outcome probability, explicit values clarification, guidance (Ottawa Decision Support Framework) Comparator: blank pages
Outcomes	Primary outcomes: knowledge, accurate risk perceptions, decisional conflict Secondary outcomes: preparation for decision making, choice, durability of decision, undecided
Notes	-

***Risk of bias***

***Risk of bias***

<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	"[C]omputer-generated random listing of two treatment allocations blocked in blocks of 2 or 4, stratified by site and infection status of <i>Burkholderia cepacia</i> " (p 2)
Allocation concealment (selection bias)	Low risk	Central allocation (p 2)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Single blinded RCT; patients and researchers were blinded but physicians were not because they were involved with patients before being randomized



**Vandemheen 2009** (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Research staff, who were blinded to treatment allocation, telephoned each patient and had them complete a follow-up questionnaire; other outcomes reported are objectively measured
Incomplete outcome data (attrition bias) All outcomes	Low risk	Baseline characteristics included (Flow diagram, p 2)
Selective reporting (reporting bias)	Low risk	Clinical trial registered with www.clinicaltrials.gov (NCT00345449)
Other bias	Low risk	Appears to be free of other potential biases

**Vodermaier 2009**

Methods	Randomized to decision aid vs usual care
Participants	74 + 78 women with breast cancer considering treatment options in Germany
Interventions	DA: Decision board administered by research psychologists and booklet on options' outcomes, clinical problem, outcome probability Comparator: booklet on clinical problem
Outcomes	Primary outcome: decisional conflict Secondary outcomes: choice, length of consultation, satisfaction with decision making, participation in decision making
Notes	-

***Risk of bias***

***Risk of bias***

<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Unclear risk	"Randomisation after the patient gave written informed consent" "Random assignment was performed by means of numbered cards in envelopes" "stratified by age group" (p 2)
Allocation concealment (selection bias)	Low risk	"[N]umbered cards in envelopes" (p 2)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Not blinded - unclear if this would introduce bias to outcome assessed

**Vodermaier 2009** (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Not blinded but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Flow diagram, p 5; baseline characteristics not included
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Low risk	Appears to be free of other potential biases

**Volk 1999**

Methods	Randomized to decision aid vs usual care
Participants	80 + 80 men considering PSA testing in the USA
Interventions	DA: Health Dialog videotape and brochure on options' outcomes, clinical problem, outcome probability, others' opinion Comparator: usual care
Outcomes	Primary outcomes: knowledge, preferred/uptake of option
Notes	-

***Risk of bias***

***Risk of bias***

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Volk 1999 (primary study), p 3: "[r]andomization by permuted blocks" "Each block included the numbers 1 through 4"; Volk 2003, p 2, Methods: Randomization by permuted blocks was used to balance the number of subjects in each arm of the study
Allocation concealment (selection bias)	Unclear risk	Volk 1999 (primary study): no information provided Volk 2003, p 2: "[d]etails of the study procedures, subjects, and 2-week follow-up results can be found elsewhere"
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Participants were not blinded to the treatment assignment, but the physicians were; therefore outcomes were unlikely to be biased

**Volk 1999** (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Interviewers were not blinded but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Volk 1999 (primary study), p 2, Procedures: baseline values included Volk 2003, p 4 Fig 1 - flow diagram; baseline data not included
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Low risk	Volk 1999 (primary study): appears to be free of other potential biases Volk 2003: appears to be free of other sources of bias

**Vuorma 2003**

Methods	Randomized to decision aid vs usual care
Participants	184 + 179 women considering treatment for menorrhagia in Finland
Interventions	DA: booklet on options' outcomes, clinical problem, outcome probability Comparator: usual care
Outcomes	Primary outcomes: uptake of option Secondary outcomes: knowledge, proportion remaining undecided, anxiety, satisfaction, health outcomes, use and cost of healthcare services
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Vuorma 2003 (primary study), p 2, Randomization: computer-generated; done by a researcher who did not participate in the planning or concealment procedures "[D]one in STAKES, by researcher separately for each hospital in computer-generated varying clusters"(p 2) Vuorma 2004: no information provided
Allocation concealment (selection bias)	Low risk	Vuorma 2003 (primary study), p 2 "sequentially numbered, opaque and sealed envelopes" Vuorma 2004, p 2 "sequentially numbered,

**Vuorma 2003** (Continued)

		opaque, sealed envelopes"
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No blinding, unclear if measurements could be influenced by lack of blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Study staff were not blinded but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Vuorma 2003 (primary study): flow chart balanced. Reasons for non-eligibility. "One women on HRT was randomized by mistake and included in analyses." Baseline characteristics included and balanced across groups (p 4-5) Vuorma 2004, flow diagram (p 3)
Selective reporting (reporting bias)	Unclear risk	Vuorma 2003 (primary study): no mention of study protocol Vuorma 2004: no information provided
Other bias	Low risk	Vuorma 2003 (primary study), p 7: "increase in knowledge in both study groups, carry-over effect; change in decision-making process of intervention group may have altered physician's negotiation with patients" appears to be free of other potential biases Vuorma 2004, p 5: "comparison of the baseline characteristics presented elsewhere" In the pre-trial group compared with the control group, there was a greater increase in the dimensions of physical role functioning and emotional role functioning of the RAND-36

**Watson 2006**

Methods	Randomized to decision aid vs usual care
Participants	475 + 522 men considering prostate cancer screening in the UK
Interventions	DA: leaflet on options' outcomes, clinical problem, outcome probability Comparator: usual care

**Watson 2006** (Continued)

Outcomes	Primary outcomes: knowledge, screening intention, attitudes Secondary outcomes: preferred role in decision making
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"[R]andom numbers generated centrally by Stata v8.2" (p 3)
Allocation concealment (selection bias)	Low risk	"[R]andom numbers generated centrally by Stata v8.2" (p 3)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No information provided
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Flow diagram (p 2); reason for exclusion from analysis mentioned. Sample characteristics of risk included
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Unclear risk	"Adjustment for multiple testing was not accounted for and hence a degree of caution with interpretation is required, particularly in relation to findings with a P-value close to 0.05" (p 3)

**Weymiller 2007**

Methods	Cluster-randomized to decision aid vs usual care
Participants	51 + 46 patients with type 2 diabetes in the USA
Interventions	DA (in consultation): 1-page decision aid options' outcomes, clinical problem, tailored outcome probability, guidance/coaching Comparator: booklet on cholesterol management
Outcomes	Primary outcomes: knowledge, decisional conflict Secondary outcomes: consultation length, acceptability of the intervention, adherence,

	estimated personal risk, trust, patient participation (OPTION), choice
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated allocation sequence (p 2) Nannenga 2009: no information provided
Allocation concealment (selection bias)	Low risk	Computer-generated allocation sequence, unavailable to personnel enrolling patients. "[W]ith concealed allocation" (Abstract); "maintained allocation concealment" (p 5); randomized by concealed central allocation (Nannenga 2009, p 2)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Participants and clinicians blinded to the study objectives, providers and patients were naive to this study objective
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Data analysts and statisticians blinded to allocation; intervention and outcomes; adequate blinding wherever possible
Incomplete outcome data (attrition bias) All outcomes	Low risk	Flow diagram (p 3); reasons for attrition mentioned (p 4); baseline characteristics included; flow diagram Nannenga 2009, p 3: reasons for attrition mentioned and study groups balanced; baseline characteristics included
Selective reporting (reporting bias)	Low risk	ClinicalTrials.gov identifier: NCT00217061
Other bias	Low risk	Enrollment of patients already receiving statin therapy and limited statin uptake decreased the precision of our results; results should best be interpreted as preliminary and requiring verification Nannenga 2009: appears to be free of other potential biases

## Whelan 2003

Methods	Randomized to decision aid vs usual care
Participants	82 + 93 women with node negative breast cancer considering adjuvant chemotherapy in Canada
Interventions	DA: decision board and booklet on options' outcomes, clinical problem, outcome probability, guidance/coaching Comparator: booklet on clinical problem
Outcomes	Primary outcomes: knowledge, satisfaction of participant Secondary outcomes: preferred option, anxiety, accurate risk perceptions, participation in decision making
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No information provided
Allocation concealment (selection bias)	Low risk	Randomization, which was performed at a central location (p 3)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unable to blind participants in our trial for practical reasons, measures were taken to minimize bias in the design of the study and the assessment of outcomes
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Flow diagram not included. "[O]ne patient excluded from analysis, determined by physician not to be candidate for chemotherapy" (p 4). Baseline data/characteristics included
Selective reporting (reporting bias)	Unclear risk	Unclear if lack of blinding contributed to potential risk of bias
Other bias	Low risk	Appears to be free of other potential biases

# Whelan 2004

Methods	Cluster-randomized to decision aid vs usual care
Participants	94 + 107 women with Stage 1 or 2 breast cancer considering surgery (cluster-RCT with 27 surgeons randomized) in Canada
Interventions	DA: decision board on options' outcomes, outcome probability, guidance/coaching Comparator: usual care
Outcomes	Primary outcomes: preferred option, knowledge, decisional conflict, satisfaction Secondary outcomes: accurate risk perceptions, anxiety
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Does not specify how the sequence was generated; a paired cluster randomization process was used (p 2, Study design and procedures)
Allocation concealment (selection bias)	Unclear risk	Randomly assigned in a concealed fashion, but method of concealment was not stated (p 2, Study design and procedures)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	"[C]hose cluster randomization method to avoid contamination that might have occurred if surgeons used decision board for some patients and not others" (p 6); unclear if this would introduce bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Baseline characteristics not included; reasons given for loss of participants
Selective reporting (reporting bias)	Unclear risk	No indication that the trial was registered in a central trials registry
Other bias	Low risk	Appears to be free of other potential biases



## Williams 2013

Methods	Randomized to decision aid at home or in clinic versus usual care at home or in clinic
Participants	134 + 138 + 134 + 137 men aged 40-70 years with no history of prostate cancer who had pre-registered for screening
Interventions	DA: content adapted from the Centers for Disease Control and Prevention's PCS educational tool. Includes clinical problem, treatment options, outcome probabilities, explicit values clarification, others' stories, summary worksheet Comparator: information booklet. A 3-page fact sheet requiring 5 minutes to read. Information presented in a Q&A format on who is recommended for testing, how to interpret results, and the limitations of testing
Outcomes	Knowledge, decisional conflict, screening outcomes, satisfaction with decision Outcomes assessed at baseline, 2 months, 13 months, except satisfaction with decision (2 months and 13 months)
Notes	No primary outcome reported; trial registration not provided

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information to judge random sequence generation
Allocation concealment (selection bias)	Unclear risk	Insufficient information to judge allocation concealment
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to judge blinding of participants and personnel
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to judge blinding of outcome assessment
Incomplete outcome data (attrition bias) All outcomes	Low risk	There does not appear to be any outcome data missing
Selective reporting (reporting bias)	Unclear risk	No registered or published protocol
Other bias	Low risk	Appears to be free of other potential biases

## Wolf 1996

Methods	Randomized to decision aid vs usual care
Participants	103 + 102 men considering PSA testing in the USA
Interventions	DA: script of options' outcomes, clinical problem, outcome probability, others' opinions Comparator: usual care (single sentence)
Outcomes	Preferred option
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Wolf 1996 (primary study): no information provided Wolf 1998, p 2: "the methodology of the randomized trial has been reported previously"
Allocation concealment (selection bias)	Unclear risk	Wolf 1996 (primary study): no information provided Wolf 1998, p 2: "The methodology of the randomized trial has been reported previously"
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Low risk	Wolf 1996 (primary study), p 2: needed a minimum sample size of 150 participants, and was achieved with total sample size of 205. Reasons for attrition mentioned; baseline characteristics included Wolf 1998: no information provided except that methodology of the randomized trial and the content of the informational intervention reported previously (p 2). Baseline characteristics included; flow of participants not included
Selective reporting (reporting bias)	Unclear risk	No indication that the trial was registered in a central trials registry

**Wolf 1996** (Continued)

Other bias	Low risk	Wolf 1996 (primary study): participant population had lower SES therefore external validity of the findings limited, but overall appears to be free of other potential biases Wolf 1998: appears to be free of other potential biases
------------	----------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

**Wolf 2000**

Methods	Randomized to decision aid vs usual care
Participants	266 + 133 elderly ( $\geq 65$ years) considering CRC screening in the USA
Interventions	DA: script of options' outcomes, clinical problem, outcome probabilities Comparator: usual care (5 sentences)
Outcomes	Primary outcome: preferred option Secondary outcomes: accurate risk perceptions
Notes	-

<i>Risk of bias</i>		<i>Risk of bias</i>
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"[P]atients were randomised" (p 2); does not indicate how
Allocation concealment (selection bias)	Unclear risk	No information provided
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Baseline data not included (p 2, Results)
Selective reporting (reporting bias)	Unclear risk	Protocol not mentioned
Other bias	Low risk	Appears to be free of other potential biases

## Wong 2006

Methods	Randomized to decision aid vs placebo control leaflet
Participants	162 + 164 women referred for pregnancy termination in the UK
Interventions	DA: decision aid leaflet on options' outcomes, clinical problem, outcome probability, explicit values clarification Comparator: placebo leaflet on contraception use post pregnancy termination
Outcomes	Primary outcomes: uptake of option, knowledge, decisional conflict, anxiety
Notes	-

### *Risk of bias*

### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"1:1 ratio, balanced block of 10"; "envelope preparation by drawing slips of paper labelled either control or intervention"; "the slip determined leaflet placed into envelope" (p 2)
Allocation concealment (selection bias)	Low risk	Consecutive numbered, opaque trial envelope (p 2, Methods)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Unclear blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Unclear blinding but outcomes were objectively measured and not subjective to interpretation
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Baseline characteristics not included (p 3) ; reasons for attrition and incompleteness mentioned
Selective reporting (reporting bias)	Unclear risk	No information provided
Other bias	Low risk	Appears to be free of other potential biases

**CHD:** coronary heart disease; **CRC:** colorectal cancer; **DA:** decision aid; **HPV:** human papilloma virus; **HRT:** hormone replacement therapy; **NSW:** New South Wales; **OA:** osteoarthritis; **PSA:** prostate-specific antigen; **PTSD:** post-traumatic stress disorder; **RCT:** randomized controlled trial; **SES:** socioeconomic status.

## Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
<a href="#">Abadie 2009</a>	Study did not evaluate the decision aid (evaluated clinician use of the decision aid in one arm of a study only)
<a href="#">Adab 2003</a>	Hypothetical choice, not at a point of decision making
<a href="#">Al Saffar 2008</a>	Study not focused on making a choice; adhering to medications only
<a href="#">Alegría 2014</a>	Not a patient decision aid
<a href="#">Altiner 2007</a>	Not a patient decision aid
<a href="#">Anderson 2011</a>	Not a randomized controlled trial
<a href="#">Arimori 2006</a>	Not a patient decision aid (not including benefits and harms)
<a href="#">Armstrong 2005</a>	Unable to ascertain whether intervention meets criteria to qualify as a patient decision aid; additional information requested from author but not provided
<a href="#">Arterburn 2013</a>	Not evaluating a patient decision aid
<a href="#">Au 2011</a>	Not a randomized controlled trial
<a href="#">Bakken 2014</a>	Not a patient decision aid; related to lifestyle choices
<a href="#">Becker 2009</a>	Hypothetical choice; not at the point of decision making
<a href="#">Belkora 2012</a>	Not a patient decision aid
<a href="#">Bellmunt 2010</a>	Not a patient decision aid
<a href="#">Bennett 2011</a>	Compares 3 versions of the same patient decision aid
<a href="#">Bieber 2006</a>	Study did not evaluate the patient decision aid (evaluated shared decision-making process); not a patient decision aid
<a href="#">Branda 2013</a>	2 patient decision aids with findings aggregated
<a href="#">Brenner 2014</a>	Not a patient decision aid
<a href="#">Breslin 2008</a>	Not a randomized controlled trial
<a href="#">Brown 2004</a>	Not focused on making a choice (no specific decision to be made)
<a href="#">Brundage 2001</a>	Not a randomized controlled trial

(Continued)

Burton 2007	Not a patient decision aid (general patient education only)
Buzhardt 2011	Not evaluating patient decision making
Campbell 2014	Not evaluating a patient decision aid
Carling 2008	Hypothetical choice, not at point of decision making
Causarano 2015	Not a patient decision aid
Chadwick 1991	Not a randomized controlled trial
Chan 2011	Not a patient decision aid
Chewning 1999	Not a randomized controlled trial
Chiew 2008	Not a randomized controlled trial
Clouston 2014	Not a patient decision aid
Col 2007	Unable to ascertain characteristics of the patient decision aid. Additional information requested from author but not provided (e.g. values clarification)
Colella 2004	Not a patient decision aid (describes model of care)
Costanza 2011	Not a randomized controlled trial
Coulter 2003	Not a randomized controlled trial (editorial)
Cox 2012	Not a randomized controlled trial
Crang-Svalenius 1996	Not a randomized controlled trial
Davison 1999	Unable to ascertain whether intervention meets criteria (values clarification) to qualify as a patient decision aid
Davison 2007	Not a patient decision aid
De Boer 2012	Not a randomized controlled trial
De Haan 2013	Not a randomized controlled trial of a patient decision aid
Deen 2012	Not a patient decision aid
Deinzer 2009	Not a patient decision aid
Denig 2014	not a patient decision aid

(Continued)

<a href="#">Deschamps 2004</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Deyo 2000</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Diefenbach 2012</a>	Not a patient decision aid
<a href="#">Dobke 2008</a>	Not focused on making a choice
<a href="#">Dodin 2001</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Donovan 2012</a>	Does not report results of the randomized controlled trial; descriptive article offering techniques of provision of information
<a href="#">Driscoll 2008</a>	Not a patient decision aid
<a href="#">Dunn 1998</a>	Quasi-RCT: randomization was by days of the week
<a href="#">Eaton 2011</a>	Not a decision aid (no decision support)
<a href="#">Eden 2009</a>	Hypothetical choice, not at point of decision making
<a href="#">Eden 2014</a>	The educational brochure (control group) provided information about the options, benefits, and harms making it a simple patient decision aid
<a href="#">Eden 2015</a>	Not a treatment or screening decision
<a href="#">Edwards 2012</a>	Hypothetical choice, not a randomized controlled trial
<a href="#">El-Jawahri 2010</a>	End of life decision
<a href="#">Ellison 2008</a>	Not a randomized controlled trial (Quasi-experimental design); unclear whether at point of decision making
<a href="#">Elwyn 2004</a>	No difference in intervention between arms; risk communication did not have values clarification
<a href="#">Emery 2007</a>	Not a patient decision aid
<a href="#">Emmett 2007</a>	Not a randomized controlled trial
<a href="#">Feldman-Stewart 2006</a>	Hypothetical choice, not at point of decision making
<a href="#">Feldman-Stewart 2012</a>	Same patient decision aid with vs without values clarification
<a href="#">Fiks 2013a</a>	Not patient decision making (uptake of vaccine)
<a href="#">Flood 1996</a>	Non-randomized allocation; wait list control

(Continued)

Francis 2009	Not a patient decision aid
Fraval 2015	Not a patient decision aid; general education material to obtain informed consent for surgery
Frosch 2001	Not a randomized controlled trial
Frosch 2003	Same decision aid delivered on the Internet versus on DVD plus booklet
Frosch 2008b	Not a randomized controlled trial
Frosch 2011	Not a patient decision aid
Frost 2009	Qualitative study for an included RCT
Fujiwara 2015	Not a patient decision aid and aims to increase screening rates
Garvelink 2013	Hypothetical decision
Genz 2012	Not a patient decision aid
Giordano 2014	Not a patient decision aid
Goel 2001	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
Graham 2000	Not a patient decision aid (general information)
Gray 2009	Hypothetical choice, not at the point of decision making
Green 2001b	Not a patient decision aid (educational intervention)
Green 2004	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
Greenfield 1985	Not focused on making a choice (intervention to increase patient involvement in care)
Griffith 2008a	Hypothetical choice, not at the point of decision making
Griffith 2008b	Not a randomized controlled trial
Gruppen 1994	Not a patient decision aid
Gummersbach 2015	Not a patient decision aid and a hypothetical decision
Hacking 2013	Not a patient decision aid
Hall 2007	Not about evaluating a patient decision aid
Hall 2011	Not a patient decision aid



(Continued)

<a href="#">Hamann 2014</a>	not a patient decision aid
<a href="#">Harmsen 2014</a>	Not a patient decision aid
<a href="#">Harwood 2011</a>	Not a randomized controlled trial
<a href="#">Healton 1999</a>	Not a patient decision aid (education to promote compliance)
<a href="#">Henderson 2013</a>	Not a treatment or screening decision
<a href="#">Herrera 1983</a>	Quasi-RCT: assigned to 1 of 2 alternating groups
<a href="#">Hess 2015</a>	Conjoint analysis for values clarification without information on options, pros and cons
<a href="#">Hewison 2001</a>	Not a patient decision aid; no values clarification
<a href="#">Heyn 2013</a>	Not a randomized controlled trial
<a href="#">Hickish 1995</a>	Not a randomized controlled trial (letter)
<a href="#">Hochlehnert 2006</a>	Not a patient decision aid (general information; no values clarification)
<a href="#">Hofbauer 2008</a>	Not a randomized controlled trial
<a href="#">Hoffman 2009</a>	Not a patient decision aid
<a href="#">Holbrook 2007</a>	Hypothetical choice, not at the point of decision making
<a href="#">Hollen 2013</a>	Not a treatment or screening decision
<a href="#">Holloway 2003</a>	Not focused on making a choice (promotes complying with a recommended option)
<a href="#">Holmes-Rovner 2011</a>	Not a randomized controlled trial
<a href="#">Holt 2009</a>	Study does not evaluate a decision aid; evaluation of spiritual versus non-spiritual framework
<a href="#">Hope 2010</a>	Same content
<a href="#">Huijbregts 2013</a>	Not a patient decision aid
<a href="#">Hunt 2005</a>	Not focused on making a choice (promotes complying with a recommended option)
<a href="#">Hunter 1999</a>	Not focused on making a choice (no specific decision)
<a href="#">Hunter 2005</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Huyghe 2009</a>	Hypothetical choice, not at point of decision making for all participants

(Continued)

Ilic 2008	No difference in content of interventions - testing mode of delivery
Isebaert 2007	Not a randomized controlled trial (English paper published in 2008 <i>Urologia Internationalis</i> )
Jackson 2011	Not a patient decision aid
Jerant 2007	Not focused on making a choice - adherence to screening
Jibaja-Weiss 2006	No comparison outcome data provided (only presents data for intervention group)
Joosten 2009	Not a patient decision aid
Joosten 2011	Not a patient decision aid
Jorm 2003	Hypothetical choice, not at point of decision making - community sample asked to evaluate information booklet on depression
Kakkilaya 2011	Hypothetical choice, not at point of decision making
Kaplan 2014a	Not a patient decision aid
Kaplan 2014b	Not randomized controlled trial results; cross-sectional analysis of baseline data
Kassan 2012	Web arm only, not a randomized controlled trial
Kellar 2008	Hypothetical choice, not at point of decision making
Kiatpongsan 2014	No specific decision to be made and not a true randomized controlled trial
Kobelka 2009	Not a randomized controlled trial; not a patient decision aid
Koelewijn-van Loon 2009	Lifestyle only
Krawczyk 2012	Uptake of a recommended option
Kripalani 2007	Not a patient decision aid
Krones 2008	Not a patient decision aid - no benefits and harms
Kuppermann 2009	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
Kurian 2009	Not a randomized controlled trial; not a patient decision aid
Köpke 2009	Not a patient decision aid
Köpke 2014	Not a patient decision aid

(Continued)

<a href="#">Labrecque 2010</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">LaCroix 1999</a>	Inadequate comparison outcome data provided, secondary report of pilot study
<a href="#">Lairson 2011</a>	Not a patient decision aid (to increase uptake of screening)
<a href="#">Lalonde 2006</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Lancaster 2009</a>	Not a patient decision aid
<a href="#">Landrey 2013</a>	Not a patient decision aid
<a href="#">Lazcano Ponce 2000</a>	Not a patient decision aid (no values clarification)
<a href="#">Legare 2003</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Leung 2004</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Levin 2011</a>	Not a patient decision aid
<a href="#">Lewis 2003</a>	Hypothetical choice, not at the point of decision making
<a href="#">Lewis 2012</a>	Uptake of a recommended option
<a href="#">Lopez-Jornet 2012</a>	Not a patient decision aid/not at point of decision-making
<a href="#">Lukens 2013</a>	Not a patient decision aid. Results in response to clinical vignettes (hypothetical scenarios)
<a href="#">Lurie 2011</a>	Not a randomized controlled trial (all patients received DA)
<a href="#">Maisels 1983</a>	Not a patient decision aid (no values clarification)
<a href="#">Mancini 2006</a>	Not about evaluating a patient decision aid
<a href="#">Manne 2009</a>	Not focused on making a choice (about adherence not decision making)
<a href="#">Manns 2005</a>	Not focused on making a choice (Promotes complying with a recommended option)
<a href="#">Markham 2003</a>	Not a patient decision aid (review of patient information pamphlets on pre-operative fasting)
<a href="#">Martin 2012</a>	Hypothetical choice, not at the point of decision making
<a href="#">Maslin 1998</a>	Insufficient outcome data provided in publication; requested from author but not provided
<a href="#">Matlock 2014</a>	End of life

(Continued)

<a href="#">Matloff 2006</a>	Not a patient decision aid - genetic counselling only
<a href="#">Mazur 1994</a>	Hypothetical choice, not at the point of decision making
<a href="#">McCaffery 2007</a>	Not a patient decision aid
<a href="#">McGinley 2002</a>	Not a patient decision aid (no values clarification)
<a href="#">McGowan 2008</a>	Not a patient decision aid
<a href="#">McInerney-Leo 2004</a>	Not a patient decision aid (no risk/benefit information; no values clarification)
<a href="#">McLaren 2012</a>	Not a patient decision aid; hypothetical choice, not at point of decision making
<a href="#">Meropol 2013</a>	Not a patient decision aid
<a href="#">Michie 1997</a>	Unable to ascertain whether intervention meets criteria (values clarification) to qualify as a patient decision aid; additional information requested but author was unable to provide the intervention
<a href="#">Miller 2014a</a>	No specific decision; related to increasing visits to healthcare provider
<a href="#">Miller 2014b</a>	Aims to increase visits to healthcare providers; intervention targeted to partners
<a href="#">Mishel 2009</a>	Not a patient decision aid (information only)
<a href="#">Mohammad 2012</a>	Not a patient decision aid; presents only benefits, not harms
<a href="#">Molenaar 2001</a>	Not a randomized controlled trial
<a href="#">Mulley 2006</a>	Not a randomized controlled trial (editorial)
<a href="#">Myers 2005a</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Myers 2005b</a>	Not a randomized controlled trial (editorial)
<a href="#">Myers 2007</a>	Not a patient decision aid
<a href="#">Myers 2011</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Myers 2013</a>	Uptake of screening
<a href="#">Neubeck 2008</a>	Study protocol, does not appear to be patient decision aid
<a href="#">Newton 2001</a>	Not a randomized controlled trial
<a href="#">O'Cathain 2002</a>	Suite of 8 decision aids (not an efficacy trial)

(Continued)

<a href="#">O'Connor 1999a</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">O'Connor 1996</a>	No patient decision aid - framing effects
<a href="#">O'Connor 1998a</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">O'Connor 2009a</a>	Not a patient decision aid
<a href="#">O'Connor 2011</a>	Not a patient decision aid
<a href="#">Owens 2014A</a>	Not an RCT; doctoral dissertation
<a href="#">Patanwala 2011</a>	Not a patient decision aid
<a href="#">Patel 2014</a>	Not an RCT
<a href="#">Pearson 2005</a>	Not a patient decision aid (focus on provision of information)
<a href="#">Peele 2005</a>	Not a patient decision aid (decision aid only supplies mortality risk information; no risk info; no values clarification)
<a href="#">Petty 2014</a>	Not a randomized controlled trial and not a patient decision aid
<a href="#">Philip 2010</a>	Not a randomized controlled trial, not a patient decision aid (promotes complying with a recommended option)
<a href="#">Phillips 1995</a>	Quasi-RCT: alternating order based on patients' initial appointment sequence
<a href="#">Pignone 2013</a>	Not a patient decision aid; compared the effect of 3 different values clarification methods
<a href="#">Pinto 2008</a>	About clinical trial entry
<a href="#">Powers 2011</a>	Not a patient decision aid
<a href="#">Proctor 2006</a>	Not a patient decision aid (general patient education resource)
<a href="#">Prunty 2008</a>	About a lifestyle choice - whether or not to have a child or have another child if I have multiple sclerosis
<a href="#">Ranta 2015</a>	Not a patient decision aid; intended to increase guideline adherence for transient ischaemic attack/stroke
<a href="#">Rapley 2006</a>	Not a randomized controlled trial
<a href="#">Raynes-Greenow 2009</a>	No difference in intervention content; comparison of presentation formats; audio-guided decision aid versus booklet only
<a href="#">Raynes-Greenow 2010</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)

(Continued)

<a href="#">Rimer 2001</a>	Not focused on making a choice (promotes complying with a recommended option)
<a href="#">Rimer 2002</a>	Not focused on making a choice (promotes complying with a recommended option)
<a href="#">Robinson 2013</a>	Not a patient decision aid
<a href="#">Ronda 2014</a>	Benefits or harms of self-testing are not provided as information on the website; values clarification exercise asks users to qualify value statements as benefits or harms
<a href="#">Rostom 2002</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Roter 2012</a>	Not a patient decision aid
<a href="#">Rothert 1997</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Rovner 2004</a>	Not a randomized controlled trial
<a href="#">Rubinstein 2011</a>	Not a patient decision aid
<a href="#">Ruddy 2009</a>	Not a patient decision aid
<a href="#">Ruehlman 2012</a>	Not a patient decision aid
<a href="#">Ruland 2013</a>	No specific decision to be made
<a href="#">Ryser 2004</a>	Not focused on making a choice (promotes complying with a recommended option)
<a href="#">Sassen 2014</a>	Not a patient decision aid evaluation study; healthcare professionals were recruited, not patients
<a href="#">Saver 2007</a>	Not a patient decision aid - general information; not a specific decision
<a href="#">Sawka 2011</a>	Not a randomized controlled trial
<a href="#">Scaffidi 2014</a>	Not an RCT
<a href="#">Schapira 2000</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Schapira 2007</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Schwartz 2009b</a>	Hypothetical choice, not at the point of decision making
<a href="#">Sears 2007</a>	About do not resuscitate versus initiating cardiopulmonary resuscitation decision
<a href="#">Sequist 2011</a>	Not a patient decision aid (promotes complying with a recommended option)
<a href="#">Shah 2012</a>	Not a patient decision aid, lifestyle choices

(Continued)

<a href="#">Sheppard 2012</a>	Not a randomized controlled trial
<a href="#">Sheridan 2004</a>	Not a randomized controlled trial
<a href="#">Sheridan 2010</a>	Hypothetical choice, not at point of decision making
<a href="#">Sheridan 2012</a>	Not a patient decision aid - no benefits and harms
<a href="#">Sherman 2014</a>	Not a randomized controlled trial
<a href="#">Shirai 2012</a>	Not a patient decision aid
<a href="#">Silver 2012</a>	Hypothetical choice, not at point of decision making
<a href="#">Siminoff 2006</a>	Not a patient decision aid (no discussion of harms)
<a href="#">Simon 2012a</a>	Not a patient decision aid
<a href="#">Simon 2012b</a>	Not a patient decision aid
<a href="#">Smith 2011a</a>	No decision regarding treatment or screening to be made (decision regarding full disclosure)
<a href="#">Smith 2011b</a>	Not a patient decision aid, not an RCT
<a href="#">Solberg 2010</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Sorenson 2004</a>	Not a randomized controlled trial
<a href="#">Sparano 2006</a>	Not a patient decision aid
<a href="#">Stalmeier 2009</a>	Not a randomized controlled trial (about instrument development)
<a href="#">Starosta 2015</a>	Not a patient decision aid - benefits and harms of screening are missing
<a href="#">Stein 2013</a>	End of life
<a href="#">Steiner 2003</a>	Not a patient decision aid (only effectiveness not cons of options; not at point of decision making)
<a href="#">Stephens 2008</a>	Not a randomized controlled trial
<a href="#">Stiggelbout 2008</a>	Not a patient decision aid
<a href="#">Stirling 2012</a>	Not a treatment or screening decision
<a href="#">Street 1995</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Street 1998</a>	Not focused on making a choice (promotes complying with a recommended option)

(Continued)

<a href="#">Sundaresan 2011</a>	Hypothetical choice, not at the point of decision making, not a randomized controlled trial
<a href="#">Tabak 1995</a>	Not a randomized controlled trial
<a href="#">Taylor 2013</a>	Not a patient decision aid - benefits and harms of screening not included
<a href="#">Ten 2008</a>	Not a patient decision aid; about stopping medication use
<a href="#">Thomas 2013</a>	Not a patient decision aid
<a href="#">Thomson 2006</a>	Not a randomized controlled trial; not at point of decision making
<a href="#">Thornton 1995</a>	Unable to ascertain whether intervention meets criteria to qualify as a patient decision aid; additional information requested from author but not provided
<a href="#">Tiller 2006</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Tinsel 2013</a>	Not a patient decision aid
<a href="#">Tomko 2015</a>	Not a patient decision aid - benefits and harms of screening are missing
<a href="#">Ukoli 2013</a>	Not an RCT
<a href="#">Valdez 2001</a>	Not a randomized controlled trial; not focused on making a choice (complying with a recommended option)
<a href="#">Van der Krieke 2013</a>	Not a patient decision aid, no benefits/harms
<a href="#">Van Roosmalen 2004</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
<a href="#">Van Steenkiste 2008</a>	Not a randomized controlled trial
<a href="#">Van Til 2009</a>	Hypothetical choice, not at the point of decision making
<a href="#">Van Tol-Geerdink 2013</a>	Not a randomized controlled trial; insufficient information to judge random sequence generation, allocation concealment, and blinding
<a href="#">Veroff 2012</a>	Not a patient decision aid
<a href="#">Volandes 2009</a>	Advanced care planning options
<a href="#">Volandes 2011</a>	Hypothetical choice, end-of-life decision
<a href="#">Volandes 2013</a>	Advanced care planning
<a href="#">Volk 2008</a>	Simple versus detailed patient decision aid (excluded in update after 2014 publication)



(Continued)

Von Wagner 2011	Not a randomized controlled trial (commentary)
Wagner 1995	Not a randomized controlled trial
Wakefield 2008a	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
Wakefield 2008b	Simple versus detailed patient decision aid (excluded in update after 2014 publication)
Wakefield 2008c	simple versus detailed patient decision aid (excluded in update after 2014 publication)
Wallston 1991	Not a patient decision aid - patient preference study
Wang 2004	Not a patient decision aid - intent of intervention to facilitate genetic counselling process, no focused decision
Warner 2015	Not a treatment or screening decision
Watts 2014	Simple versus detailed patient decision aid
Welschen 2012	Not a patient decision aid
Wennberg 2010	Same decision aid in both groups
Westermann 2013	Not a patient decision aid
Weymann 2015	Patients not at the point of decision making
Wilhelm 2009	Not a patient decision aid
Wilkes 2013	Unable to ascertain characteristics of the patient decision aid. Additional information requested from author but not provided (e.g. values clarification)
Wilkie 2013	Not treatment or screening decision
Wilkins 2006	Not a randomized controlled trial
Willemsen 2006	Lifestyle change
Williams-Piehot 2008	Not a randomized controlled trial
Williamson 2014	Lifestyle decision - not treatment or screening
Woltmann 2011	Not a patient decision aid
Wroe 2005	Not focused on making a choice - promotes complying with a recommended option
Yee 2014	Not a patient decision aid

(Continued)

<a href="#">Yun 2011</a>	End-of-life decision
<a href="#">Zajac 2012</a>	Hypothetical
<a href="#">Zapka 2004</a>	Not focused on making a choice - promotes complying with a recommendation
<a href="#">Zikmund-Fisher 2008</a>	No difference in intervention content - comparison of presentation of probabilities
<a href="#">Zoffman 2012</a>	Not a randomized controlled trial, not a patient decision aid

### Characteristics of ongoing studies *[ordered by study ID]*

#### [ACTRN12615000523505](#)

Trial name or title	The motherhood choices decision aid for women with rheumatoid arthritis increases knowledge and reduces decisional conflict: a randomized controlled study
Methods	RCT
Participants	130 women diagnosed with rheumatoid arthritis and currently under the care of a rheumatologist
Interventions	Patient decision aid vs usual care
Outcomes	Decisional conflict, knowledge, anxiety, depression, self-efficacy
Starting date	May 2015
Contact information	Tanya Meade; School of Social Science and Psychology University of Western Sydney; Sydney, Australia
Notes	Trial #: ACTRN12615000523505

#### [ACTRN12615000843550](#)

Trial name or title	Evaluation of decision aids for parents about the benefits and harms of antibiotic use for coughs and colds in children
Methods	Pilot RCT
Participants	108 adult parents or primary caregivers of a child
Interventions	Patient decision aid vs usual care
Outcomes	Informed choice, knowledge, attitudes towards antibiotic use, intention to use antibiotic, decisional conflict, confidence in decision-making, usability and accessibility of the written materials
Starting date	August 2015

**ACTRN12615000843550** (Continued)

Contact information	Mr Peter D Coxeter; pcoxeter@bond.edu.au; Bond University, Queensland, Australia
Notes	ACTRN12615000843550

**Al-Itejawi 2015**

Trial name or title	(Cost-)effectiveness and implementation of a decision aid for patients with prostate cancer
Methods	Stepped wedge cluster RCT
Participants	Newly diagnosed adult participants with localized prostate cancer
Interventions	Patient decision aid vs usual care
Outcomes	Decisional conflict, quality of life, treatment preferences, participation in decision making, knowledge, patient-provider communication
Starting date	May 2015
Contact information	Hoda Al-Itejawi; Afdeling Urologie, Amsterdam, the Netherlands
Notes	Trial #: NTR5177

**Anderson 2014**

Trial name or title	Shared decision making in the emergency department: Chest Pain Choice Trial (CPC)
Methods	RCT
Participants	Presenting to the emergency department with chest pain
Interventions	Chest Pain Choice decision aid vs usual care
Outcomes	Knowledge, patient engagement, decisional conflict, satisfaction, adverse events, admissions, healthcare utilization
Starting date	October 2013
Contact information	Erik P Hess, Mayo Clinic
Notes	NCT01969240; verified September 2014, estimated study completion March 2016

**Aslani 2014**

Trial name or title	Computerized decision aid on mode of delivery
Methods	Cluster RCT
Participants	Pregnant Iranian women
Interventions	Computerized decision aid
Outcomes	Decisional conflict, knowledge
Starting date	Not reported
Contact information	Azam Aslani, Mashhad University, Iran
Notes	-

**Buhse 2013**

Trial name or title	Efficacy of an evidence-based informed shared decision making program for prevention of myocardial infarction in type 2 diabetes
Methods	RCT
Participants	154 patients with type 2 diabetes
Interventions	Shared decision-making programme consisting of a decision aid booklet and a curriculum for group counselling vs placebo counselling
Outcomes	Knowledge, sustainability of knowledge, achievement of individual treatment goals, achievement of treatment goals prioritized by individual patients, medication uptake
Starting date	March 2013
Contact information	Matthias Lenz, University of Hamburg
Notes	ISRCTN84636255

**Carroll 2012**

Trial name or title	Development of and feasibility testing of decision support for patients who are candidates for an implantable defibrillator
Methods	RCT
Participants	Referred for consideration of an implantable cardioverter-defibrillators (non-cardiac resynchronization therapy) for a primary prevention indication

**Carroll 2012** (Continued)

Interventions	Patient decision aid provided prior to the consultation with the physician, which provides a lay summary that outlines the facts, risks, benefits (including probabilities), specific to the option of an implantable defibrillator or the option of medical management vs usual care
Outcomes	Decision aid development and evaluation, decisional conflict and decision quality, sure test, reparation for decision-making scale, medical outcomes trust short form (SF-36v2)
Starting date	June 2012
Contact information	Sandra Carroll, McMaster University
Notes	Trial #: NCT01876173

**Chambers 2008**

Trial name or title	ProsCan for Men: randomized controlled trial of a decision support intervention for men with localised prostate cancer
Methods	RCT
Participants	700 men newly diagnosed with localized prostate cancer
Interventions	A tele-based nurse delivered 5-session decision support/psychosocial intervention vs usual care
Outcomes	Cancer threat appraisal; decision-related distress and bother from treatment side effects; involvement in decision making; satisfaction with health care; healthcare utilization; use of healthcare resources; and a return to previous activities
Starting date	Not yet assessed
Contact information	Suzanne K Chambers, Griffith University
Notes	Trials #: ACTRN012607000233426

**Coylewright 2012**

Trial name or title	Shared decision making in patients with stable coronary artery disease: PCI Choice
Methods	RCT
Participants	-
Interventions	-
Outcomes	-
Starting date	-

**Coylewright 2012** (Continued)

Contact information	Megan Coylewright, Mayo Clinic
Notes	Upcoming RCT

**Cuypers 2015**

Trial name or title	Prostate cancer patient-centered care: impact of a treatment decision aid in a pragmatic, cluster randomized controlled trial
Methods	Pragmatic RCT
Participants	400 men newly diagnosed with early stage prostate cancer
Interventions	Decision aid (online) vs usual care
Outcomes	Decisional conflict, decisional regret, treatment satisfaction, decision making role, knowledge, satisfaction with decision-making process, preparation for decision-making, health-related quality of life, personality (anxiety, depression, optimism), skills measures (self-efficacy, health literacy, numeracy)
Starting date	May 2014
Contact information	M Cuypers; M.Cuypers@uvt.nl; Tilburg University Social and Behavioral Sciences Tilburg, the Netherlands
Notes	NTR4554

**Den Ouden 2015**

Trial name or title	Shared decision-making in type 2 diabetes with a support decision tool that takes into account clinical factors, the intensity of treatment and patient preferences
Methods	Cluster RCT
Participants	150 adults with type 2 diabetes mellitus for 8-15 years
Interventions	Patient decision aid with training vs usual care
Outcomes	Achievement of diabetes-specific health goals, satisfaction with treatment, quality of life, well-being, coping, evidence of shared decision-making
Starting date	March 2012
Contact information	h.denouden@umcutrecht.nl; Henk den Ouden; Julius Centre for Health Sciences and Primary Care, University Medical Centre, Utrecht, the Netherlands
Notes	Trial #: NCT02285881

**Dirmaier 2013**

Trial name or title	Tailored, dialogue-based health communication application for patients with chronic low back pain
Methods	RCT
Participants	414 patients with self-reported chronic low back pain
Interventions	Web-based interactive health communication application (IHCA) vs control (standard info)
Outcomes	Knowledge, patient empowerment, website usage, preparation for decision making, decisional conflict
Starting date	2012
Contact information	Martin Härter, University Medical Center Hamburg-Eppendorf
Notes	International Clinical Trials Registry DRKS00003322

**Geiger 2011**

Trial name or title	Investigating a training supporting Shared Decision Making (IT'S SDM 2011): study protocol for a randomized controlled trial
Methods	RCT
Participants	40 physicians that contribute a sequence of 4 medical consultations including a diagnostic or treatment decision
Interventions	A training curriculum for the doctors - intend to stimulate efforts to involve their patients in the decision-making process
Outcomes	Physician-patient communication, effect of SDM on perceived quality of the decision process and on the elaboration of the decision, decisional conflict
Starting date	Not yet assessed
Contact information	Friedemann Geiger, University Medical Center Schleswig - Holstein
Notes	Trials #: ISRCTN78716079

**Hersch 2014**

Trial name or title	Effect of information about over detection of breast cancer on women's decision-making about mammography screening
Methods	RCT
Participants	970 women aged 48-50

**Hersch 2014** (Continued)

Interventions	Intervention (evidence-based information booklet including over detection, breast cancer mortality reduction and false positives) vs control information booklet (including mortality reduction and false positives only)
Outcomes	Knowledge, consistency between attitudes and intentions, decision conflict, confidence, regret, anxiety, perceived risk, quality of life
Starting date	June 2014
Contact information	Kirsten McCaffery, University of Sydney
Notes	Australian New Zealand Clinical Trials Registry ACTRN12613001035718

**Hess 2014**

Trial name or title	Shared decision making in parents of children with head trauma: head CT choice
Methods	RCT
Participants	1004 parent-child dyad, seeking care for a child who had blunt trauma above the eyebrows and is positive for at least 1 PECARN clinical prediction rules
Interventions	Patient decision aid vs usual care
Outcomes	Knowledge, engagement in decision-making process, decisional conflict, trust in the physician, satisfaction with the decision-making process, choice, healthcare utilization 7-days post ER visit, rate of clinically important traumatic brain injury
Starting date	April 2014
Contact information	Erik Hess; Mayo Clinic; Rochester, MN
Notes	Trial #: NCT02063087

**Jimbo 2012**

Trial name or title	Decision aid to technologically enhance shared decision making
Methods	RCT
Participants	Patients who are not current with colorectal cancer screening
Interventions	Web based decision aid + interactive component (preferences and risk assessment) vs web based decision aid only
Outcomes	Uptake of screening on patient determinants/preference/intention before the patient-physician encounter, and on shared decision making, concordance and patient intention during/after the patient-physician encounter



**Jimbo 2012** (Continued)

Starting date	May 2012
Contact information	Masahito Jimbo, University of Michigan
Notes	Trial # :NCT01514786; last updated December 2013, estimated study completion October 2014

**Layton 2012**

Trial name or title	Effects of a web-based decision aid on African American men's prostate screening knowledge and behavior
Methods	-
Participants	128 African American men
Interventions	-
Outcomes	-
Starting date	-
Contact information	Beverly Layton, Walden University
Notes	Unpublished thesis

**LeBlanc 2013**

Trial name or title	Translating comparative effectiveness of depression medications into practice by comparing the depression medication choice decision aid to usual care: study protocol for a randomized controlled trial
Methods	RCT
Participants	300 patients
Interventions	Use of the Depression Medication Choice decision aid by patients and their primary care clinician during the clinical encounter vs usual care
Outcomes	Decisional conflict, knowledge, satisfaction, preference in decision making style, patient involvement in decision making, depression outcomes, medication adherence
Starting date	December 2011
Contact information	Victor Montori, Mayo Clinic, USA
Notes	NCT01502891

### **Mann 2012**

Trial name or title	Increasing efficacy of primary care-based counselling for diabetes prevention: rationale and design of the ADAPT (Avoiding Diabetes Thru Action Plan Targeting) trial
Methods	RCT
Participants	Primary care providers
Interventions	Using the ADAPT (Avoiding Diabetes Thru Action Plan Targeting) system to enhance providers' effectiveness to counsel about lifestyle behaviour changes
Outcomes	Outcome measurements are designed to detect changes in patient behaviours that are most likely to result from the use of ADAPT tool: difference between intervention and control patients in the change in mean steps per day at baseline and after 6 months, and 6 month difference of differences in haemoglobin A1C and self-reported diet between the 2 groups
Starting date	Not yet assessed
Contact information	Devin Mann, Boston University School of Medicine
Notes	Trial #: NCT01473654

### **NCT00813033**

Trial name or title	Use of a patient decision aid for gastrologic endoscopy in a paediatric setting
Methods	Interventional efficacy study
Participants	80 parents considering gastro-endoscopy for child
Interventions	Not yet assessed
Outcomes	Knowledge, expectations of outcomes, clarity of values, decision, decision conflict
Starting date	December 2008
Contact information	Nancy Neilan, Children's Mercy Hospital, Kansas City
Notes	Trials #: NCT00813033; completed March 2011

### **NCT01077037**

Trial name or title	Shared decision making in the emergency department: the Chest Pain Choice Trial
Methods	RCT
Participants	1500 adults admitted to the emergency department for chest pain, being considered by the treating clinical for admission for cardiac testing

**NCT01077037** (Continued)

Interventions	Patient decision aid vs usual care
Outcomes	Knowledge, healthcare utilization (rate of hospital admission, rate of cardiac testing, etc), patient engagement in decision-making process, decisional conflict, trust in the physician, satisfaction with decision, safety (major adverse cardiac events within 30 days)
Starting date	October 2013
Contact information	hess.erik@mayo.edu; Mayo Clinic, Rochester, Minnesota, USA
Notes	Trial #: NCT01969240

**NCT01152294**

Trial name or title	Measuring quality of decisions about treatment of menopausal symptoms
Methods	RCT
Participants	Patients talked with healthcare provider about ways to manage menopause or seriously considered taking medicine or supplement to manage menopause
Interventions	Decision aid (DVD/booklet) vs usual care
Outcomes	Knowledge, value concordance
Starting date	June 2010
Contact information	Karen R Sepucha, Massachusetts General Hospital
Notes	NCT01152294; completed, study results on <a href="http://clinicaltrials.gov">clinicaltrials.gov</a>

**NCT01152307**

Trial name or title	Measuring quality of decisions about treatment of depression
Methods	RCT
Participants	Patients that talked to a healthcare provider about starting or stopping a treatment (prescription medicine for depression or counselling)
Interventions	Decision aid (DVD/booklet) vs usual care
Outcomes	Knowledge, value concordance
Starting date	June 2010

**NCT01152307** (Continued)

Contact information	Karen R Sepucha, Massachusetts General Hospital
Notes	NCT01152307; completed, study results on clinicaltrials.gov

**NCT01447186**

Trial name or title	Informed decisions about lung cancer screening
Methods	RCT
Participants	500 adults between 55 and 77 years olds who are currently smoking or quit within the past 15 years
Interventions	Patient decision aid vs standard educational information
Outcomes	Decisional conflict: value subscale and informed subscale
Starting date	March 2015
Contact information	MD Anderson Cancer Center; USA
Notes	Trial #: NCT02286713

**NCT01618097**

Trial name or title	Evaluation of DVD and Internet decision aids for hip and knee osteoarthritis: focus on health literacy
Methods	RCT
Participants	Osteoarthritis patients
Interventions	DVD decision aid vs Internet-based decision aid
Outcomes	Decisional conflict, decision self-efficacy, knowledge
Starting date	January 2012
Contact information	Kelli D Allen, Duke University
Notes	Trial #: NCT01618097; last updated March 2014, study completion date January 2014

**NCT01713894**

Trial name or title	Utility of a clinically relevant decision aid, for parents facing extremely premature delivery
Methods	RCT
Participants	300 women who are receiving counselling at the limits of viability
Interventions	Decision aid vs usual care
Outcomes	Decisional conflict, knowledge
Starting date	May 2013
Contact information	<a href="mailto:uguillen@christianacare.org">uguillen@christianacare.org</a> ; Ursula Guillen, Christiana Care Health Systems; University of Michigan
Notes	Trial # NCT01713894

**NCT01771536**

Trial name or title	Study to test use of a decision aid in a clinical visit to help patients choose a diabetes medication. Translating Information on Comparative Effectiveness Into Practice (TRICEP)
Methods	RCT
Participants	Type 2 diabetes mellitus patients
Interventions	Diabetes medication decision aid vs usual care
Outcomes	Patient satisfaction and knowledge. Physician adoption and satisfaction with the decision aid
Starting date	January 2011
Contact information	Nilay D Shah, Mayo Clinic
Notes	NCT01293578; estimated completion date December 2014

**NCT01851785**

Trial name or title	Behavioral and social science research on understanding and reducing health disparities: African American preference for knee replacement: a patient-centred intervention (ACTION)
Methods	RCT
Participants	African-American participants referred to orthopaedic doctor with presence of knee OA
Interventions	Decision aid video + communication, skill-building intervention vs educational programme (an NIH-developed booklet) that summarizes how to live with knee OA but does not mention joint replacement

**NCT01851785** (Continued)

Outcomes	Recommendation and receipt of knee joint replacement
Starting date	July 2010
Contact information	Said A Ibrahim, University of Pennsylvania
Notes	Trial #: NCT01851785; last verified May 2013, estimated completion date June 2015

**NCT01941186**

Trial name or title	A family centered intervention to promote optimal child development
Methods	RCT
Participants	64 parent-child dyad in which the child is aged 0-36 months screening positive for developmental concern
Interventions	Patient decision aid vs usual care
Outcomes	Evaluation by early intervention specialist, attitudes, knowledge, uncertainty, intervention acceptability, intervention feasibility
Starting date	December 2013
Contact information	Children's Hospital of Philadelphia Philadelphia, PN, USA, 19104
Notes	Trial #: NCT01941186

**NCT01976325**

Trial name or title	Incorporation of the 'Ottawa Malaria Decision Aid' into the pre-travel consultation process
Methods	RCT
Participants	100 adults attending a travel clinic before travelling to an area with known chloroquine-resistant malaria
Interventions	Decision aid vs usual care
Outcomes	Knowledge, decisional conflict, preparation for decision-making, medication adherence
Starting date	January 2014
Contact information	amccarthy@toh.on.ca; Anne E McCarthy; Ottawa Hospital Research Institute
Notes	Trial # NCT01976325

**NCT02026102**

Trial name or title	A pilot trial of patient decision aids for implantable cardioverter-defibrillators (ICDs)
Methods	RCT
Participants	60 patients with heart failure referred for primary prevention implantable cardioverter-defibrillators
Interventions	Decision aid toolkit vs usual care
Outcomes	Intervention acceptability, decision quality (knowledge and values concordance), quality of life, depressive symptoms, health status, spiritual well-being
Starting date	September 2014
Contact information	amy.jenkins@ucdenver.edu; University of Colorado Hospital (UCH)
Notes	Trial #: NCT02026102

**NCT02084290**

Trial name or title	Evaluating a prediction tool and decision aid for patients with Crohn's disease
Methods	RCT
Participants	300 adults with Crohn's disease
Interventions	Patient decision aid and SDM programme vs usual care
Outcomes	Preferred choice, actual choice, adherence, cost of care, remission, patient on steroids, surgeries, Crohn's disease related hospitalizations
Starting date	March 2014
Contact information	corey.a.siegel@hitchcock.org; Corey A Siegel; Dartmouth-Hitchcock Medical Center
Notes	Trial #: NCT02084290

**NCT02110979**

Trial name or title	Validation of a patient decision aid for type 2 diabetes
Methods	RCT
Participants	200 type 2 diabetes patients
Interventions	Patient decision aid vs usual care
Outcomes	Knowledge, decisional conflict

**NCT02110979** (Continued)

Starting date	April 2014
Contact information	EPI-Q Inc, Oak Brook, IL, USA, 60523 <a href="http://www.epi-q.com/our-approach">www.epi-q.com/our-approach</a>
Notes	Trial #: NCT02110979

**NCT02145481**

Trial name or title	Decisional quality for patients with stable coronary artery disease
Methods	RCT
Participants	846 adults with stable coronary artery disease
Interventions	Patient decision aid vs standard education
Outcomes	Quality of the decision-making process, knowledge, communication, involvement, treatment preferences
Starting date	May 2014
Contact information	R. Adams Dudley; University of California, San Francisco
Notes	Trial # NCT02145481

**NCT02198690**

Trial name or title	Randomized trial of a mammography decision aid for women aged 75 and older
Methods	RCT
Participants	550 women aged 75-89 years
Interventions	Decision aid vs usual care
Outcomes	Receipt of mammography screening, acceptability, anxiety, decision-making role, decisional conflict, home safety, home safety discussions, knowledge, preparation for decision-making, screening discussions, screening intentions
Starting date	September 2014
Contact information	Mara A Schonberg, MD, MPH; <a href="mailto:mschonbe@bidmc.harvard.edu">mschonbe@bidmc.harvard.edu</a> ; Beth Israel Deaconess Medical Center; Boston, MA, USA
Notes	NCT02198690



**NCT02235571**

Trial name or title	iChoose kidney decision aid for treatment options among end-stage renal disease (ESRD) patients
Methods	RCT
Participants	450 adults with end-stage renal disease on dialysis for < 1 year and being evaluated for kidney transplant
Interventions	Patient decision aid vs usual care
Outcomes	Knowledge, evidence of shared decision-making, access to transplant, treatment preferences
Starting date	September 2014
Contact information	Rachel Patzer; Emory Transplant Center; Atlanta, GA, USA
Notes	Trial # NCT02235571

**NCT02248974**

Trial name or title	Development and user testing of a decision aid for left ventricular assist device (LVAD) placement
Methods	RCT
Participants	144 adults who are candidates for a left ventricular assist device
Interventions	Patient decision aid vs. standard education
Outcomes	Knowledge, decisional conflict, control preferences scale, CollaboRATE score, perceived quality of care, satisfaction with decision-making process, decisional regret, satisfaction with life, preparation for decision-making, usability and acceptability of the intervention
Starting date	February 2014
Contact information	Jennifer Blumenthal-Barby; Baylor College of Medicine; Houston, TX
Notes	Trial #: NCT02248974

**NCT02259699**

Trial name or title	Ovarian cancer patient-centered decision aid
Methods	RCT
Participants	221 women with stage III optimally debulked advanced ovarian cancer
Interventions	Patient decision aid vs usual care
Outcomes	Satisfaction with decision, evidence of shared decision-making, quality of life, satisfaction with care and satisfaction with cancer treatment

**NCT02259699** (Continued)

Starting date	December 2014
Contact information	lwenzel@uci.edu; Lari Wenzel; University of California, Irvine, USA
Notes	Trial #: NCT02259699

**NCT02308592**

Trial name or title	Patient decision aid for antidepressant use in pregnancy
Methods	RCT
Participants	50 women aged 18 years or older planning a pregnancy or <30 weeks pregnant
Interventions	Patient decision aid vs standard resource sheet
Outcomes	depression, anxiety, decisional conflict, knowledge, intervention acceptability, choice, satisfaction with DA
Starting date	January 2015
Contact information	<a href="mailto:simone.vigod@wchospital.ca">simone.vigod@wchospital.ca</a> Women's College Hospital, Toronto, Ontario, Canada
Notes	Trial #: NCT02308592

**NCT02319525**

Trial name or title	
Methods	
Participants	
Interventions	
Outcomes	
Starting date	
Contact information	
Notes	

**NCT02326597**

Trial name or title	Decision aid for therapeutic options in sickle cell disease
Methods	RCT
Participants	120 individuals with sickle cell disease ages 8 to 80 years
Interventions	Decision aid vs usual care
Outcomes	Knowledge, self-efficacy, decisional conflict, values, realistic expectations, preparation for decision-making, choice predisposition, stage of decision-making, decisional regret
Starting date	September 2014
Contact information	diana.ross@emory.edu; principal investigator Lakshmanan Krishnamurti; Emory University, Atlanta, GA, USA
Notes	Trial # NCT02326597

**NCT02344576**

Trial name or title	A multicenter trial of a shared decision support intervention for patients and their caregivers offered destination therapy for end-stage heart failure
Methods	RCT
Participants	400 adults advanced heart failure and are being evaluated for destination left ventricular assist device
Interventions	Patient decision aid vs usual care
Outcomes	Knowledge, values, decisional conflict, decisional regret, stress, anxiety, depression, quality of life, control preferences scale, illness acceptance, health status
Starting date	May 2015
Contact information	jocelyn.thompson@ucdenver.edu; University of Colorado, Denver
Notes	Trial #: NCT02344576

**NCT02488317**

Trial name or title	Empowering patients on choices for renal replacement therapy
Methods	RCT
Participants	150 adults with kidney disease
Interventions	Patient decision aid vs usual care

**NCT02488317** (Continued)

Outcomes	Preference for shared decision-making (CPS), decisional conflict, decision self-efficacy, knowledge, preparation for decision making
Starting date	May 2015
Contact information	Francesca Tentori; Arbor Research Collaborative for Health; Ann Arbor, MI
Notes	Trial #: NCT02488317

**NCT02488603**

Trial name or title	Utilization of decision aids for tamoxifen treatment in breast cancer patients
Methods	RCT
Participants	360 breast cancer patients referred for tamoxifen treatment
Interventions	Patient decision aid vs usual care
Outcomes	Knowledge, decisional conflict scale, satisfaction with decision, quality of life
Starting date	August 2015
Contact information	Eun Sook Lee; National Cancer Center, Korea
Notes	Trial # NCT02488603

**NCT02492009**

Trial name or title	Patient decision aid for antidepressant use in pregnancy
Methods	RCT
Participants	50 women aged 18 years or older planning a pregnancy or < 30 weeks pregnant
Interventions	Patient decision aid vs standard resource sheet
Outcomes	Depression, anxiety, decisional conflict, knowledge, intervention acceptability, choice
Starting date	June 2015
Contact information	hind.khalifeh@kcl.ac.uk or ruth.brauer@kcl.ac.uk Section of Women's Mental Health, King's College London
Notes	Trial #: NCT02492009

**NCT02503553**

Trial name or title	Decision aids in cerebral aneurysm treatment
Methods	RCT
Participants	60 patients undergoing treatment for cerebral aneurysm
Interventions	Patient decision aid vs usual care
Outcomes	Participation in the shared-decision making process; stress levels, patient satisfaction level
Starting date	August 2015
Contact information	Kimon Bekelis; Dartmouth-Hitchcock Medical Center; New Hampshire, USA
Notes	Trial #: NCT02503553

**NCT02516449**

Trial name or title	Assessment of shared decision making aids in asthma
Methods	RCT
Participants	51 adults with mild to severe asthma
Interventions	Patient decision aid vs usual care
Outcomes	Knowledge, decisional conflict, treatment adherence, asthma control
Starting date	March 2013
Contact information	Centre de recherche de l'Institut universitaire de cardiologie et de pneumologie de Québec, Québec, Canada, G1V 4G5
Notes	Trial # NCT02516449

**NCT02540044**

Trial name or title	Supporting patient care with electronic resource (SuPER): efficacy of an online decision aid for patients considering biologic therapy for rheumatoid arthritis
Methods	RCT
Participants	144 adults with rheumatoid arthritis whose rheumatologists have recommended initiating a biologic/subsequent entry biologic or switching to another biologic agent
Interventions	Online patient decision aid vs online standard information

**NCT02540044** (Continued)

Outcomes	Decisional conflict, knowledge, self-efficacy, self-management behaviours, health resource utilization, choice, evidence of shared decision-making
Starting date	January 2016
Contact information	Linda Li; University of British Columbia; Vancouver, Canada
Notes	Trial #: NCT02540044

**NCT02611050**

Trial name or title	Treatment decisions for multi-vessel CAD
Methods	RCT
Participants	160 adults with stable multi-vessel CAD at relative equipoise for at least 2 potential treatment options
Interventions	Option grid decision aid vs usual care
Outcomes	Decisional conflict, CollaboRATE score, knowledge, patient experience, treatment received
Starting date	December 2015
Contact information	Elizabeth L Nichols; the Dartmouth Institute
Notes	Trial #: NCT02611050

**Oostendorp 2011**

Trial name or title	Assessing the information desire of patients with advanced cancer by providing information with a decision aid, which is evaluated in a randomized trial: a study protocol
Methods	RCT
Participants	Patients with advanced colorectal, breast, or ovarian cancer and have started treatment with first-line palliative chemotherapy
Interventions	Patients are randomized to receive either usual care or usual care + decision aid
Outcomes	Not yet assessed
Starting date	Not yet assessed
Contact information	Linda JM Oostendorp, Radbound University
Notes	Netherlands Trial Register (NTR): NTR1113

## Yu 2015

Trial name or title	Impact of an interprofessional shared decision-making and goal setting decision aid for patients with diabetes
Methods	Cluster-randomized controlled trial
Participants	112 patients with diabetes
Interventions	Multicomponent patient decision aid toolkit vs patient education pamphlet
Outcomes	Decisional conflict, diabetes distress, health-related quality of life, chronic illness care, intention to engage in SDM
Starting date	April 2015
Contact information	yuca@smh.ca
Notes	Trial # NCT02379078

**CA-125:** cancer antigen 125; **CAD:** coronary artery disease; **CT:** computerized tomography; **NIH:** National Institutes of Health; **NSW:** New South Wales; **OA:** osteoarthritis; **RCT:** randomized controlled trial; **SDM:** shared decision making.

## DATA AND ANALYSES

### Comparison 1. Knowledge

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Knowledge - all studies	52	13316	Mean Difference (IV, Random, 95% CI)	13.27 [11.32, 15.23]
2 Knowledge - subgroup by timing of intervention (in consultation versus in preparation for consultation)	52		Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 In consultation	8	922	Mean Difference (IV, Random, 95% CI)	10.57 [4.79, 16.36]
2.2 In preparation for consultation	44	12394	Mean Difference (IV, Random, 95% CI)	13.77 [11.61, 15.93]
3 Knowledge - studies without high risk of bias	47	12327	Mean Difference (IV, Random, 95% CI)	13.43 [11.37, 15.49]

### Comparison 2. Accurate risk perceptions

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Accurate risk perceptions - all studies	17	5096	Risk Ratio (M-H, Random, 95% CI)	2.10 [1.66, 2.66]
2 Accurate risk perceptions - subgroup by timing of intervention (in consultation versus in preparation for consultation)	17		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.1 In consultation	6	898	Risk Ratio (M-H, Random, 95% CI)	1.79 [1.28, 2.52]
2.2 In preparation for consultation	11	4198	Risk Ratio (M-H, Random, 95% CI)	2.25 [1.65, 3.07]
3 Accurate risk perceptions - studies without high risk of bias	15	4732	Risk Ratio (M-H, Random, 95% CI)	2.02 [1.57, 2.59]



**Comparison 3. Informed values-choice congruence**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Informed values-choice congruence - all studies	10	4626	Risk Ratio (M-H, Random, 95% CI)	2.06 [1.46, 2.91]
2 Informed values-choice congruence - actual choice only	8	4154	Risk Ratio (M-H, Random, 95% CI)	2.13 [1.44, 3.14]
3 Informed values-choice congruence -using MMIC	8	4365	Risk Ratio (M-H, Random, 95% CI)	2.08 [1.40, 3.08]
4 Informed values-choice congruence - heterogeneous measures	2	261	Risk Ratio (M-H, Random, 95% CI)	2.02 [1.44, 2.83]
5 Informed values-choice congruence - without studies of high risk of bias	10	4626	Risk Ratio (M-H, Random, 95% CI)	2.06 [1.46, 2.91]

**Comparison 4. Decisional conflict**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Decisional conflict - all studies	42		Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Total decisional conflict score	38	8785	Mean Difference (IV, Random, 95% CI)	-7.22 [-9.12, -5.31]
1.2 Uninformed subscale	27	5707	Mean Difference (IV, Random, 95% CI)	-9.28 [-12.20, -6.36]
1.3 Unclear values subscale	23	5068	Mean Difference (IV, Random, 95% CI)	-8.81 [-11.99, -5.63]
1.4 Uncertainty subscale	28	6200	Mean Difference (IV, Random, 95% CI)	-4.04 [-6.27, -1.81]
1.5 Unsupported subscale	24	5214	Mean Difference (IV, Random, 95% CI)	-6.27 [-8.86, -3.68]
1.6 Ineffective choice subscale	24	5241	Mean Difference (IV, Random, 95% CI)	-6.31 [-8.93, -3.70]
2 Decisional conflict - in consultation	6		Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Uncertainty subscale	2	310	Mean Difference (IV, Random, 95% CI)	-6.45 [-18.29, 5.38]
2.2 Uninformed subscale	4	545	Mean Difference (IV, Random, 95% CI)	-6.37 [-14.58, 1.85]
2.3 Unclear values subscale	1	204	Mean Difference (IV, Random, 95% CI)	-17.2 [-23.77, -10.63]
2.4 Unsupported subscale	2	354	Mean Difference (IV, Random, 95% CI)	-7.16 [-13.28, -1.03]
2.5 Ineffective choice subscale	2	307	Mean Difference (IV, Random, 95% CI)	-2.37 [-7.31, 2.58]
2.6 Total decisional conflict score	5	735	Mean Difference (IV, Random, 95% CI)	-6.46 [-12.78, -0.14]
3 Decisional conflict - in preparation for consultation	36		Mean Difference (IV, Random, 95% CI)	Subtotals only
3.1 Uncertainty subscale	26	5890	Mean Difference (IV, Random, 95% CI)	-3.83 [-6.12, -1.55]
3.2 Uninformed subscale	23	5162	Mean Difference (IV, Random, 95% CI)	-9.81 [-13.00, -6.61]
3.3 Unclear values subscale	22	4864	Mean Difference (IV, Random, 95% CI)	-8.40 [-11.59, -5.21]
3.4 Unsupported subscale	22	4860	Mean Difference (IV, Random, 95% CI)	-6.18 [-8.96, -3.40]
3.5 Ineffective choice subscale	22	4934	Mean Difference (IV, Random, 95% CI)	-6.75 [-9.59, -3.90]

3.6 Total decisional conflict score	33	8050	Mean Difference (IV, Random, 95% CI)	-7.32 [-9.35, -5.28]
4 Decisional conflict - without studies having high risk of bias	39		Mean Difference (IV, Random, 95% CI)	Subtotals only
4.1 Uncertainty subscale	26	5809	Mean Difference (IV, Random, 95% CI)	-4.53 [-6.87, -2.18]
4.2 Uninformed subscale	25	5316	Mean Difference (IV, Random, 95% CI)	-9.96 [-13.13, -6.78]
4.3 Unclear values subscale	21	4677	Mean Difference (IV, Random, 95% CI)	-9.55 [-13.08, -6.02]
4.4 Unsupported subscale	22	4823	Mean Difference (IV, Random, 95% CI)	-7.00 [-9.76, -4.24]
4.5 Ineffective choice subscale	22	4850	Mean Difference (IV, Random, 95% CI)	-6.97 [-9.76, -4.18]
4.6 Total decisional conflict score	35	8240	Mean Difference (IV, Random, 95% CI)	-7.81 [-9.84, -5.77]

### Comparison 5. Participation in decision making

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Participation in decision making - all studies	16		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Clinician-controlled decision making	16	3180	Risk Ratio (M-H, Random, 95% CI)	0.68 [0.55, 0.83]
1.2 Patient-controlled decision making	15	3009	Risk Ratio (M-H, Random, 95% CI)	1.28 [1.05, 1.55]
1.3 Shared decision making	15	2973	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.83, 1.10]
2 Participation in decision making - in consultation	3		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.1 Clinician-controlled decision making - in consultation	3	650	Risk Ratio (M-H, Random, 95% CI)	0.89 [0.70, 1.12]
2.2 Patient-controlled decision making - in consultation	2	479	Risk Ratio (M-H, Random, 95% CI)	1.01 [0.80, 1.27]
2.3 Shared decision making - in consultation	2	479	Risk Ratio (M-H, Random, 95% CI)	1.14 [0.84, 1.55]
3 Participation in decision making - in preparation for consultation	13		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
3.1 Clinician-controlled decision making	13	2530	Risk Ratio (M-H, Random, 95% CI)	0.60 [0.48, 0.75]
3.2 Patient-controlled decision making	13	2530	Risk Ratio (M-H, Random, 95% CI)	1.37 [1.08, 1.73]
3.3 Shared decision making	13	2494	Risk Ratio (M-H, Random, 95% CI)	0.94 [0.80, 1.09]

**Comparison 6. Proportion undecided**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Proportion undecided - all studies	22	5256	Risk Ratio (M-H, Random, 95% CI)	0.64 [0.52, 0.79]

**Comparison 7. Satisfaction**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Satisfaction with the choice - all studies	11		Mean Difference (IV, Random, 95% CI)	Totals not selected
2 Satisfaction with the choice - in consultation	1		Mean Difference (IV, Random, 95% CI)	Totals not selected
3 Satisfaction with the choice - in preparation for consultation	10		Mean Difference (IV, Random, 95% CI)	Totals not selected
4 Satisfaction with the decision making process - all studies	9		Mean Difference (IV, Random, 95% CI)	Totals not selected
5 Satisfaction with the decision making process - in consultation	1		Mean Difference (IV, Random, 95% CI)	Totals not selected
6 Satisfaction with the decision making process - in preparation for consultation	8		Mean Difference (IV, Random, 95% CI)	Totals not selected

**Comparison 8. Choice**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Choice: surgery over conservative option	18		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Per-protocol analysis	18	3286	Risk Ratio (M-H, Random, 95% CI)	0.87 [0.75, 1.01]
1.2 Intention-to-treat analysis	18	3844	Risk Ratio (M-H, Random, 95% CI)	0.86 [0.75, 1.00]
1.3 Per-protocol analysis without prophylactic mastectomy	17	3108	Risk Ratio (M-H, Random, 95% CI)	0.84 [0.73, 0.97]
2 Choice for screening	25		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.1 PSA screening	10	3996	Risk Ratio (M-H, Random, 95% CI)	0.88 [0.80, 0.98]
2.2 Colorectal cancer screening	10	4529	Risk Ratio (M-H, Random, 95% CI)	1.12 [0.95, 1.31]
2.3 Breast cancer genetic testing	3	738	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.71, 1.38]

2.4 Prenatal diagnostic testing	2	1100	Risk Ratio (M-H, Random, 95% CI)	0.99 [0.91, 1.09]
3 Choice: diabetes medication (uptake new medication)	4	447	Risk Ratio (M-H, Random, 95% CI)	1.65 [1.06, 2.56]

## ADDITIONAL TABLES

Table 1. Decision aids evaluated in the trials

Study	Topic	Availability	Source	Contact Information
<a href="#">Allen 2010</a>	Prostate cancer screening	No	Allen, Center for Community-Based Research, Dana-Farber Cancer Institute, Boston, MA, USA, 2010	Requested access
<a href="#">Arterburn 2011</a>	Bariatric surgery	Yes	Informed Medical Decisions Foundation, MA, USA, 2010	<a href="http://informedmedicaldecisions.org/imdf/decision-aid/making-decisions-about-weight-loss-surgery/">informedmedicaldecisions.org/imdf/decision-aid/making-decisions-about-weight-loss-surgery/</a>
<a href="#">Auvinen 2004</a>	Prostate cancer treatment	Yes	Auvinen, Helsinki, Finland, 1993	Included in publication
<a href="#">Barry 1997</a>	Benign prostate disease treatment	Yes	Informed Medical Decisions Foundation, MA, USA, 2001	<a href="http://informedmedicaldecisions.org/imdf/decision-aid/treatment-options-for-benign-prostatic-hyperplasia/">informedmedicaldecisions.org/imdf/decision-aid/treatment-options-for-benign-prostatic-hyperplasia/</a>
<a href="#">Bekker 2004</a>	Prenatal screening	Yes	Bekker, Leeds, UK, 2003	Included in publication
<a href="#">Bernstein 1998</a>	Ischaemic heart disease treatment	Yes	Informed Medical Decisions Foundation, MA, USA, 2002	<a href="http://informedmedicaldecisions.org/imdf/decision-aid/treatment-choices-for-carotid-artery-disease/">informedmedicaldecisions.org/imdf/decision-aid/treatment-choices-for-carotid-artery-disease/</a>
<a href="#">Berry 2013</a>	Prostate cancer treatment	No	Berry, Phyllis F. Cantor Center, MA, USA, 2011	donna.berry@dfci.harvard.edu
<a href="#">Bjorklund 2012</a>	Antenatal Down syndrome screening	Yes	Södersjukhuset, Department of Obstetrics and Gynecology, Stockholm, Sweden	<a href="https://vimeo.com/34600615/">vimeo.com/34600615/</a>

**Table 1. Decision aids evaluated in the trials** (Continued)

Bozic 2013	Osteoarthritis of the knee or hip	No	Informed Medical Decisions Foundation and Health Dialog; USA	<a href="http://www.healthdialog.com">www.healthdialog.com</a>
Brazell 2014	Pelvic Organ Prolapse	Yes	Healthwise, USA	<a href="http://decisionaid.ohri.ca">decisionaid.ohri.ca</a>
Chabrera 2015	Prostate cancer treatment	No	C Chabrera. School of Health Sciences, Department of Nursing, Mataro, Spain	cchabrera@tecnocampus.cat
Chambers 2012	Healthcare personnel's influenza immunization	Yes	A McCarthy. Ottawa Influenza Decision Aid Planning Group, CA, 2008	<a href="http://decisionaid.ohri.ca/decaids.html#oida">decisionaid.ohri.ca/decaids.html#oida</a>
Clancy 1988	Hepatitis B Vaccine	No	Clancy, Richmond VA, USA, 1983	-
Davison 1997	Prostate cancer treatment	No	Davison, Manitoba CA, 1992-1996	-
De Achaval 2012	Total knee arthroplasty treatment	Yes	Informed Medical Decisions Foundation, MA, USA	<a href="http://informedmedicaldecisions.org/imdf/decision-aid/treatment-choices-for-knee-osteoarthritis/">informedmedicaldecisions.org/imdf/decision-aid/treatment-choices-for-knee-osteoarthritis/</a>
Dolan 2002	Colon cancer screening	No	Dolan, Rochester NY, USA, 1999	-
Evans 2010	Prostate cancer screening	Yes	Elwyn, Cardiff, UK	<a href="http://www.prosdex.com">www.prosdex.com</a>
Fagerlin 2011	Breast cancer prevention	Yes	Fagerlin, Ann Arbor, MI, USA	-
Fraenkel 2007	Osteoarthritis knee treatment	No	Fraenkel, New Haven CT, USA	Author said DA never fully developed, all info in paper
Fraenkel 2012	Atrial fibrillation	No	Veterans Affairs Connecticut Healthcare System, USA	Obtained from author <a href="mailto:terri.fried@yale.edu">terri.fried@yale.edu</a>
Frosch 2008a	Prostate cancer screening	No	Frosch, Los Angeles, USA	Screenshots from author
Gattellari 2003	Prostate cancer screening	Yes	Gatellari, Sydney, AU, 2003	included in publication

**Table 1. Decision aids evaluated in the trials** (Continued)

<a href="#">Gattellari 2005</a>	Prostate cancer screening	Yes	Gatellari, Sydney, AU, 2003	Included in publication
<a href="#">Green 2001</a>	Breast cancer genetic testing	Yes	Green, Hershey PA, USA, 2000	1-800-757-4868 dwc@mavc.com
<a href="#">Hamann 2006</a>	Schizophrenia treatment	Yes	Hamann, Munich, GER	Emailed by author (in German)
<a href="#">Hanson 2011</a>	Feeding options in advanced dementia	Yes	Mitchell, Tetroe, O'Connor; 2001 (updated 2008)	<a href="http://decisionaid.ohri.ca/decaids.html#feedingtube">decisionaid.ohri.ca/decaids.html#feedingtube</a>
<a href="#">Heller 2008</a>	Breast reconstruction	Yes	University of Texas MD Anderson Cancer Center, Houston TX, USA, 2003	Disc mailed
<a href="#">Hess 2012</a>	Stress testing for chest pain	Yes	Hess, Rochester, MN, USA, 2012	Included in publication
<a href="#">Jibaja-Weiss 2011</a>	Breast cancer treatment	Yes	Jibaja-Weiss, Baylor College of Medicine, 2010	<a href="http://www.bcm.edu/patchworkofflife">www.bcm.edu/patchworkofflife</a>
<a href="#">Johnson 2006</a>	Endodontic treatment	Yes	Johnson, Chicago, USA, 2004	Included in publication
<a href="#">Kasper 2008</a>	Multiple sclerosis	No	Jürgen Kasper	-
<a href="#">Kennedy 2002</a>	Abnormal uterine bleeding treatment	No	Kennedy/Coulter, London UK, 1996	-
<a href="#">Knops 2014</a>	Asymptomatic Abdominal Aortic Aneurysm treatment	Yes	Amsterdam, The Netherlands	<a href="http://www.keuzehulp.info/amc/AAA/landing-page">www.keuzehulp.info/amc/AAA/landing-page</a>
<a href="#">Krist 2007</a>	Prostate cancer screening	Yes	Krist, Fairfax VA, USA	<a href="http://www.familymedicine.vcu.edu/research/misc/psa/index.html">www.familymedicine.vcu.edu/research/misc/psa/index.html</a>
<a href="#">Kupke 2013</a>	Dental - posterior tooth decay	Yes	University of Cologne, Cologne, Germany	jana.kupke@uk-koeln.de
<a href="#">Kuppermann 2014</a>	Prenatal screening	No	Kuppermann, San Francisco CA, USA	Interactive web-based decision aid
<a href="#">Lam 2013</a>	Breast cancer treatment	Yes	Kwong Wah Hospital, Hong Kong, China	Obtained from author. wwtlam@hku.hk

**Table 1. Decision aids evaluated in the trials** (Continued)

Langston 2010	Contraceptive method choice	Yes	World Health Organization, 2005	<a href="http://www.who.int/reproductivehealth/publications/family_planning/9241593229index/en/index.html">www.who.int/reproductivehealth/publications/family_planning/9241593229index/en/index.html</a>
Laupacis 2006	Pre-operative autologous blood donation	No	Laupacis, Ottawa, CA, 2001	<a href="http://Decisionaid.ohri.ca/decaids-archive.html">Decisionaid.ohri.ca/decaids-archive.html</a>
LeBlanc 2015	Treatment for osteoporosis	Yes	Mayo Clinic	-
Legare 2008a	Natural health products	No	Legare, Quebec City, CA, 2006	-
Legare 2011	Use of antibiotics for acute respiratory infections	Yes	Legare, Quebec City, CA, 2007	<a href="http://www.decision.chaire.fmed.ulaval.ca/index.php?id=192&amp;L=2">www.decision.chaire.fmed.ulaval.ca/index.php?id=192&amp;L=2</a>
Legare 2012	Antibiotics for acute respiratory infections	Yes	Legare, Quebec City, CA	<a href="http://www.decision.chaire.fmed.ulaval.ca/index.php?">www.decision.chaire.fmed.ulaval.ca/index.php?</a>
Leighl 2011	Advanced colorectal cancer chemotherapy	Yes	Princess Margaret Hospital, Toronto, 2011	Natasha.Leighl@uhn.on.ca
Lepore 2012	Prostate cancer screening	Yes	Sally Weinrich University of Louisville, USA	Obtained from author slepore@temple.edu
Lerman 1997	Breast cancer genetic testing	No	Lerman/Schwartz, Washington DC, USA, 1997	-
Lewis 2010	Colorectal cancer screening	Yes	Lewis, University of North Carolina, Chapel Hill, NC, USA, 2010	<a href="http://decisionsupport.unc.edu/CHOICE6/">decisionsupport.unc.edu/CHOICE6/</a>
Loh 2007	Depression treatment	Yes	Loh, Freiburg, GER	Emailed to us by author - in German
Man-Son-Hing 1999	Atrial fibrillation treatment	No	McAlister/Laupacis, Ottawa CA, 2000	<a href="http://decisionaid.ohri.ca/decaids-archive.html">decisionaid.ohri.ca/decaids-archive.html</a>
Mann D 2010	Diabetes treatment - statins	Yes	Montori, Rochester MN, USA	<a href="http://mayoresearch.mayo.edu/mayo/research/ker/unit/form.cfm">mayoresearch.mayo.edu/mayo/research/ker/unit/form.cfm</a>

**Table 1. Decision aids evaluated in the trials** (Continued)

<a href="#">Mann E 2010</a>	Diabetes screening	Yes	Marteau, King's College London, London, England, 2010	Additional file 2 of publication
<a href="#">Marteau 2010</a>	Diabetes screening	Yes	Marteau, King's College London, London, England, 2010	Provided by author, same DA as <a href="#">Mann E 2010</a>
<a href="#">Mathieu 2007</a>	Mammography	Yes	Mathieu, Sydney, AU	DA emailed by author
<a href="#">Mathers 2012</a>	Diabetes treatment	Yes	The University of Sheffield, Sheffield, UK, 2008	Obtained from author C.Ng@sheffield.ac.uk
<a href="#">Mathieu 2010</a>	Mammography	Yes	Mathieu, University of Sydney, AUS, 2010	<a href="http://www.psych.usyd.edu.au/comped/com'decision'aids.shtml">www.psych.usyd.edu.au/comped/com'decision'aids.shtml</a>
<a href="#">McAlister 2005</a>	Atrial fibrillation treatment	No	McAlister/Laupacis, Ottawa CAN, 2000	<a href="http://decisionaid.ohri.ca/decaids-archive.html">decisionaid.ohri.ca/decaids-archive.html</a>
<a href="#">McBride 2002</a>	Hormone replacement therapy	Yes, update in progress	Sigler/Bastien, Durham NC, USA, 1998	basti001@mc.duke.edu
<a href="#">McCaffery 2010</a>	Screening after mildly abnormal pap smear	Yes	Screening & test evaluation program, School of public health, University of Sydney 2007	kirstenm@health.usyd.edu.au
<a href="#">Miller 2005</a>	BRCA1/BRCA2 gene testing	No	Miller, Fox Chase PA, USA	-
<a href="#">Miller 2011</a>	Colorectal cancer screening	Yes	University of North Carolina, Chapel Hill, NC, USA, 2007	<a href="http://intmedweb.wakehealth.edu/choice/choice.html">intmedweb.wakehealth.edu/choice/choice.html</a> (no longer available)
<a href="#">Montgomery 2003</a>	Hypertension treatment	No	Montgomery, UK, 2000	-
<a href="#">Montgomery 2007</a>	Birthing options after caesarean	Yes	Montgomery, Bristol, UK, last update 2004	<a href="http://www.computing.dundee.ac.uk/acstaff/cjones/diamond/Information.html">www.computing.dundee.ac.uk/acstaff/cjones/diamond/Information.html</a>
<a href="#">Montori 2011</a>	Osteoporosis treatment	Yes	Montori, Mayo Foundation for Medical Education and Research, 2007	<a href="http://shareddecisions.mayoclinic.org/decision-aids-for-diabetes/other-decision-">shareddecisions.mayoclinic.org/decision-aids-for-diabetes/other-decision-</a>



**Table 1. Decision aids evaluated in the trials** (Continued)

				aids/
<a href="#">Morgan 2000</a>	Ischaemic heart disease treatment	Yes	Informed Medical Decisions Foundation, MA, USA, 2002	<a href="http://informedmedicaldecisions.org/imdf/decision%20aid/treatment-choices-for-carotid-artery-disease/">informedmedicaldecisions.org/imdf/decision%20aid/treatment-choices-for-carotid-artery-disease/</a>
<a href="#">Mott 2014</a>	PTSD treatment	Yes	Michael E DeBakey Veterans Affairs Medical Center, Houston, USA	Obtained from author <a href="mailto:juliette.mott@va.gov">juliette.mott@va.gov</a>
<a href="#">Mullan 2009</a>	Diabetes treatment	Yes	Montori or Mayo Foundation(?) Rochester MN, USA,	Included in publication
<a href="#">Murray 2001a</a>	Benign prostate disease treatment	Yes	Informed Medical Decisions Foundation, MA, USA, 2001	<a href="http://informedmedicaldecisions.org/imdf/decision%20aid/treatment-options-for-benign-prostatic-hyperplasia/">informedmedicaldecisions.org/imdf/decision%20aid/treatment-options-for-benign-prostatic-hyperplasia/</a>
<a href="#">Murray 2001b</a>	Hormone replacement therapy	No, update in progress	Informed Medical Decisions Foundation, MA, USA	<a href="http://informedmedicaldecisions.org/imdf/decision%20aid/treatment-choices-for-managing-menopause/">informedmedicaldecisions.org/imdf/decision%20aid/treatment-choices-for-managing-menopause/</a>
<a href="#">Nagle 2008</a>	Prenatal screening	Yes	Nagle, Victoria, AU	<a href="http://www.mcrci.edu.au/Downloads/PrenatalTestingDecisionAid.pdf">www.mcrci.edu.au/Downloads/PrenatalTestingDecisionAid.pdf</a>
<a href="#">Nassar 2007</a>	Birth breech presentation	Yes	Nassar, West Perth WA, AU	<a href="http://sydney.edu.au/medicine/public-health/shdg/resources/decision%20aids.php">sydney.edu.au/medicine/public-health/shdg/resources/decision%20aids.php</a>
<a href="#">Oakley 2006</a>	Osteoporosis treatment	No	Cranney, Ottawa CA, 2002	<a href="http://decisionaid.ohri.ca/decaids-archive.html">decisionaid.ohri.ca/decaids-archive.html</a>
<a href="#">Ozanne 2007</a>	Breast cancer prevention	No	Ozanne, Boston MA, USA	-
<a href="#">Partin 2004</a>	Prostate cancer screening	Yes	Informed Medical Decisions Foundation, MA, USA, 2001	<a href="http://informedmedicaldecisions.org/imdf/decision%20aid/deciding-if-the-psa-test-is-right-for-you/">informedmedicaldecisions.org/imdf/decision%20aid/deciding-if-the-psa-test-is-right-for-you/</a>

**Table 1. Decision aids evaluated in the trials** (Continued)

<a href="#">Pignone 2000</a>	Colon cancer screening	Yes	Pignone, Chapel Hill NC, USA, 1999	<a href="http://www.med.unc.edu/medicine/edusrc/colon.htm">www.med.unc.edu/medicine/edusrc/colon.htm</a>
<a href="#">Protheroe 2007</a>	Menorrhagia treatment	No	Protheroe, Manchester, UK	Computerized decision aid, Clinical Guidance Tree - no longer in existence, author sent chapter in thesis
<a href="#">Rubel 2010</a>	Prostate cancer screening	No	Centers for Disease Control and Prevention (CDC), USA, 2010	No longer available
<a href="#">Ruffin 2007</a>	Colorectal cancer screening	Yes	Regents of the University of Michigan (copyright info), Ann Arbor MI, USA, 2006	<a href="http://colorectalweb.org">colorectalweb.org</a>
<a href="#">Sawka 2012</a>	Adjuvant radioactive iodine treatment for patients with early-stage papillary thyroid cancer	No	University Health Network, Toronto, Canada, 2009	-
<a href="#">Schroy 2011</a>	Colorectal cancer screening	Yes	Schroy III, Boston, USA	<a href="mailto:Paul.schroy@bmc.org">Paul.schroy@bmc.org</a>
<a href="#">Schwalm 2012</a>	Coronary angiogram access site	Yes	Schwalm, Hamilton, ON, Canada, 2009	<a href="http://www.phri.ca/workfiles/studies/presentations/PtDA%20Vascular%20Access%202023-May-2012.pdf">www.phri.ca/workfiles/studies/presentations/PtDA%20Vascular%20Access%202023-May-2012.pdf</a>
<a href="#">Schwartz 2001</a>	Breast cancer genetic testing	No	Schwartz/Lerman, Washington DC, USA, 1997	-
<a href="#">Schwartz 2009a</a>	BRCA mutation prophylactic surgery	No	Schwartz, Washington DC, USA	-
<a href="#">Sheridan 2006</a>	Cardiovascular prevention	Yes	Sheridan, Chapel Hill, NC, USA	<a href="http://www.med-decisions.com/cvtool/">www.med-decisions.com/cvtool/</a>
<a href="#">Sheridan 2011</a>	Coronary heart disease prevention	Yes	Sheridan, University of North Carolina at Chapel Hill, Division of General Internal Medicine, North Carolina, USA, 2011	<a href="http://www.med-decisions.com/h2hv3/">www.med-decisions.com/h2hv3/</a>

**Table 1. Decision aids evaluated in the trials** (Continued)

<a href="#">Shorten 2005</a>	Birth options after previous caesarean	Yes (updated 2006)	Shorten, Wollongong, AU, 2000	<a href="mailto:ashorten@uow.edu.au">ashorten@uow.edu.au</a> or <a href="http://www.capersbookstore.com.au/product.asp?id=301">www.capersbookstore.com.au/product.asp?id=301</a>
<a href="#">Shourie 2013</a>	Measles mumps and rubella vaccination	Yes	University of Leeds, UK & NSIRS Australia	<a href="http://www.leedsmmr.co.uk">www.leedsmmr.co.uk</a>
<a href="#">Smith 2010</a>	Bowel cancer screening	Yes	Smith, Sydney, AU 2008	<a href="http://sydney.edu.au/medicine/public-health/shdg/resources/decision_aids.php">sydney.edu.au/medicine/public-health/shdg/resources/decision_aids.php</a>
<a href="#">Stacey 2014a</a>	Osteoarthritis of the hip and knee	No	Informed Medical Decisions Foundation and Health Dialog; USA	<a href="http://www.healthdialog.com">www.healthdialog.com</a>
<a href="#">Steckelberg 2011</a>	Colorectal cancer screening	Yes	Steckelberg, Hamburg, Germany	-
<a href="#">Taylor 2006</a>	Prostate cancer screening	Yes	Georgetown University Medical Center, Washington DC, USA, 2000	Obtained from author <a href="mailto:taylorkl@georgetown.edu">taylorkl@georgetown.edu</a>
<a href="#">Thomson 2007</a>	Atrial fibrillation treatment	Yes	Thomson, Newcastle Upon Tyne, UK	Disc sent by mail
<a href="#">Trevena 2008</a>	Colorectal cancer screen	Yes	Trevena, Sydney, AU	<a href="http://sydney.edu.au/medicine/public-health/shdg/resources/decision_aids.php">sydney.edu.au/medicine/public-health/shdg/resources/decision_aids.php</a>
<a href="#">Van Peperstraten 2010</a>	Embryos transplant	Yes	Radboud University Nijmegen Medical Centre; 2006	<a href="http://www.umcn.nl/ivfda-en">www.umcn.nl/ivfda-en</a>
<a href="#">Vandemheen 2009</a>	Cystic Fibrosis referral transplant	Yes	Aaron, Ottawa ON, CA, 2009 (last update 2011)	<a href="http://decisionaid.ohri.ca/decaids.html#cfda">decisionaid.ohri.ca/decaids.html#cfda</a>
<a href="#">Vodermaier 2009</a>	Breast cancer surgery	Yes	Vodermaier, Vancouver BC, CA	Received by email (in German)
<a href="#">Volk 1999</a>	Prostate cancer screening	Yes	Informed Medical Decisions Foundation, MA, USA, 1999	<a href="http://informedmedicaldecisions.org/imdf/decision_aid/deciding-if-the-psa-test-is-right-for-you/">informedmedicaldecisions.org/imdf/decision_aid/deciding-if-the-psa-test-is-right-for-you/</a>

**Table 1. Decision aids evaluated in the trials** (Continued)

<a href="#">Vuorma 2003</a>	Menorrhagia treatment	No	Vuorma, Helsinki Finland, 1996	-
<a href="#">Watson 2006</a>	Prostate cancer screening	Yes	Oxford, UK	Included in publication
<a href="#">Weymiller 2007</a>	Diabetes mellitus type 2 treatment	Yes	Montori, Rochester MN, USA	<a href="http://mayoresearch.mayo.edu/mayo/research/kerunit/form.cfm">mayoresearch.mayo.edu/mayo/research/kerunit/form.cfm</a>
<a href="#">Williams 2013</a>	Prostate cancer screening	Yes	Georgetown University, Washington, DC, USA	Obtained from author <a href="mailto:taylorlkl@georgetown.edu">taylorlkl@georgetown.edu</a>
<a href="#">Whelan 2003</a>	Breast cancer chemotherapy	Yes	Whelan, Hamilton CA, 1995	Included in publication
<a href="#">Whelan 2004</a>	Breast cancer surgery	Yes	Whelan, Hamilton CA, 1997	Included in publication
<a href="#">Wolf 1996</a>	Prostate cancer screening	Yes	Wolf, Charlottesville VA, USA, 1996	Script in publication
<a href="#">Wolf 2000</a>	Colon cancer screening	Yes	Wolf, Charlottesville VA, USA, 2000	Script in publication
<a href="#">Wong 2006</a>	Pregnancy termination	No	Bekker, Leeds, UK, 2002	-

**Table 2. Knowledge**

Study	Scale used	Timing	N decision aid	Decision aid - mean	N comparison	Comparison - mean	Notes
<a href="#">Bozic 2013</a>	Decision quality instrument, 19 items re knowledge (> 50%)	After 1st consultation with surgeon	60	58.3%	60	33.3%	P = 0.01
<a href="#">Evans 2010</a>	12 true or false questions; scores ranging from -12 to 12	Immediately post	89	4.9	103	2.17	P < 0.001
<a href="#">Fagerlin 2011</a>	Insufficient (≤ 50% correct)	Immediately post	383	31.8%	102	93.1%	P < 0.001

**Table 2. Knowledge** (Continued)

	Sufficient	Immediately post	383	61.9%	102	6.9%	-
Fraenkel 2012	Open-ended questions about medication options to reduce stroke - knows medications	Postintervention	66	61%	62	31%	OR 3.5 (95% CI: 1.6 to 7.7, P = 0.001)
	Open-ended questions about side effects of medications - knows side effects	Postintervention	53	49%	46	37%	OR 1.9 (95%CI: 0.9 to 4.0; P = 0.07)
Hamann 2006	7-item multiple choice knowledge test (unable to standardize results)	On discharge (~ 1 month)	49	15 (4.4 SD)	58	10.9 (5.4 SD)	P = 0.01
Heller 2008	12-item multiple choice	Pre-operatively	66	14%*	67	8%*	*mean increase from baseline P = 0.02
LeBlanc 2015 (in consultation)	13-item questionnaire (median, IQR) total score	Immediately post	32	7 (4.5 to 9.0)	45	5.5 (2.5 to 8.0)	P = 0.11
	9-items knowledge based on decision aid	Immediately post	32	6 (3.5 to 6.5)	45	4 (2.0 to 8.0)	P = 0.01
Legare 2008a	10-item yes/no/unsure general knowledge test about natural health products (not	Change scores from baseline to 2 weeks	43	0.86 ± 1.77 P = 0.002	41	0.51 ± 1.47 P = 0.031	No difference between groups (P = 0.162)

**Table 2. Knowledge** (Continued)

	specific to outcomes of options)						
Mann D 2010 (in consultation)	14-item survey	Immediately post	-	-	-	-	No difference in level of knowledge between groups
Mathers 2012	Correctly answers question about best option to lower blood sugar	6 months postintervention	95	51.6%	80	28.8%	P < 0.001
	Correctly answers question about best option to lower complications	6 months postintervention	95	31.0%	80	29%	P = 0.90
Mathieu 2007	9-item - 4 concept questions and 5 numeric questions	-	351	-	357	-	Significantly higher mean increase for the intervention group (2.62 ) compared to control group (0.68) from baseline, P < 0.001
Miller 2005	8-item survey	2-week, 2-month, and 6-month follow-ups	-	-	-	-	Intervention type had no impact on general or specific knowledge
Nagle 2008	Good level knowledge was scored higher than the mid point of the knowledge scale (greater than 4)	-	-	-	-	-	88% (147/167) in DA group compared to 72% (123/171) pamphlet group. OR 3.43 (95% CI 1.79 to 6.58)

**Table 2. Knowledge** (Continued)

<a href="#">Ozanne 2007</a> (in consultation)	Change in knowledge from baseline	Post-test	15	48% to 64%	15	45% to 57%	change in knowledge score was significant for decision aid ( $P = 0.01$ ) but not control ( $P = 0.13$ )
<a href="#">Partin 2004</a>	10-item knowledge index score	2 weeks	308	7.44	290	6.9	$P = 0.001$
<a href="#">Rubel 2010</a>	24-items adapted from existing prostate cancer knowledge measures	Immediately post	100	-	100	-	The total mean standardized knowledge score was 84.38 (SD 12.38)
<a href="#">Trevena 2008</a>	Adequate knowledge (positive score: understanding benefits/harms)	1 month	134	28/134	137	8/137	$P = 0.0001$
<a href="#">Watson 2006</a>	12-item true/false/don't know	Post-test	468	75% (range 0 to 100)	522	25% (range 0 to 100)	$P < 0.0001$
<a href="#">Weymiller 2007</a> (in consultation)	14-item - 9 addressed by decision aid; 5 were not	Immediately post	52		46	-	Mean difference between groups 2.4 (95% CI 1.5 to 3.3) $P < 0.05$ (when decision aid administered during the consultation only - not if prior to the consultation)

CI: confidence interval; DA: decision aid; OR: odds ratio; SD: standard deviation.

**Table 3. Accurate risk perceptions**

Study	Scale used	Timing	N decision aid	Decision aid - mean	N comparison	Comparison - mean	Notes
Fraenkel 2012	Accuracy of stroke risk (reported by taking the absolute value of the difference between the participant's risk as estimated by the DA and the estimate provided by the participant - out of 100; lower score indicates more accurate estimation of risk)	Postintervention	69	9.1 (SD 13.3)	66	14.2 (SD 13)	P = 0.002
	Accuracy of bleeding risk (reported same as above)	Postintervention	69	8.7 (SD 12.5)	66	13.1 (SD 12.2)	P = 0.004
Hanson 2011	Expectation of benefit index 11 items score from 1 to 4 with lower score indicating better knowledge	Post (after reviewing DA)	127	2.3	129	2.6	P = 0.001
Kuppermann 2014	Correct estimate of amniocentesis miscarriage risk	3-6 months postintervention	357	263 (73.8%)	353	208 (59.0%)	P < 0.001
	Correct estimate of Down syndrome risk	3-6 months postintervention	357	210 (58.7%)	353	163 (46.1%)	P = 0.001



**Table 3. Accurate risk perceptions** (Continued)

Mann E 2010	3 of 8 multiple choice items in the knowledge test (question 4, 5, 7)	2 weeks post	-	-	-	-	Total knowledge reported only
Mathieu 2010	5 item numerical questions (max = 5)	Post	113	3.02	189	2.45	P < 0.001
Miller 2005	-	2-week, 2-month, and 6-month follow-ups	-	-	-	-	Intervention type had no impact on risk perceptions
Smith 2010	8 numerical questions (max = 8)	-	357	2.93 (SD 2.91)	173	0.58 (SD 1.28)	P < 0.001
Weymiller 2007 (in consultation)	-	Immediately	52	-	46	-	Difference between group OR 22.4 (95% CI 5.9 to 85.8) when decision aid administered during the consultation only (not if prior to) OR 6.7 (95% CI 2.2 to 19.7) when the decision aid administered prior to or during the consultation

CI: confidence interval; DA: decision aid; OR: odds ratio; SD: standard deviation.

**Table 4. Values congruent with chosen option**

Study	Scale used	Timing	N decision aid	Decision aid - mean	N comparison	Comparison - mean	Notes
<a href="#">Arterburn 2011</a>	Percent match procedures described by Sepucha et al (2007; 2008). For values items were most predictive and used to specify logistic models to estimate predicted probability of selecting surgery > 0.5	Postintervention	75	-	77	-	The intervention group experienced a more rapid early improvement in value concordance immediately after the intervention compared to control
<a href="#">Berry 2013</a>	Concordant when men reported: a) sexual function influenced decision and they had radiation therapy; b) bowel function influenced decision and they had surgery; c) all effects influenced decision and they had surveillance	6 months postintervention	239	-	209	-	No difference OR = 0.82; 95% CI 0.56 to 1.2
<a href="#">Frosch 2008a</a>	Concordance between participant's preferences and values for potential outcomes related to the decision and the choice	within weeks	155	-	151	-	Men assigned to the decision aid who chose not to have a PSA test rated their concern about prostate cancer

**Table 4. Values congruent with chosen option** (Continued)

	made						lower than did men who requested a PSA test. Men assigned to usual care provided similar ratings of concern about prostate cancer regardless of their PSA decision. There was no statistically significant difference between groups
Legare 2008a	-	-	-	-	-	-	Women valuing of non-chemical aspect of natural health products was positively associated with their choice of nature health products, $P = 0.006$ . No difference between groups
Lerman 1997	Association between values and choice	-	-	-	-	-	No difference; between-group differences were not reported
Vandemheen 2009	Congruence between personal values and decision	3 weeks	70	-	70	-	Patient choices were consistent with their values across both randomized groups

DA: decision aid; SD: standard deviation.

**Table 5. Decisional Conflict Score**

Study	Scale used	Timing	N decision aid	Decision aid - mean	N comparison	Comparison - mean	Notes
Arterburn 2011	Total decisional conflict- change from baseline (standardised values)	Immediately post	75	Mean -20 SD 19.44	77	Mean -11.8 SD 22.83	P = 0.03
Berry 2013	Decisional conflict scale	Uncertainty	-	-3.61 units	-	-	P = 0.04
		Uninformed	-	-	-	-	No significant difference
		Unclear values	-	-3.57 units	-	-	P = 0.002
		Unsupported	-	-	-	-	No significant difference
		Ineffective decision	-	-	-	-	No significant difference
		Total	-	-1.75 units	-	-	P = 0.07
Fagerlin 2011	Decisional conflict scale	Immediately post	-	-	-	-	DCS was higher in the intervention group compared to control, P < 0.001
Frosch 2008a	Decisional conflict - subscales only	Feeling uninformed	155	23.37	151	29.68	P < 0.05
		Feeling unclear values	155	32.25	151	37.93	P < 0.05
		Feeling supported	155	30.51	151	35.21	P < 0.05
		Feeling uncertain	155	-	151	-	No difference
		Effective decisions	155	-	151	-	No difference

**Table 5. Decisional Conflict Score** (Continued)

Knops 2014	Decisional conflict (total score)	4 months	73	19 SD 14	81	22 SD 17	No difference
		10 months	73	21 SD 17	81	18 SD 17	No difference
Krist 2007	Decisional conflict	Immediately after office visit	196	1.54	75	1.58	No difference
LeBlanc 2015 (in consult)	Decision conflict (overall) median, IQR	Immediately post	28	10.9 (95% CI 1.6 to 26.6)	36	22.7 (95% CI 7.8 to 28.5)	P = 0.18
	Informed subscale	Immediately post	28	4.2 (95% CI 0 to 25)	36	20.8 (95% CI 0 to 33.3)	P = 0.14
	Values subscale	Immediately post	28	16.7 (95% CI 0 to 25)	36	25.0 (95% CI 8.3 to 33.3)	P = 0.25
	Support subscale	Immediately post	28	8.3 (95% CI 0 to 25)	36	16.7 (95% CI 0 to 25)	P = 0.35
	Certainty subscale	Immediately post	28	8.3 (95% CI 0 to 25)	36	25 (95% CI 0 to 25)	P = 0.3
	Effectiveness subscale	Immediately post	28	12.5 (95% CI 0 to 25)	36	18.8 (95% CI 0 to 25)	P = 0.15
Legare 2012 (in consult)	Decisional conflict - proportion who had a value of 2.5 or more on the 1–5 DCS. (n, %)	Immediately post	163	4.6% (95% CI 2.6 to 7.4)	165	6.3% (95% CI 0 to 12.8)	Absolute difference 1.7; RR 0.8 (95% CI 0.2 to 2.4)
Leighl 2011	Decisional conflict scale median (range)	1-2 weeks postintervention	107	26 (range 0-79)	100	26 (range 0-67)	No difference
Mathieu 2010	Based on approaches suggested by Marteau et al. (informed choice)	Immediately after intervention	91	71%	110	64%	P = 0.24

**Table 5. Decisional Conflict Score** (Continued)

<a href="#">Ozanne 2007</a> (in consult)	Decisional conflict	Postconsultation	15	-	15	-	Both groups showed lower decisional conflict postconsultation ( $P < 0.001$ ) but no difference between groups
<a href="#">Rubel 2010</a>	Decisional conflict	Immediately post	-	-	-	-	The total mean score was 24.5 with a SD of 15.25 ( $N = 200$ )
<a href="#">Schwartz 2009a</a>	Decisional conflict	12 of 16 items of the original scale	-	-	-	-	Significant longitudinal impact of the decision aid was moderated by baseline decision status; decision aid led to significant decreases in decisional conflict for those who were undecided at the time of randomisation
<a href="#">Thomson 2007</a> (in consult)	Decisional conflict	Postconsultation	53	-	56	-	Difference between decision aid and control group were $-0.18$ (95% CI $-0.34$ to $-0.01$ ). $P = 0.036$
		3-months post	51	-	55	-	Difference between decision aid and control group

**Table 5. Decisional Conflict Score** (Continued)

							were -0.15 (95% CI -0.37 to 0.06), no significant difference
Van Peperstraten 2010	15 item questionnaire (1-5) - satisfaction-uncertainty	Postintervention, pre IVF	124	72.5	128	75	P = 0.76
	15 item questionnaire (1-5) - informed (includes some items from DCS)	Postintervention, pre IVF	124	77.5	128	87.5	P = 0.001
Weymiller 2007 (in consult)	Decisional conflict	Immediately post	52	-	46	-	Mean difference indicates statistically significantly lower decisional conflict for decision aid compared to usual care Total DCS -10.6 (95% CI -15.4 to -5.9) Uncertain -12.8 (95% CI -18.4 to -7.3) Informed -17.3 (95% CI -22.6 to -12.0) if administered during consult -6.6 (95% CI -14.3 to 1.1) if adminis-

**Table 5. Decisional Conflict Score** (Continued)

							tered prior to consult Values clarity −8.5 (95% CI −15.7 to −1.3) Support −9.4 (95% CI −14.8 to −3.9) Effective decision −10.0 (95% CI −15.0 to −5.0)
--	--	--	--	--	--	--	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------

CI: confidence interval; DA: decision aid; DCS: decisional conflict scale; IVF: in vitro fertilisation; SD: standard deviation.

**Table 6. Decisional Conflict Score - low literacy version**

Study	Scale used	Timing	N decision aid	Decision aid - mean	N comparison	Comparison - mean	Notes
Fraenkel 2012	Informed	Immediately post	69	13.0	66	24.8	P = 0.01
	Values	Immediately post	69	6.4	66	21.0	P <.001
Smith 2010	Total DCS	2 week follow-up	357	13.63 (SD 20.55)	173	14.91 (SD 18.34)	P = 0.02
Taylor 2006	Total DCS	Used 8 of 10 items only 1 month post	80	24.1% high	74	41.9% high	Results were dichotomized (items removed choosing without pressure from others; know what options are available to you)
Williams 2013	Total DCS	2 months post	153	27.5%	136	38.2%	Significant decrease for DA group compared to usual care in the home condition site



**Table 6. Decisional Conflict Score - low literacy version** (Continued)

		13 months post	153	38.6%	136	31.6%	No difference
--	--	----------------	-----	-------	-----	-------	---------------

DA: decision aid; DCS: decisional conflict scale; SD: standard deviation.

**Table 7. Decisional Conflict Score - SURE test**

Study	Scale used	Timing	N decision aid	Decision aid - mean	N comparison	Comparison - mean	Notes
Stacey 2014a	SURE tool Item: 'Feels sure about the best choice'	Postintervention; prior to surgical consult	65	72.3%	66	80.3%	No difference
	'Knows the benefits and harms . . .'	Postintervention; prior to surgical consult	65	92.3%	66	66.7%	No difference
	'Clear about which benefits and harms . . .'	Postintervention; prior to surgical consult	65	87.7%	66	74.2%	No difference
	'Has enough support and advice . . .'	Postintervention; prior to surgical consult	65	76.9%	66	77.3%	No difference
	Total SURE score	Postintervention; prior to surgical consult	65	69.2%	66	57.6%	No difference

**Table 8. Patient-clinician communication**

Study	Scale used	Timing	N decision aid	Decision aid - mean	N comparison	Comparison - mean	Notes
Fraenkel 2012	Discussed risk of stroke	Immediately post	69	71%	66	12%	P < 0.001
	Discussed risk of major bleeding	Immediately post	69	69%	66	20%	P < 0.001

**Table 8. Patient-clinician communication** (Continued)

Hanson 2011	Discussed feeding with physician, nurse clinician, or physician's assistant	3 months	126	46%	127	33%	P = 0.04
	Discussed feeding with other nursing home staff	3 months	126	64%	127	71%	P = 0.42
Hess 2012 (in consult)	OPTION scale	Analysis of the consultation using video-recordings	101	Mean 26.6% (95% CI 24.9 to 8.2)	103	Mean 7% (95% CI 5.9 to 8.1)	Significantly greater in the intervention arm
LeBlanc 2015 (in consult)	OPTION scale	Analysis of the consultation using video-recordings	25	Mean 57% (95% CI 50 to 64)	13	Mean 43% (95% CI 37 to 48)	P = 0.001
Lepore 2012	Discussed PSA testing with physician postintervention	8 months postintervention	215	15.8%	216	8.3%	P < 0.001
Montori 2011 (in consult)	OPTION 100-point scale	Analysis of the consultation using video-recorded consultations	38	49.8	32	27.3	P < 0.001
Mullan 2009 (in consult)	OPTION scale	Analysis of the consultation using video-recorded consultations	48 used decision aid within consultation	Mean 49.7% (SD 17.74)	37 usual care	Mean 27.7% (SD 11.75)	MD 21.8 (95% CI 13.0 to 30.5) for decision aid vs usual care. All but 2 of the 12 items significantly favoured the decision aid
Sheridan 2006	Discussed CHD with doctor	Patient reported Immediately post	16/41 decision aid pre-consult	-	8/34 usual care	-	Absolute difference 16% (95% CI

**Table 8. Patient-clinician communication** (Continued)

			with summary report to bring to consult				–4 to 37)
	Plan to reduce CHD risk and discussed with doctor	Patient reported Immediately post	15/41 decision aid pre-consult with summary report to bring to consult	-	8/34 usual care	-	Abso-lute difference 13% (95% CI –7 to 34).
	Plan to reduce CHD risk and not discussed with doctor	Patient reported Immediately post	37/41 decision aid pre-consult with summary report to bring to consult	-	25/34 usual care	-	Abso-lute difference 16% (95% CI –1 to 33)
Sheridan 2011	Had CHD discussion with provider	Patient reported Immediately post	79	89%	78	58%	Abso-lute difference 31% (95% CI 15 to 45; P < 0.001)
	Patient-raised discussion	Patient reported Immediately post	79	63%	78	35%	Abso-lute difference 28% (95% CI 9 to 45; P = 0.02)
	Mod-ified Health-care Climate Question-naire: 1. “My provider provided me with choices and options about lowering my chances of heart disease”	Patient reported Immediately post	79	91%	78	76%	Abso-lute difference 15% (95% CI –0.1 to 31; P = 0.02)
	2. “My provider under-stands how I see things with respect to low-	Patient reported Immediately post	79	95%	78	86%	Abso-lute difference 9% (95% CI –7 to 25; P = 0.21)

**Table 8. Patient-clinician communication** (Continued)

	ering my chances of heart disease.”						
	3. “My provider conveyed confidence in my ability to make changes regarding lowering my chances of heart disease”	patient reported Immediately post	79	88%	78	77%	Abso-lute difference 11% (95% CI –5 to 27; P = 0.15)
	4. “My provider encouraged me to ask questions”	Patient reported Immediately post	79	78%	78	67%	Abso-lute difference 11% (95% CI –4% to 27%; P = 0.13)
	5. “My provider listened to how I would like to do things”	Patient reported Immediately post	79	92%	78	71%	Abso-lute difference 21% (CI 95% 6 to 37; P < 0.01)
	6. “My provider tried to understanding how I see things before suggesting new ways to lower my chances of heart disease.”	Patient reported Immediately post	79	84%	78	69%	Abso-lute difference 15% (CI 95% –0.3 to 31; P = 0.05)
Weymiller 2007 (in consult)	OPTION Scale	Analysis of the consultation using video-recorded consultations	1/2 used decision aid prior to consult and 1/2 used it during consult	-	Usual care	-	Greater patient participation MD 4.4 (95% CI 2.9 to 6.0) in decision aid compared to usual care

CHD: coronary heart disease; CI: confidence interval; DA: decision aid; DCS: decisional conflict scale; ICC: intraclass correlation coefficient; MD: mean difference; OPTION scale: observing patient involvement scale; RR: risk ratio; SD: standard deviation

**Table 9. Participation in decision making**

Study	Scale used	Timing	N decisionaid	Decision aid - mean	N comparison	Comparison - mean	Notes
Allen 2010	Control preferences - patients choosing active/collaborative decision making	Postintervention	291	95%	334	92%	No difference
	Control preferences did not change	Postintervention	291	92%	334	87%	No difference
	Control preferences changed to passive	Postintervention	291	3%	334	5%	No difference
	Control preferences changed to active/ collaborative	Postintervention	291	3%	334	7%	No difference
Hamann 2006	COMRADE used to measure patients' perceived involvement in decisions	Postconsultation	49	79.5 (SD 18.6) 76.8 (SD 20.9)	58	69.7 (SD 20.0) 73.5 (SD 19.3)	Increased patient involvement in decision aid group postintervention compared to usual care at baseline. At discharge there was no difference between groups
Hanson 2011	Surrogates feeling somewhat or very involved in decision making	Postintervention	-	83%	-	77%	P = 0.18
Leighl 2011	Achieved decision involvement	Postintervention	-	32%	-	35%	No difference

**Table 9. Participation in decision making** (Continued)

Loh 2007 (in consult)	Patients' perceived involvement in decision making	Postconsultation	191	26.3 pre 28.0 post	96	24.5 pre 25.5 post	Improved patient participation from baseline to post exposure to the decision aid ( $P = 0.010$ ) and in comparison to the usual care group ( $P = 0.003$ ) but there was no change in the control group for the pre-post comparison
Rubel 2010	Adapted from the Control Preferences Scale	Postintervention	-	-	-	-	The total mean scores were: 2.74 (SD 1.25) ( $N = 99$ ) pre and 2.83 (SD 1.16) ( $N = 199$ ) post, no statistically significant difference
Sheridan 2011	Patient participation: 'Any'	Immediately post	79	79%	78	51%	Absolute difference 28% (95% CI 9 to 45; $P = 0.01$ )
	'None'	Immediately post	79	21%	78	49%	Absolute difference -28% (95% CI -45 to -9)
Van Peperstraten 2010	Decision Evaluation scale (15 item questionnaire) Decision control subscale	Postconsultation	124	85	128	87.5	$P = 0.33$

DA: decision aid; SD: standard deviation.

**Table 10. Proportion undecided**

Study	Scale used	Timing	N decisionaid	Decision aid - mean	N comparison	Comparison - mean	Notes
Kasper 2008	Single item - ranging from '0 = completely undecided' to '100 = made my decision'	-	-	-	-	-	No difference
Sawka 2012	Answer "I don't know" to question "I favor taking adjuvant radioactive iodine"	Im- mediately post - treatment preference	37	10.8%	37	21.6%	-
		6.3 months (mean) post - actual decision	37	13.5%	37	8.1%	-
	Answer "I don't know" to question "I favor not taking adjuvant radioactive iodine"	Im- mediately post - treatment preference	37	43.2%	37	37.8%	-
		6.3 months (mean) post - actual decision	37	40.5%	37	51.4%	-

DA: decision aid

**Table 11. Satisfaction with the choice**

Study	Scale used	Timing	N decisionaid	Decision aid - mean	N comparison	Comparison - mean	Notes
Heller 2008	1- item; pleased with treatment choice	1 month post-surgery	62/66	-	55/67	-	P = 0.03
Legare 2012 (in consult)	Single question Likert scale to assess the quality of the decision made (0 = very low qual-	Immediately post	162	8.54 (SD 1.56)	159	8.53 (SD 1.51)	No difference; MD 0.0 (95% CI -0.4 to 0.4)

**Table 11. Satisfaction with the choice** (Continued)

	ity; 10 = very high quality)						
Leighl 2011	Satisfaction with decision scale: median (range)	1 month post-intervention	107	22 (13-25)	100	21(15-25)	No difference
Marteau 2010	Scale: ranging from 1–7 and standardized out 100	4 weeks	-	91.17 (SD 14)	-	91.33 (SD 14.50)	No difference
Schwartz 2009b	6-item	1, 6, 12 months	100	-	114	-	Overall, no difference between groups; decision aid led to significantly increased satisfaction compared to usual care among those who were undecided at randomization but not among those who had made a decision before randomization; (only graph in paper with no raw data)
Taylor 2006	Single item - "Are you satisfied with your decision about prostate cancer testing?"	1 month	80	79.7%	74	75.7%	-
Trevena 2008	Satisfaction with the decision	Immediately post	134	-	137	-	No difference (P = 0.56)



**Table 11. Satisfaction with the choice** (Continued)

Williams 2013	6-item Satisfaction with Decision Scale	Baseline	-	> 95%	-	> 95%	-
---------------	-----------------------------------------	----------	---	-------	---	-------	---

DA: decision aid.

**Table 12. Satisfaction with the decision-making process**

Study	Scale used	Timing	N decisionaid	Decision aid - mean	N comparison	Comparison - mean	Notes
<i>Satisfaction with the decision-making process</i>							<i>Satisfaction with decision-making process</i>
Hess 2012 (in consult)	Satisfaction with decision process (0 for strongly agree to 5 for strongly disagree)	-	101	-	103	-	Patients in DA group reported greater satisfaction with the DM process (strongly agree, 61% DA vs 40% usual care)
Vodermaier 2009	Satisfied with process	1 week follow-up	53	42	56	50	High satisfaction with no difference by group
<i>Satisfaction with participating in decision making</i>							<i>Satisfaction with participating in decision making</i>
Kennedy 2002	Measured satisfaction with opportunities to participate in decision making using a single item	-	-	-	-	-	Compared to usual care, women who received the decision aid followed by nurse coaching were significantly more satisfied with the opportunities to

**Table 12. Satisfaction with the decision-making process** (Continued)

							participate in decision making (OR 1.5, 95% CI 1.1 to 2.0)
<i>Satisfaction with the information provided</i>							<i>Satisfaction with information provided</i>
LeBlanc 2015 (in consult)	Amount of information was just right	Postconsultation	29	25 (86%)	37	34 (92%)	P = 0.69
	Information received was clear	Postconsultation	27	17 (63%)	36	26 (72%)	P = 0.43
	Information received was helpful	Postconsultation	28	21 (75%)	34	23 (68%)	P = 0.53
	Would recommend method to others	Postconsultation	28	24 (86%)	35	27 (77%)	P = 0.52
Laupacis 2006	Satisfaction with information received subscale 4-item (0 to 100; low to high)	Average 10 days	54	76 (15.5 SD)	56	59 (23.3 SD)	P = 0.001
Montori 2011 (in consult)	(7 point scales)	Postintervention	49	6.6	46	6.3	P = 0.798
				6		6	P = 0.296
	Participants' satisfaction with knowledge transfer			6		5.8	P = 0.624
	• Amount of information			6.1		5.8	P = 0.248
	• Clarity of information			6.4		6.2	P = 0.435
	• Helpfulness of the information						

**Table 12. Satisfaction with the decision-making process** (Continued)

	<ul style="list-style-type: none"> <li>• Would want other decisions</li> <li>• Recommend to others</li> </ul>						
	<i>Clinicians' satisfaction with knowledge transfer</i>	Postintervention	39	5.8 6.1 5.9	33	5.2 4.9 4.8	P = 0.006 P < 0.001 P < 0.001
	<ul style="list-style-type: none"> <li>• Helpfulness of the information               <ul style="list-style-type: none"> <li>• Would want other decisions</li> </ul> </li> <li>• Recommend to others</li> </ul>						
Oakley 2006	Satisfaction with information about medicines	4 months post	16	10.4 (SD 2.9)	17	10.1 (SD 2.2)	No difference
<i>Satisfaction with the clinician</i>							<i>Satisfaction with the clinician</i>
Laupacis 2006	Satisfaction with practitioner treatment during decision process sub-scale 4-item (0 to 100; low to high)	Average days	10	54	69 (25.3 SD)	56	54 (26.7 SD) P = 0.004
Miller 2005	Satisfaction with cancer information service 1-item (1 to 5; low to high)	2 weeks	-	4.37 (0.84 SD)	-	4.38 (0.86 SD)	No difference
		6 months	-	4.51 (0.75 SD)	-	4.51 (0.64 SD)	No difference

**Table 12. Satisfaction with the decision-making process** (Continued)

Vodermaier 2009	• Physician helped me understand	1 week follow-up	53	49 (92.5%)	56	53 (94.6%)	High satisfaction with no difference by group
				47		50	
				47		51	
				44		45	
				36		36	
	• Physician understood important to me						
	• Physician answered questions						
	• Satisfied with involvement						
	• Satisfied with physician's involvement						

DA: decision aid; SD: standard deviation.

**Table 13. Preparation for decision making**

Study	Scale used	Timing	N decisionaid	Decision aid - mean	N comparison	Comparison - mean	Notes
Fraenkel 2007	Preparation for Decision Making Scale	Pre-consultation	43	35 (median)	40	20.5 (median)	P < 0.001
Stacey 2014a	Preparation for Decision Making Scale item (5-point scale from: 1 not at all to 5 a great deal) 'Help recognize decision to be made'	Postintervention; pre-consultation	66	4.12 (SD 1.21)	64	3.78 (SD 1.25)	No difference